



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2021 – 02:58 PM EDT

PDB ID : 1CCI
Title : HOW FLEXIBLE ARE PROTEINS? TRAPPING OF A FLEXIBLE LOOP
Authors : Cao, Y.; Musah, R.A.; Wilcox, S.K.; Goodin, D.B.; Mcrec, D.E.
Deposited on : 1996-12-18
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	294	83% 14% ..

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3003 atoms, of which 513 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CYTOCHROME C PEROXIDASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	291	2855	1494	512	392	451	6	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	53	ILE	THR	variant	UNP P00431
A	152	GLY	ASP	variant	UNP P00431
A	202	GLY	PHE	engineered mutation	UNP P00431

- Molecule 2 is 2,3-DIMETHYLIMIDAZOLIUM ION (three-letter code: DMI) (formula: $C_5H_9N_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	N		
2	A	1	8	5	1	2	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CYTOCHROME C PEROXIDASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	107.72Å 76.63Å 51.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	5.00 – 2.40 46.60 – 2.01	Depositor EDS
% Data completeness (in resolution range)	67.1 (5.00-2.40) 73.2 (46.60-2.01)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 2.01Å)	Xtrriage
Refinement program	XTALVIEW, X-PLOR 3.1	Depositor
R, R_{free}	(Not available) , (Not available) 0.370 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	19.6	Xtrriage
Anisotropy	0.297	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 46.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.76	EDS
Total number of atoms	3003	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, DMI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.84	0/2408	1.64	46/3259 (1.4%)

There are no bond length outliers.

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	31	ARG	NE-CZ-NH2	-15.03	112.78	120.30
1	A	48	ARG	NE-CZ-NH1	11.72	126.16	120.30
1	A	166	ARG	NE-CZ-NH2	-11.50	114.55	120.30
1	A	31	ARG	NE-CZ-NH1	9.34	124.97	120.30
1	A	162	ASN	CA-C-N	-8.23	99.09	117.20
1	A	223	TRP	CD1-CG-CD2	8.19	112.85	106.30
1	A	127	ARG	CB-CG-CD	-8.16	90.38	111.60
1	A	101	TRP	CD1-CG-CD2	8.13	112.80	106.30
1	A	143	ARG	NE-CZ-NH2	-8.13	116.24	120.30
1	A	48	ARG	NE-CZ-NH2	-8.04	116.28	120.30
1	A	191	TRP	CD1-CG-CD2	8.00	112.70	106.30
1	A	72	ARG	NE-CZ-NH1	7.79	124.20	120.30
1	A	51	TRP	CD1-CG-CD2	7.75	112.50	106.30
1	A	101	TRP	CE2-CD2-CG	-7.59	101.23	107.30
1	A	211	TRP	CD1-CG-CD2	7.59	112.37	106.30
1	A	51	TRP	CE2-CD2-CG	-7.58	101.23	107.30
1	A	166	ARG	NE-CZ-NH1	7.51	124.06	120.30
1	A	191	TRP	CE2-CD2-CG	-7.42	101.37	107.30
1	A	126	TRP	CD1-CG-CD2	7.21	112.07	106.30
1	A	71	TYR	CB-CG-CD2	-7.09	116.75	121.00
1	A	223	TRP	CE2-CD2-CG	-7.07	101.65	107.30
1	A	211	TRP	CE2-CD2-CG	-7.02	101.69	107.30
1	A	127	ARG	NE-CZ-NH1	6.93	123.76	120.30
1	A	126	TRP	CE2-CD2-CG	-6.91	101.77	107.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	127	ARG	NE-CZ-NH2	-6.60	117.00	120.30
1	A	14	ARG	NE-CZ-NH2	-6.58	117.01	120.30
1	A	57	TRP	CD1-CG-CD2	6.51	111.51	106.30
1	A	57	TRP	CE2-CD2-CG	-6.37	102.20	107.30
1	A	67	TYR	CB-CG-CD1	-6.31	117.22	121.00
1	A	162	ASN	N-CA-C	6.28	127.96	111.00
1	A	195	ASN	N-CA-CB	-5.84	100.08	110.60
1	A	126	TRP	CB-CG-CD1	-5.79	119.48	127.00
1	A	126	TRP	CG-CD2-CE3	5.71	139.04	133.90
1	A	231	MET	CA-CB-CG	5.56	122.75	113.30
1	A	51	TRP	CB-CG-CD1	-5.52	119.82	127.00
1	A	166	ARG	CG-CD-NE	-5.35	100.57	111.80
1	A	223	TRP	CG-CD1-NE1	-5.34	104.76	110.10
1	A	143	ARG	NE-CZ-NH1	5.29	122.95	120.30
1	A	5	VAL	CG1-CB-CG2	-5.21	102.57	110.90
1	A	126	TRP	CG-CD1-NE1	-5.19	104.91	110.10
1	A	120	GLN	N-CA-C	5.18	125.00	111.00
1	A	162	ASN	CA-C-O	5.18	130.99	120.10
1	A	230	MET	CA-CB-CG	5.07	121.92	113.30
1	A	119	MET	CG-SD-CE	5.05	108.28	100.20
1	A	101	TRP	CG-CD1-NE1	-5.02	105.08	110.10
1	A	237	SER	CB-CA-C	-5.00	100.60	110.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2343	512	2222	10	3
2	A	7	1	9	0	0
3	A	43	0	30	0	0
4	A	97	0	0	2	0
All	All	2490	513	2261	10	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:CYS:HA	4:A:326:HOH:O	1.81	0.81
1:A:130:ARG:NH1	4:A:326:HOH:O	2.37	0.56
1:A:20:GLN:HE22	1:A:287:LYS:H	1.64	0.45
1:A:276:PHE:HA	1:A:277:PRO:HD3	1.92	0.43
1:A:48:ARG:HG2	1:A:85:LEU:HD21	2.00	0.43
1:A:119:MET:O	1:A:120:GLN:HG2	2.19	0.42
1:A:166:ARG:HH21	1:A:257:LYS:NZ	2.19	0.40
1:A:10:VAL:HG13	1:A:128:CYS:SG	2.61	0.40
1:A:40:ILE:HD11	1:A:194:ALA:HB3	2.03	0.40
1:A:77:PHE:CE1	1:A:86:GLN:HG3	2.57	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:ARG:HH21	1:A:213:LEU:O[4_477]	1.16	0.44
1:A:14:ARG:HH12	1:A:221:GLU:OE2[4_477]	1.31	0.29
1:A:14:ARG:NH2	1:A:213:LEU:O[4_477]	2.00	0.20

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	289/294 (98%)	278 (96%)	9 (3%)	2 (1%)	22 32

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	162	ASN
1	A	193	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	248/251 (99%)	239 (96%)	9 (4%)	35 54

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	LEU
1	A	35	GLU
1	A	72	ARG
1	A	74	LYS
1	A	120	GLN
1	A	171	LEU
1	A	255	GLN
1	A	260	LYS
1	A	278	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	24	ASN
1	A	78	ASN
1	A	220	ASN
1	A	240	GLN
1	A	292	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HEM	A	500	1,4	27,50,50	1.48	4 (14%)	17,82,82	1.94	5 (29%)
2	DMI	A	499	-	6,7,7	1.81	1 (16%)	3,9,9	1.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	500	1,4	-	0/6/54/54	-
2	DMI	A	499	-	-	-	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	499	DMI	C4-N3	-3.89	1.30	1.37
3	A	500	HEM	C3B-CAB	-3.89	1.40	1.47
3	A	500	HEM	C3C-CAC	-3.32	1.41	1.47
3	A	500	HEM	CBC-CAC	2.66	1.46	1.29
3	A	500	HEM	CBB-CAB	2.60	1.46	1.29

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	500	HEM	CMC-C2C-C3C	4.23	132.60	124.68
3	A	500	HEM	CMB-C2B-C3B	3.36	130.97	124.68
3	A	500	HEM	C4A-C3A-C2A	-2.63	105.16	107.00
3	A	500	HEM	C1D-C2D-C3D	-2.56	105.21	107.00
3	A	500	HEM	CMD-C2D-C3D	2.12	128.95	124.94


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

Unable to reproduce the depositor's R factor - this section is therefore empty.