

Full wwPDB NMR Structure Validation Report (i)

Nov 10, 2024 – 12:32 am GMT

PDB ID : 1DWL

Title : The Ferredoxin-Cytochrome complex using heteronuclear NMR and docking

simulation

Authors: Morelli, X.; Guerlesquin, F.; Czjzek, M.; Palma, P.N.

Deposited on : 1999-12-08

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

wwPDB-ShiftChecker : v1.2

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR,\ THEORETICAL\ MODEL$

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$egin{array}{c} { m NMR \ archive} \ (\#{ m Entries}) \end{array}$
Clashscore	210492	14027
Ramachandran outliers	207382	12486
Sidechain outliers	206894	12463

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain						
1	A	59	63%	27%	8% •				
2	В	79	53%	34%	10% •				



2 Ensemble composition and analysis (i)

This entry contains 3 models. The atoms present in the NMR models are not consistent. Some calculations may have failed as a result. All residues are included in the validation scores. Model 3 is the overall representative, medoid model (most similar to other models). The authors have identified model 2 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues								
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid mo								
1	A:1-A:59 (59)	1.36	3					
2	B:3-B:79 (77)	2.06	3					

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 1 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3



3 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 1273 atoms, of which 215 are hydrogens and 0 are deuteriums.


• Molecule 1 is a protein called FERREDOXIN I.

Mol	Chain	Residues	Atoms					Trace	
1	Λ	50	Total	С	Н	N	О	S	0
1	A	59	503	261	73	64	97	8	0

• Molecule 2 is a protein called CYTOCHROME C553.

Mol	Chain	Residues		Atoms					Trace
9	D	79	Total	С	Н	N	О	S	0
	Б	19	715	357	138	98	115	7	U


• Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms			
9	Λ	1	Total	Fe	S	
3	A	1	8	4	4	

• Molecule 4 is HEME C (three-letter code: HEC) (formula: $C_{34}H_{34}FeN_4O_4$).





Mol	Chain	Residues	Atoms					
1	D	1	Total	С	Fe	Н	N	О
4	Б	1	47	34	1	4	4	4




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.






4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


• Molecule 1: FERREDOXIN I






4.2.2 Score per residue for model 2

• Molecule 1: FERREDOXIN I



4.2.3 Score per residue for model 3 (medoid)

• Molecule 1: FERREDOXIN I





Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: MAPPING OF THE CHEMICAL SHIFT VARIATIONS.

Of the? calculated structures, 3 were deposited, based on the following criterion: COMBINATION OF NMR SHIFT VARIATION AND A SOFT DOCKING ALGORITHM.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.851

No chemical shift data was provided.



6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HEC, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		E	ond lengths	Bond angles		
MIOI	Chain	RMSZ	#Z>5	RMSZ	#Z>5	
1	A	0.97 ± 0.17	$2\pm1/434$ ($0.4\pm~0.2\%$)	1.59 ± 0.25	$7\pm 3/588$ ($1.1\pm 0.5\%$)	
2	В	1.10 ± 0.07	$5\pm1/570~(~0.8\pm~0.2\%)$	1.66 ± 0.17	$16\pm 3/751$ ($2.2\pm 0.4\%$)	
All	All	1.06	19/3014 (0.6%)	1.64	69/4023 (1.7%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	1.0 ± 0.0	3.3 ± 1.2
2	В	0.0 ± 0.0	7.0 ± 1.4
All	All	3	31

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Dag	Trme	Atoma	\mathbf{Z}	Observed (Å)	$Ideal(\mathring{A})$	Mod	dels
MIOI	Chain	Res	Type	Atoms		$\mathbf{Observed}(\mathbf{\mathring{A}})$	Ideal(A)	Worst	Total
1	A	14	SER	C-N	-15.25	0.98	1.34	1	1
2	В	64	TYR	C-N	-13.41	1.03	1.34	3	2
2	В	78	LYS	C-N	10.75	1.58	1.34	2	1
2	В	24	GLY	C-N	10.18	1.57	1.34	3	2
1	A	11	GLY	C-N	-9.03	1.13	1.34	1	1
2	В	21	ALA	C-N	8.15	1.52	1.34	2	2
2	В	65	SER	C-N	8.00	1.52	1.34	2	1
2	В	75	TYR	C-N	7.99	1.52	1.34	1	1
2	В	66	ASP	C-N	6.68	1.49	1.34	3	1
2	В	65	SER	C-O	6.67	1.36	1.23	2	1
2	В	67	GLU	N-CA	6.52	1.59	1.46	1	1
1	A	14	SER	N-CA	6.00	1.58	1.46	3	1




Continued from previous page...

Mol	Chain	Dec	Type	$\mathbf{z} = \mathbf{z} = \mathbf{z}$	$Ideal(\mathring{A})$	${f Models}$			
MIOI	Chain	nes	туре	Atoms	Z	Observed(A)	Ideal(A)	Worst	Total
1	A	28	GLY	C-N	-5.73	1.20	1.34	1	1
2	В	66	ASP	CG-OD2	5.71	1.38	1.25	1	1
2	В	75	TYR	C-O	5.53	1.33	1.23	1	1
1	A	29	GLU	C-N	-5.42	1.21	1.34	2	1

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mal	Clasia.	Das	Т	A +	7	Ob(0)	T.J 1(0)	Mod	dels
Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$	Worst	Total
1	A	14	SER	O-C-N	-17.69	94.39	122.70	1	1
1	A	11	GLY	O-C-N	-16.66	96.05	122.70	1	2
2	В	65	SER	O-C-N	16.59	149.24	122.70	2	2
2	В	24	GLY	O-C-N	-12.90	102.06	122.70	2	2
2	В	75	TYR	O-C-N	12.90	143.34	122.70	1	1
1	A	11	GLY	C-N-CA	11.96	151.59	121.70	1	1
1	A	14	SER	CA-C-N	11.69	142.91	117.20	1	1
2	В	21	ALA	C-N-CA	-11.54	92.86	121.70	2	2
1	A	11	GLY	CA-C-N	11.49	142.49	117.20	1	1
2	В	24	GLY	C-N-CA	11.46	150.35	121.70	2	1
2	В	23	MET	O-C-N	11.42	142.61	123.20	1	2
1	A	14	SER	C-N-CA	11.22	149.76	121.70	1	1
2	В	21	ALA	O-C-N	10.05	138.78	122.70	2	1
2	В	65	SER	CA-C-N	-9.69	95.87	117.20	2	1
2	В	67	GLU	O-C-N	8.78	136.75	122.70	3	1
1	A	29	GLU	O-C-N	-8.46	109.17	122.70	2	1
2	В	4	ALA	O-C-N	-8.23	109.53	122.70	3	1
2	В	78	LYS	O-C-N	8.09	135.64	122.70	2	1
2	В	24	GLY	CA-C-N	-8.07	99.44	117.20	2	1
2	В	75	TYR	CB-CG-CD2	-7.83	116.30	121.00	2	1
2	В	23	MET	CA-C-N	-7.83	100.55	116.20	1	1
2	В	66	ASP	CB-CA-C	-7.57	95.25	110.40	1	1
2	В	24	GLY	N-CA-C	-7.37	94.68	113.10	1	1
2	В	21	ALA	CA-C-N	-7.26	101.22	117.20	2	1
2	В	22	ALA	O-C-N	7.12	134.09	122.70	2	1
1	A	59	GLU	N-CA-C	-7.10	91.82	111.00	1	3
2	В	79	LEU	N-CA-C	-6.98	92.16	111.00	3	3
2	В	23	MET	C-N-CA	6.87	136.73	122.30	1	1
2	В	78	LYS	CA-C-N	-6.77	102.31	117.20	2	1
2	В	75	TYR	CA-C-N	-6.59	102.70	117.20	1	1
2	В	65	SER	C-N-CA	6.52	137.99	121.70	2	1
1	A	51	CYS	CA-CB-SG	-6.45	102.40	114.00	2	1
		•					Continued	on nert	mage





Continued from previous page...

Mal	Chain	Dec	Trme	Atoma	\mathbf{Z}	Observed(0)	Ideal(0)	Mod	dels
Mol	Chain	Res	Type	Atoms	L	$Observed(^o)$	$\operatorname{Ideal}(^{o})$	Worst	Total
2	В	14	HIS	C-N-CA	-6.37	108.93	122.30	2	1
1	A	28	GLY	C-N-CA	6.31	137.47	121.70	1	1
1	A	39	SER	N-CA-CB	-6.12	101.32	110.50	3	3
1	A	9	CYS	CA-CB-SG	-6.03	103.15	114.00	2	1
2	В	9	SER	N-CA-CB	-5.90	101.65	110.50	3	2
2	В	64	TYR	C-N-CA	5.82	136.25	121.70	1	1
2	В	67	GLU	CA-C-N	-5.81	104.42	117.20	3	1
2	В	23	MET	CG-SD-CE	5.68	109.28	100.20	2	2
2	В	66	ASP	N-CA-CB	-5.66	100.41	110.60	3	1
2	В	26	ALA	N-CA-C	-5.62	95.83	111.00	3	1
2	В	21	ALA	N-CA-C	-5.52	96.09	111.00	1	1
2	В	34	ALA	O-C-N	-5.44	114.00	122.70	3	1
2	В	64	TYR	O-C-N	-5.43	114.02	122.70	3	1
2	В	53	ARG	O-C-N	5.39	131.33	122.70	1	1
2	В	54	LYS	N-CA-CB	-5.34	100.99	110.60	1	1
2	В	53	ARG	NE-CZ-NH2	5.30	122.95	120.30	1	1
2	В	49	TYR	N-CA-C	-5.24	96.86	111.00	2	1
1	A	6	HIS	N-CA-C	-5.19	96.98	111.00	3	2
1	A	28	GLY	O-C-N	5.16	130.96	122.70	2	1
2	В	50	GLY	N-CA-C	-5.14	100.26	113.10	3	3
2	В	75	TYR	N-CA-CB	-5.03	101.55	110.60	1	1

All unique chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms	Models (Total)
1	A	26	ILE	СВ	3

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	11	GLY	Peptide	3
1	A	19	CYS	Peptide	2
2	В	64	TYR	Sidechain, Peptide, Mainchain	2
2	В	76	MET	Mainchain,Peptide	2
2	В	78	LYS	Peptide	2
1	A	51	CYS	Peptide	2
2	В	24	GLY	Mainchain,Peptide	2
1	A	4	ILE	Peptide	1
1	A	5	ASP	Peptide	1
1	A	54	GLU	Peptide	1



Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
2	В	20	LYS	Peptide	1
2	В	65	SER	Peptide	1
2	В	10	CYS	Peptide	1
2	В	17	ASP	Mainchain	1
2	В	49	TYR	Sidechain	1
2	В	75	TYR	Sidechain	1
2	В	7	TYR	Sidechain	1
2	В	21	ALA	Peptide	1
2	В	32	GLN	Peptide	1
2	В	44	TYR	Sidechain	1

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	430	73	401	14±6
2	В	564	132	563	22±6
3	A	8	0	0	1±0
4	В	43	4	30	3±1
All	All	3135	630	2978	105

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance (Å)	Mod	dels
Atom-1	Atom-2	Clash(A)	$oxed{ ext{Distance}(ext{Å}) }$	Worst	Total
2:B:23:MET:HG2	2:B:53:ARG:CG	1.57	1.20	1	1
2:B:23:MET:CG	2:B:53:ARG:HG3	1.47	1.35	1	1
1:A:13:GLU:OE1	1:A:13:GLU:O	1.13	1.65	3	1
1:A:13:GLU:OE1	1:A:13:GLU:C	1.07	1.92	3	1
2:B:62:LYS:O	2:B:65:SER:OG	1.02	1.77	2	1
2:B:65:SER:O	2:B:67:GLU:N	0.98	1.95	3	1
2:B:23:MET:SD	2:B:56:MET:HE1	0.94	2.02	1	1
1:A:13:GLU:HB2	2:B:56:MET:HG2	0.91	1.42	3	1
2:B:23:MET:CG	2:B:53:ARG:CG	0.91	2.16	1	1
1:A:14:SER:HB3	4:B:80:HEC:HBC1	0.88	1.46	1	1



Atom 1		Clash(Å)	Distance (Å)	Mod	dels
Atom-1	Atom-2	Clash(Å)	$\operatorname{Distance}(\operatorname{\AA})$	Worst	Total
2:B:23:MET:SD	2:B:56:MET:CE	0.88	2.62	1	1
1:A:12:CYS:HB2	1:A:14:SER:OG	0.88	1.67	1	1
2:B:23:MET:HG2	2:B:53:ARG:CD	0.82	2.03	1	1
1:A:13:GLU:HB3	4:B:80:HEC:HBC1	0.81	1.50	3	1
2:B:33:GLY:O	2:B:35:GLU:N	0.80	2.14	3	1
1:A:13:GLU:HB3	4:B:80:HEC:CBC	0.78	2.08	3	1
2:B:23:MET:SD	2:B:53:ARG:HB3	0.74	2.21	1	1
1:A:13:GLU:OE2	1:A:16:VAL:HB	0.74	1.81	3	1
2:B:51:GLY:N	2:B:54:LYS:HG3	0.70	2.00	1	1
2:B:23:MET:HG2	2:B:53:ARG:CB	0.69	2.16	1	1
2:B:15:GLY:O	2:B:17:ASP:N	0.68	2.25	2	1
2:B:51:GLY:O	2:B:54:LYS:HE3	0.68	1.89	1	1
2:B:23:MET:CG	2:B:53:ARG:CD	0.67	2.70	1	1
2:B:23:MET:CB	2:B:53:ARG:HG3	0.67	2.20	1	1
2:B:20:LYS:HE2	2:B:23:MET:HG2	0.66	1.67	3	1
1:A:13:GLU:OE1	1:A:16:VAL:HB	0.66	1.90	3	1
1:A:12:CYS:O	1:A:14:SER:OG	0.65	2.14	3	1
2:B:13:CYS:O	2:B:24:GLY:HA3	0.65	1.90	2	1
1:A:13:GLU:CD	1:A:16:VAL:HB	0.65	2.12	3	1
2:B:60:ALA:HB1	4:B:80:HEC:HBB2	0.64	1.67	2	2
2:B:23:MET:SD	2:B:53:ARG:CB	0.64	2.86	1	1
2:B:66:ASP:O	2:B:67:GLU:HG3	0.63	1.93	3	1
2:B:64:TYR:C	2:B:68:GLU:HB2	0.63	2.14	3	1
1:A:12:CYS:O	1:A:14:SER:N	0.63	2.32	3	2
1:A:49:ASP:HA	1:A:52:PRO:HA	0.62	1.70	2	1
1:A:18:LEU:HB3	1:A:43:CYS:HB2	0.61	1.72	1	1
1:A:13:GLU:HG3	1:A:25:MET:SD	0.61	2.35	2	1
2:B:17:ASP:HB2	2:B:75:TYR:HE2	0.61	1.55	1	1
2:B:17:ASP:CB	2:B:75:TYR:HE2	0.61	2.08	1	1
2:B:50:GLY:CA	2:B:54:LYS:HG3	0.60	2.25	1	1
2:B:15:GLY:HA2	2:B:22:ALA:N	0.60	2.12	2	1
2:B:22:ALA:HB3	4:B:80:HEC:HMD3	0.60	1.74	3	1
1:A:9:CYS:HA	3:A:60:SF4:S3	0.59	2.37	3	2
1:A:19:CYS:HB3	1:A:22:VAL:HB	0.59	1.74	3	2
2:B:41:MET:HA	2:B:44:TYR:HD2	0.56	1.58	1	1
1:A:13:GLU:OE2	1:A:16:VAL:CB	0.56	2.52	3	1
2:B:10:CYS:HB3	4:B:80:HEC:CHC	0.56	2.31	2	1
2:B:18:GLY:HA3	2:B:29:VAL:HB	0.55	1.79	3	1
2:B:50:GLY:HA2	2:B:54:LYS:HG3	0.54	1.80	1	1
2:B:14:HIS:HA	2:B:26:ALA:HB3	0.54	1.79	2	1
1:A:48:ILE:HG12	1:A:56:ILE:HB	0.54	1.79	1	1



Atom-1	Atom-2	Clash(Å)	Distance(Å)	Mod	
		` ′	,	Worst	Total
2:B:64:TYR:HA	2:B:68:GLU:OE1	0.53	2.04	3	1
1:A:48:ILE:HG23	1:A:54:GLU:HG2	0.53	1.80	2	1
2:B:75:TYR:HD2	2:B:76:MET:HG3	0.52	1.64	1	1
1:A:16:VAL:O	1:A:20:PRO:HA	0.52	2.05	1	1
1:A:16:VAL:HG11	1:A:25:MET:HB2	0.52	1.81	3	2
1:A:19:CYS:SG	1:A:43:CYS:HB2	0.52	2.45	2	1
2:B:64:TYR:O	2:B:69:LEU:HG	0.51	2.06	3	1
2:B:22:ALA:C	2:B:24:GLY:N	0.50	2.63	2	1
2:B:17:ASP:CB	2:B:75:TYR:CE2	0.49	2.95	1	1
2:B:50:GLY:C	2:B:54:LYS:HG3	0.49	2.28	1	1
2:B:23:MET:CG	2:B:53:ARG:CB	0.49	2.85	1	1
1:A:23:PHE:HA	1:A:34:VAL:HA	0.49	1.84	2	1
1:A:13:GLU:OE2	1:A:16:VAL:CG1	0.48	2.61	3	1
2:B:69:LEU:HA	2:B:72:LEU:HD12	0.48	1.83	3	1
1:A:12:CYS:C	1:A:14:SER:H	0.48	2.10	1	1
2:B:51:GLY:N	2:B:54:LYS:CG	0.47	2.76	1	1
1:A:2:ILE:HG12	1:A:58:LYS:HG2	0.47	1.85	3	1
2:B:64:TYR:O	2:B:68:GLU:HB2	0.47	2.09	3	1
2:B:51:GLY:H	2:B:54:LYS:HG3	0.47	1.69	1	1
1:A:9:CYS:HB3	1:A:30:GLU:C	0.47	2.29	3	1
2:B:23:MET:HB2	4:B:80:HEC:HMD3	0.46	1.87	1	1
2:B:22:ALA:C	2:B:24:GLY:H	0.46	2.14	2	1
2:B:22:ALA:O	2:B:24:GLY:N	0.46	2.48	2	1
2:B:15:GLY:HA3	2:B:20:LYS:O	0.46	2.10	2	1
1:A:13:GLU:C	1:A:13:GLU:CD	0.45	2.70	3	1
2:B:32:GLN:O	2:B:77:SER:HA	0.45	2.11	2	1
2:B:65:SER:C	2:B:67:GLU:N	0.45	2.68	3	1
2:B:61:VAL:HA	2:B:69:LEU:HD21	0.45	1.88	3	1
2:B:34:ALA:HB2	2:B:77:SER:HB3	0.44	1.89	3	1
1:A:9:CYS:HB3	1:A:30:GLU:CA	0.44	2.43	3	1
1:A:22:VAL:HG22	1:A:39:SER:HB2	0.43	1.90	3	1
1:A:47:ALA:HA	1:A:50:ALA:HB3	0.43	1.89	2	1
1:A:2:ILE:HD12	1:A:22:VAL:HG11	0.43	1.89	3	1
2:B:3:GLY:O	2:B:6:LEU:N	0.43	2.45	3	1
2:B:45:ALA:HA	2:B:58:THR:HG23	0.43	1.90	1	1
1:A:27:ASP:C	1:A:29:GLU:H	0.43	2.15	2	1
2:B:34:ALA:H	2:B:77:SER:HA	0.43	1.73	3	1
2:B:41:MET:HA	2:B:44:TYR:CD2	0.43	2.44	1	1
2:B:66:ASP:OD1	2:B:66:ASP:O	0.43	2.37	1	1
1:A:14:SER:HA	2:B:56:MET:SD	0.43	2.54	2	1
2:B:44:TYR:HE1	2:B:54:LYS:HB2	0.42	1.74	1	1
	l .	I.	I.		1



Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models		
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total	
2:B:72:LEU:HD13	4:B:80:HEC:HBB3	0.42	1.90	1	1	
2:B:14:HIS:HA	2:B:26:ALA:CB	0.42	2.45	2	1	
2:B:20:LYS:HE2	2:B:23:MET:CG	0.41	2.42	3	1	
1:A:19:CYS:SG	1:A:44:ALA:HB2	0.41	2.56	3	1	
2:B:7:TYR:HB2	2:B:72:LEU:HD22	0.40	1.94	2	1	
2:B:13:CYS:O	2:B:24:GLY:CA	0.40	2.66	2	1	
2:B:20:LYS:CE	2:B:23:MET:HG2	0.40	2.44	3	1	
1:A:9:CYS:HA	3:A:60:SF4:S2	0.40	2.56	2	1	

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Pe	Percentiles	
1	A	57/59 (97%)	44±1 (78±2%)	10±1 (17±2%)	3±0 (5±0%)		3	22
2	В	76/79~(96%)	57±2 (75±3%)	13±1 (18±2%)	5±2 (7±2%)		2	16
All	All	397/414 (96%)	303 (76%)	69 (17%)	25 (6%)		2	18

All 18 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	14	SER	3
1	A	29	GLU	2
2	В	65	SER	2
2	В	75	TYR	2
1	A	13	GLU	2
2	В	66	ASP	2
1	A	40	THR	1
2	В	54	LYS	1
2	В	15	GLY	1
2	В	16	ALA	1
2	В	17	ASP	1
2	В	23	MET	1
1	A	30	GLU	1



Mol	Chain	Res	Type	Models (Total)
2	В	4	ALA	1
2	В	8	LYS	1
2	В	26	ALA	1
2	В	34	ALA	1
2	В	78	LYS	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	49/49 (100%)	42±0 (86±1%)	7±0 (14±1%)	5	45	
2	В	54/55 (98%)	48±1 (90±2%)	6±1 (10±2%)	8	53	
All	All	309/312 (99%)	272 (88%)	37 (12%)	6	49	

All 21 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	1	THR	3
1	A	14	SER	3
1	A	39	SER	3
1	A	43	CYS	3
2	В	9	SER	3
2	В	19	SER	3
1	A	35	THR	2
2	В	48	SER	2
2	В	54	LYS	2
2	В	58	THR	2
1	A	12	CYS	1
1	A	26	ILE	1
2	В	25	SER	1
2	В	56	MET	1
2	В	75	TYR	1
1	A	40	THR	1
1	A	57	SER	1
2	В	7	TYR	1
2	В	44	TYR	1



Mol	Chain	Res	Type	Models (Total)
1	A	13	GLU	1
1	A	51	CYS	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mal	Tuno	Chain	Dag	Link	Bond lengths			
WIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	#Z>2	
4	HEC	В	80	2	32,50,50	1.39 ± 0.04	2±0 (6±0%)	
3	SF4	A	60	1	0,12,12	0.00 ± 0.00	-	

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.



	Mal	Trens	Chain	Dag	Timle	Bond angles Counts RMSZ #Z>2			
	IVIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	#Z>2	
	4	HEC	В	80	2	24,82,82	1.24 ± 0.14	2±1 (9±3%)	
Ī	3	SF4	A	60	1	-	-	-	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SF4	A	60	1	-	-	$0\pm0,6,5,5$
4	HEC	В	80	2	-	$0\pm0,10,54,54$	-

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mal	Chain	Dec	Tuno	Atoma	Atoms Z Observed(Å		$d(A) \mid Ideal(A)$		Models	
MIOI	Chain	nes	Туре	Atoms	L	Observed(A)	Ideal(A)	Worst	Total	
4	В	80	HEC	C3C-C2C	4.54	1.36	1.40	3	3	
4	В	80	HEC	C2B-C3B	4.34	1.36	1.40	3	3	

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Mol Chain		Tuno	Atoma	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$	Models	
MIOI	Chain	Res	Type	${f Atoms}$	L	Observed(')	Ideal(*)	Worst	Total
4	В	80	HEC	CMC-C2C-C1C	2.73	124.26	128.46	2	1
4	В	80	HEC	C4C-C3C-C2C	2.71	103.42	106.35	2	1
4	В	80	HEC	CBA-CAA-C2A	2.39	116.62	112.60	1	1
4	В	80	HEC	CMD-C2D-C1D	2.30	124.93	128.46	1	1
4	В	80	HEC	O1A-CGA-CBA	2.09	116.38	123.08	2	2
4	В	80	HEC	CMB-C2B-C1B	2.02	125.36	128.46	1	1

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be



highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

The following chains have linkage breaks:



Mol	Chain	Number of breaks
1	1-A	2
2	1-B	1
2	3-B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	В	66:ASP	С	67:GLU	N	2.97
1	A	11:GLY	С	12:CYS	N	1.13
3	В	64:TYR	С	65:SER	N	1.03
1	A	14:SER	С	15:CYS	N	0.99



7 Chemical shift validation (i)

No chemical shift data were provided

