

Full wwPDB X-ray Structure Validation Report (i)

Nov 11, 2024 – 04:47 AM EST

PDB ID	:	1ED9
Title	:	STRUCTURE OF E. COLI ALKALINE PHOSPHATASE WITHOUT THE
		INORGANIC PHOSPHATE AT 1.75A RESOLUTION
Authors	:	Stec, B.; Holtz, K.M.; Kantrowitz, E.R.
Deposited on	:	2000-01-27
Resolution	:	1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Matria	Whole archive	Similar resolution
Metric	$(\# {\rm Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
Clashscore	180529	3097 (1.76-1.76)
Ramachandran outliers	177936	3072(1.76-1.76)
Sidechain outliers	177891	3072(1.76-1.76)
RSRZ outliers	164620	2887 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	449	86%	12%	•
1	В	449	83%	15%	•



Entry composition (i) $\mathbf{2}$

There are 5 unique types of molecules in this entry. The entry contains 7240 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Δ	440	Total	С	Ν	0	\mathbf{S}	0	0	0
1 A	449	3304	2042	581	669	12	0	0	0	
1	D	440	Total	С	Ν	0	S	0	0	0
1	Б	449	3304	2042	581	669	12	0	0	0

• Molecule 1 is a protein called ALKALINE PHOSPHATASE.

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	2	Total Zn 2 2	0	0
2	В	2	Total Zn 2 2	0	0

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0

• Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O_4S).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	359	Total O 359 359	0	0
5	В	257	Total O 257 257	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: ALKALINE PHOSPHATASE



4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants	194.47Å 167.30Å 76.55Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Bosolution (Å)	15.00 - 1.75	Depositor
Resolution (A)	15.00 - 1.75	EDS
% Data completeness	92.0 (15.00-1.75)	Depositor
(in resolution range)	90.7(15.00-1.75)	EDS
R _{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$0.87 (at 1.75 \text{\AA})$	Xtriage
Refinement program	SHELXL-97	Depositor
R R.	0.196 , 0.224	Depositor
n, n_{free}	0.189 , (Not available)	DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor $(Å^2)$	17.4	Xtriage
Anisotropy	0.240	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.35 , 69.8	EDS
L-test for $twinning^2$	$ < L >=0.51, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7240	wwPDB-VP
Average B, all atoms $(Å^2)$	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.93	0/3359	1.26	9/4560~(0.2%)	
1	В	0.85	0/3359	1.22	7/4560~(0.2%)	
All	All	0.89	0/6718	1.24	16/9120~(0.2%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1
1	В	1	4
All	All	1	5

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	940	TYR	CB-CG-CD1	-9.18	115.49	121.00
1	В	584	TYR	CB-CG-CD1	7.87	125.72	121.00
1	В	732	ARG	NE-CZ-NH1	-7.76	116.42	120.30
1	А	62	ARG	NE-CZ-NH2	7.46	124.03	120.30
1	В	940	TYR	CB-CG-CD2	6.58	124.95	121.00
1	А	62	ARG	NE-CZ-NH1	-6.55	117.02	120.30
1	А	292	ARG	NE-CZ-NH1	-6.54	117.03	120.30
1	А	199	ARG	NE-CZ-NH1	6.49	123.55	120.30
1	В	504	MET	CB-CA-C	6.45	123.30	110.40
1	А	293	ASN	N-CA-C	6.06	127.37	111.00
1	А	375	GLN	N-CA-CB	5.62	120.71	110.60
1	В	504	MET	N-CA-CB	5.60	120.67	110.60



Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	А	87	TYR	CB-CG-CD1	5.30	124.18	121.00
1	В	822	GLU	OE1-CD-OE2	-5.23	117.03	123.30
1	А	284	VAL	CG1-CB-CG2	5.12	119.10	110.90
1	А	268	TRP	CB-CG-CD1	5.07	133.59	127.00

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	В	504	MET	CA

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	226	ARG	Sidechain
1	В	726	ARG	Sidechain
1	В	767	ARG	Sidechain
1	В	792	ARG	Sidechain
1	В	851	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3304	0	3248	32	0
1	В	3304	0	3245	39	0
2	А	2	0	0	0	0
2	В	2	0	0	0	0
3	А	1	0	0	0	0
3	В	1	0	0	0	0
4	А	5	0	0	0	0
4	В	5	0	0	1	0
5	А	359	0	0	8	0
5	В	257	0	0	9	0
All	All	7240	0	6493	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.



A 1 - 1		Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:A:289:ASN:O	1:A:292:ARG:HG2	1.94	0.68
1:A:73:LYS:HE3	5:A:1142:HOH:O	1.95	0.65
1:B:859:GLU:OE2	1:B:861:ASN:N	2.29	0.65
1:A:312:LYS:HE2	5:A:1137:HOH:O	1.96	0.65
1:A:137:LYS:HE3	1:A:199:ARG:O	1.98	0.64
1:A:120:LEU:O	1:A:162:HIS:HA	2.00	0.61
1:A:128:ASP:OD1	1:A:188:LYS:HE3	2.01	0.60
1:A:314:GLU:HA	1:A:314:GLU:OE2	2.00	0.60
1:B:620:LEU:O	1:B:662:HIS:HA	2.02	0.59
1:B:793:ASN:OD1	1:B:794:ASP:N	2.36	0.58
1:B:792:ARG:NH1	4:B:958:SO4:O2	2.35	0.57
1:B:907:GLU:HB2	5:B:1547:HOH:O	2.03	0.56
1:B:668:CYS:SG	1:B:677:LYS:HB2	2.45	0.56
1:B:688:LYS:HD2	5:B:1313:HOH:O	2.04	0.56
1:B:859:GLU:OE1	1:B:861:ASN:HB2	2.05	0.56
1:B:739:ASP:OD2	1:B:742:SER:OG	2.18	0.55
1:A:267:ARG:HG2	1:A:268:TRP:CD1	2.43	0.54
1:B:794:ASP:O	1:B:794:ASP:OD1	2.28	0.52
1:A:252:GLN:OE1	1:A:252:GLN:N	2.39	0.52
1:A:122:VAL:HA	1:A:127:LYS:O	2.10	0.52
1:B:793:ASN:HB3	1:B:796:VAL:HG23	1.91	0.52
1:B:515:ASP:O	1:B:521:GLY:HA3	2.10	0.52
1:B:905:SER:O	5:B:1599:HOH:O	2.20	0.50
1:B:506:VAL:HG22	5:B:1446:HOH:O	2.11	0.50
1:B:854:GLU:O	1:B:858:LYS:HG3	2.11	0.49
1:A:224:THR:OG1	1:A:227:GLU:HG3	2.11	0.49
1:B:798:THR:OG1	1:B:801:GLN:HG3	2.12	0.49
1:B:501:THR:HG23	1:B:502:PRO:HD2	1.95	0.48
1:A:365:ILE:HD13	1:A:438:LEU:HD11	1.94	0.48
1:A:228:GLN:O	1:A:232:ARG:HG3	2.15	0.47
1:B:699:ARG:NH2	1:B:754:PRO:HD3	2.29	0.47
1:B:501:THR:O	1:B:503:GLU:N	2.47	0.47
1:A:293:ASN:HB3	1:A:295:SER:OG	2.15	0.47
1:B:591:LYS:O	5:B:1444:HOH:O	2.21	0.47
1:B:672:SER:HG	1:B:713:GLU:CD	2.18	0.47
1:B:879:PRO:HA	1:B:899:VAL:HG21	1.96	0.47
1:A:31:ALA:HB1	5:A:1537:HOH:O	2.16	0.45
1:A:291:GLN:HG3	5:A:1174:HOH:O	2.17	0.45
1:A:403:GLY:HA2	5:A:1125:HOH:O	2.17	0.45
1:B:699:ARG:HD2	5:B:1245:HOH:O	2.17	0.45

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:666:ARG:NH2	5:B:1546:HOH:O	2.47	0.44
1:A:331:HIS:ND1	1:A:410:GLN:O	2.45	0.44
1:A:188:LYS:HE2	5:A:1149:HOH:O	2.17	0.44
1:B:616:TYR:CZ	1:B:619:ALA:HB2	2.52	0.44
1:B:637:LYS:NZ	1:B:699:ARG:HB3	2.34	0.43
1:B:793:ASN:ND2	1:B:795:SER:OG	2.50	0.43
1:A:264:MET:HB3	1:A:265:PRO:HD2	2.00	0.43
1:B:859:GLU:OE2	1:B:861:ASN:HB2	2.18	0.43
1:A:345:LEU:O	1:A:349:VAL:HG23	2.19	0.43
1:B:513:GLN:HG2	1:B:523:ARG:O	2.19	0.43
1:A:38:SER:OG	1:A:40:LYS:HB2	2.19	0.43
1:B:678:CYS:HB3	1:B:681:ASN:OD1	2.19	0.43
1:A:250:ASN:OD1	1:A:252:GLN:OE1	2.37	0.42
1:B:714:THR:HA	1:B:724:THR:HA	2.00	0.42
1:B:610:SER:O	1:B:631:THR:HA	2.19	0.42
1:A:128:ASP:OD1	1:A:188:LYS:CE	2.65	0.42
1:B:505:PRO:HD2	5:B:1422:HOH:O	2.19	0.41
1:B:750:ASN:OD1	1:B:752:GLN:HB2	2.20	0.41
1:A:250:ASN:CG	1:A:252:GLN:OE1	2.59	0.41
1:A:171:PRO:HD2	1:A:213:GLU:OE1	2.20	0.41
1:B:628:ASP:OD1	1:B:688:LYS:NZ	2.49	0.41
1:B:742:SER:H	1:B:742:SER:HG	1.61	0.41
1:A:294:ASP:HB2	5:A:1571:HOH:O	2.20	0.41
1:A:166:ARG:NH2	5:A:1544:HOH:O	2.53	0.41
1:B:627:LYS:HG2	1:B:628:ASP:N	2.33	0.41
1:B:767:ARG:NH2	1:B:847:GLU:OE2	2.43	0.40
1:A:149:ALA:HB2	1:A:324:ALA:CB	2.51	0.40
1:A:424:PRO:O	1:A:425:HIS:HB2	2.20	0.40
1:A:434:ASP:O	1:A:437:ASP:HB2	2.20	0.40
1:B:883:ALA:HB1	5:B:1112:HOH:O	2.20	0.40
1:A:124:ILE:HG21	1:A:124:ILE:HD13	1.88	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	447/449~(100%)	440 (98%)	7~(2%)	0	100	100
1	В	447/449~(100%)	434 (97%)	12 (3%)	1 (0%)	44	28
All	All	894/898~(100%)	874 (98%)	19~(2%)	1 (0%)	48	32

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	793	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	340/340~(100%)	327~(96%)	13 (4%)	28 10
1	В	340/340~(100%)	317~(93%)	23 (7%)	13 2
All	All	680/680~(100%)	644 (95%)	36 (5%)	19 4

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	8	GLU
1	А	40	LYS
1	А	84	TYR
1	А	92	LYS
1	А	141	LEU
1	А	177	LYS
1	А	236	LEU
1	А	261	ASP
1	А	291	GLN
1	А	294	ASP
1	A	304	ASP
1	А	330	ASP



Mol	Chain	Res	Type
1	А	408	ASP
1	В	503	GLU
1	В	504	MET
1	В	533	LEU
1	В	540	LYS
1	В	543	LYS
1	В	584	TYR
1	В	592	LYS
1	В	677	LYS
1	В	685	LYS
1	В	709	LYS
1	В	723	LYS
1	В	735	GLN
1	В	736	LEU
1	В	742	SER
1	В	753	LYS
1	В	814	GLU
1	В	815	LYS
1	В	858	LYS
1	В	872	HIS
1	В	882	LYS
1	В	909	SER
1	В	911	GLU
1	В	949	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	752	GLN
1	В	838	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Turne	Chain	Their Dec	Tink	Bond lengths			Bond angles		
	Unam	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2	
4	SO4	В	958	-	4,4,4	0.52	0	$6,\!6,\!6$	0.09	0
4	SO4	А	458	-	4,4,4	0.55	0	6,6,6	0.19	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	958	SO4	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	449/449~(100%)	0.05	12 (2%) 56 62	11, 20, 44, 115	0
1	В	449/449~(100%)	0.38	27 (6%) 29 32	13, 26, 58, 134	0
All	All	898/898~(100%)	0.21	39 (4%) 40 45	11, 23, 53, 134	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	502	PRO	7.7
1	А	1	THR	6.7
1	А	2	PRO	5.7
1	В	501	THR	5.0
1	В	507	LEU	4.5
1	В	904	ASN	4.0
1	А	293	ASN	3.4
1	В	949	LYS	3.3
1	В	793	ASN	3.2
1	А	408	ASP	3.1
1	В	731	ALA	3.0
1	В	508	GLU	2.9
1	В	506	VAL	2.8
1	В	503	GLU	2.8
1	В	753	LYS	2.7
1	В	509	ASN	2.6
1	В	909	SER	2.5
1	В	746	VAL	2.5
1	В	505	PRO	2.5
1	A	294	ASP	2.5
1	A	407	GLU	2.4
1	В	526	THR	2.4
1	В	720	TRP	2.3
1	В	736	LEU	2.3



Mol	Chain	Res	Type	RSRZ
1	В	903	GLY	2.3
1	В	592	LYS	2.2
1	В	908	ASP	2.2
1	А	404	ASN	2.2
1	А	3	GLU	2.2
1	В	540	LYS	2.2
1	А	40	LYS	2.2
1	В	740	ALA	2.1
1	А	312	LYS	2.1
1	А	252	GLN	2.1
1	В	716	THR	2.1
1	В	529	GLN	2.0
1	A	403	GLY	2.0
1	В	813	ASN	2.0
1	В	752	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
4	SO4	В	958	5/5	0.88	0.10	60,75,82,84	5
4	SO4	А	458	5/5	0.90	0.15	34,54,59,62	0
3	MG	А	462	1/1	0.97	0.04	16,16,16,16	0
2	ZN	А	450	1/1	0.99	0.07	22,22,22,22	0
3	MG	В	962	1/1	0.99	0.04	21,21,21,21	0
2	ZN	А	451	1/1	0.99	0.07	18,18,18,18	0
2	ZN	В	951	1/1	0.99	0.07	23,23,23,23	0
2	ZN	В	950	1/1	1.00	0.06	27,27,27,27	0



6.5 Other polymers (i)

There are no such residues in this entry.

