

Full wwPDB X-ray Structure Validation Report (i)

Dec 15, 2024 - 06:48 AM EST

PDB ID	:	1EUA
Title	:	SCHIFF BASE INTERMEDIATE IN KDPG ALDOLASE FROM ES-
		CHERICHIA COLI
Authors	:	Allard, J.; Grochulski, P.; Sygusch, J.
Deposited on	:	2000-04-14
Resolution	:	1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\#Entries)$	Similar resolution $(\#Entries, resolution range(Å))$
R _{free}	164625	3187 (1.96-1.96)
Clashscore	180529	3412 (1.96-1.96)
Ramachandran outliers	177936	3390 (1.96-1.96)
Sidechain outliers	177891	3390 (1.96-1.96)
RSRZ outliers	164620	3186 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain	
1	А	213	9%	22%
1	В	213	32%	16%
1	С	213	31%	24% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	А	2002	-	-	Х	-
4	PYR	А	2101	-	Х	-	-
4	PYR	В	2102	-	Х	-	-
4	PYR	С	2103	-	Х	-	-



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 5355 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Λ	012	Total	С	Ν	0	\mathbf{S}	0	0	0
1	A	213	1566	1006	260	293	7	0		
1	Р	012	Total	С	Ν	0	S	0	0	0
1	I D	213	1566	1006	260	293	7	0	0	0
1	C	012	Total	С	Ν	0	S	0	0	0
	213	1566	1006	260	293	7	0	0	0	

• Molecule 1 is a protein called KDPG ALDOLASE.

• Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
2	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0

• Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O_4S).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
3	С	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	А	1	$\begin{array}{cc} {\rm Total} & {\rm C} \\ 6 & 3 \end{array}$	O 3	0	0

Continued on next page...

BANK

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
4	С	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	193	Total O 193 193	0	0
5	В	248	Total O 248 248	0	0
5	С	175	Total O 175 175	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: KDPG ALDOLASE



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	54.41Å 85.72Å 133.19Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Bosolution (Å)	50.37 - 1.95	Depositor
Resolution (A)	50.37 - 1.95	EDS
% Data completeness	97.1 (50.37-1.95)	Depositor
(in resolution range)	96.1 (50.37 - 1.95)	EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$4.70 (at 1.95 \text{\AA})$	Xtriage
Refinement program	CNS	Depositor
B B.	0.201 , 0.243	Depositor
n, n_{free}	0.263 , 0.300	DCC
R_{free} test set	2280 reflections $(5.01%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	25.7	Xtriage
Anisotropy	0.527	Xtriage
Bulk solvent $k_{sol}(e/A^3), B_{sol}(A^2)$	0.36 , 53.1	EDS
L-test for $twinning^2$	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5355	wwPDB-VP
Average B, all atoms $(Å^2)$	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.11% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, PYR, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.31	0/1594	0.60	0/2168
1	В	0.33	0/1594	0.63	0/2168
1	С	0.30	0/1594	0.59	0/2168
All	All	0.31	0/4782	0.61	0/6504

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1566	0	1630	35	0
1	В	1566	0	1630	28	0
1	С	1566	0	1629	52	0
2	А	8	0	6	3	0
3	А	5	0	0	0	0
3	В	5	0	0	0	0
3	С	5	0	0	0	0
4	А	6	0	0	0	0
4	В	6	0	0	0	0
4	С	6	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes			
5	А	193	0	0	6	1			
5	В	248	0	0	3	2			
5	С	175	0	0	11	0			
All	All	5355	0	4895	112	2			

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (112) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:A:36:LEU:HD23	1:A:44:LEU:HD22	1.57	0.87	
1:C:187:VAL:HG12	1:C:192:LEU:HD21	1.61	0.83	
1:A:160:PRO:HG2	1:A:181:ILE:HG22	1.64	0.79	
1:C:35:ALA:HB1	5:C:2368:HOH:O	1.82	0.78	
1:A:212:LYS:O	1:A:213:LEU:HB2	1.86	0.74	
1:A:136:PRO:HG2	1:A:139:ALA:HB3	1.69	0.74	
1:A:51:GLU:H	1:A:51:GLU:CD	1.91	0.73	
1:A:189:ALA:O	1:A:193:GLU:HG3	1.87	0.72	
1:C:21:ILE:HG21	1:C:32:MET:HE1	1.70	0.71	
1:C:76:ASN:OD1	1:C:78:GLN:HG3	1.90	0.71	
1:C:136:PRO:HG2	1:C:139:ALA:HB3	1.72	0.71	
1:B:1:MET:CE	1:B:105:THR:HG22	2.20	0.71	
1:B:155:GLN:HB2	5:B:2323:HOH:O	1.92	0.68	
1:C:187:VAL:CG1	1:C:192:LEU:HD21	2.23	0.68	
1:A:165:SER:HB2	1:A:166:PRO:HD2	1.75	0.68	
1:B:1:MET:HE2	1:B:105:THR:HG22	1.76	0.67	
1:A:5:LYS:HD3	1:A:5:LYS:C	2.15	0.67	
1:C:192:LEU:HD22	1:C:192:LEU:H	1.61	0.65	
1:A:102:LYS:O	1:A:106:GLU:HG3	1.97	0.64	
1:A:190:ASP:HA	1:A:193:GLU:OE1	1.99	0.62	
1:C:165:SER:HB2	1:C:166:PRO:HD2	1.81	0.62	
1:C:201:THR:N	5:C:2368:HOH:O	2.32	0.62	
1:C:189:ALA:HB3	5:C:2311:HOH:O	1.98	0.61	
1:C:160:PRO:HG2	1:C:181:ILE:HG22	1.83	0.60	
1:A:19:PRO:HD2	1:A:43:VAL:O	2.02	0.59	
1:B:145:ALA:O	1:B:149:ILE:HD13	2.01	0.59	
1:C:1:MET:CE	1:C:105:THR:HG22	2.33	0.59	
1:C:152:PRO:HG2	1:C:153:PHE:CD1	2.38	0.59	
1:C:192:LEU:HD22	1:C:192:LEU:N	2.19	0.57	
1:B:143:VAL:O	1:B:147:GLN:HG3	2.05	0.56	



		Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:C:62:LYS:HG3	5:C:2278:HOH:O	2.06	0.55	
1:C:12:LEU:HD21	1:C:112:ILE:HD13	1.87	0.55	
1:B:188:PRO:HG2	1:B:191:ALA:HB2	1.88	0.55	
1:C:114:GLY:HA2	1:C:133:LYS:HB3	1.89	0.55	
1:C:202:LYS:HG2	1:C:206:GLU:OE2	2.07	0.54	
1:A:136:PRO:HG3	1:B:149:ILE:HD12	1.91	0.52	
1:A:114:GLY:HA2	1:A:133:LYS:O	2.09	0.52	
1:B:188:PRO:HG2	1:B:191:ALA:CB	2.39	0.52	
1:A:1:MET:HB3	5:A:2335:HOH:O	2.08	0.52	
1:B:19:PRO:HD2	1:B:43:VAL:O	2.10	0.52	
1:C:138:GLU:HB2	5:C:2258:HOH:O	2.09	0.52	
1:B:76:ASN:HB2	1:B:77:PRO:HD2	1.92	0.52	
1:C:26:LEU:O	1:C:26:LEU:HD23	2.09	0.52	
1:C:181:ILE:HD12	1:C:181:ILE:C	2.31	0.50	
1:A:25:LYS:H	2:A:2002:ACT:CH3	2.24	0.50	
1:C:188:PRO:O	1:C:191:ALA:HB3	2.11	0.50	
1:A:130:LYS:HG3	1:A:131:GLU:OE2	2.12	0.49	
1:C:192:LEU:H	1:C:192:LEU:CD2	2.23	0.49	
1:B:68:ILE:HA	1:B:89:GLN:NE2	2.28	0.49	
1:B:143:VAL:HG13	1:B:175:LEU:HD11	1.94	0.49	
1:C:114:GLY:HA2	1:C:133:LYS:O	2.13	0.49	
1:C:130:LYS:NZ	5:C:2307:HOH:O	2.46	0.49	
1:B:9:GLU:O	1:B:13:THR:HG23	2.13	0.49	
1:C:161:THR:HG21	5:C:2378:HOH:O	2.13	0.48	
1:A:213:LEU:HD22	5:A:2377:HOH:O	2.13	0.48	
1:B:25:LYS:HB2	1:B:28:HIS:HD2	1.79	0.48	
1:B:181:ILE:C	1:B:181:ILE:HD12	2.34	0.48	
1:B:80:LEU:HD21	1:B:104:ALA:HA	1.96	0.47	
1:C:169:TYR:HB2	1:C:181:ILE:HG12	1.95	0.47	
1:C:102:LYS:HD3	5:C:2340:HOH:O	2.15	0.47	
1:C:1:MET:HE1	1:C:105:THR:HG22	1.95	0.47	
1:C:19:PRO:HD2	1:C:43:VAL:O	2.15	0.47	
1:C:23:VAL:HG23	1:C:23:VAL:O	2.15	0.46	
1:C:54:VAL:HG12	5:C:2275:HOH:O	2.14	0.46	
1:A:25:LYS:H	2:A:2002:ACT:H3	1.79	0.46	
1:C:102:LYS:HB2	1:C:102:LYS:NZ	2.31	0.46	
1:A:105:THR:HG21	5:A:2315:HOH:O	2.15	0.46	
1:A:131:GLU:HG3	1:A:157:ARG:HB2	1.98	0.46	
1:B:95:GLY:HA2	1:B:114:GLY:O	2.16	0.45	
1:B:76:ASN:HB2	1:B:77:PRO:CD	2.46	0.45	
1:C:92:ILE:CG2	1:C:133:LYS:HD2	2.47	0.45	



		Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:C:41:VAL:HG23	1:C:204:ALA:HB1	1.98	0.45	
1:C:166:PRO:HD3	1:C:185:TRP:CE2	2.52	0.45	
1:B:1:MET:HG3	1:B:4:TRP:CZ3	2.52	0.45	
1:B:96:LEU:HD12	1:B:96:LEU:O	2.16	0.45	
1:B:1:MET:HE1	1:B:105:THR:HG22	1.99	0.45	
1:A:135:PHE:CD1	1:A:136:PRO:HA	2.52	0.45	
1:A:202:LYS:HD2	5:A:2249:HOH:O	2.17	0.45	
1:A:94:PRO:O	1:B:118:VAL:HG21	2.17	0.44	
1:C:114:GLY:CA	1:C:133:LYS:HB3	2.47	0.44	
1:A:39:GLY:HA2	1:A:205:ARG:HG3	2.00	0.44	
1:C:202:LYS:HE2	1:C:206:GLU:OE2	2.17	0.44	
1:C:26:LEU:HD23	1:C:26:LEU:C	2.38	0.44	
1:A:118:VAL:HG21	1:C:94:PRO:O	2.17	0.44	
1:A:181:ILE:C	1:A:181:ILE:HD12	2.38	0.44	
1:B:5:LYS:HB2	5:B:2291:HOH:O	2.18	0.44	
1:B:64:VAL:N	1:B:65:PRO:HD3	2.32	0.44	
1:A:160:PRO:HD2	1:A:178:VAL:HG13	2.00	0.43	
1:C:25:LYS:HB2	1:C:28:HIS:CD2	2.54	0.43	
1:A:132:PHE:O	1:A:158:PHE:HA	2.19	0.43	
1:C:25:LYS:HB2	1:C:28:HIS:HD2	1.84	0.43	
1:C:30:VAL:N	1:C:31:PRO:HD2	2.34	0.43	
1:C:213:LEU:HD23	1:C:213:LEU:H	1.83	0.43	
1:B:25:LYS:HB2	1:B:28:HIS:CD2	2.54	0.42	
1:A:157:ARG:HB3	1:A:179:LEU:HG	2.01	0.42	
1:A:170:ARG:NH1	1:A:210:GLY:O	2.51	0.42	
1:A:141:GLY:HA3	1:A:145:ALA:HB2	2.01	0.42	
1:C:115:ILE:C	1:C:115:ILE:HD12	2.40	0.42	
1:C:73:THR:HG22	1:C:73:THR:O	2.18	0.42	
1:A:149:ILE:HD11	5:A:2395:HOH:O	2.18	0.42	
1:B:187:VAL:HB	5:B:2278:HOH:O	2.19	0.42	
1:C:161:THR:CG2	5:C:2378:HOH:O	2.67	0.42	
1:B:5:LYS:HB2	1:B:5:LYS:HE2	1.94	0.42	
1:C:73:THR:HA	1:C:94:PRO:HD3	2.00	0.42	
1:A:115:ILE:HD12	1:A:115:ILE:C	2.40	0.41	
1:A:25:LYS:HB2	2:A:2002:ACT:H3	2.01	0.41	
1:C:159:CYS:HA	1:C:180:CYS:O	2.19	0.41	
1:A:62:LYS:HG3	5:A:2260:HOH:O	2.20	0.41	
1:C:5:LYS:HB2	5:C:2297:HOH:O	2.20	0.41	
1:B:101:LEU:O	1:B:105:THR:HG23	2.21	0.41	
1:C:166:PRO:HA	1:C:169:TYR:CD2	2.56	0.41	
1:C:213:LEU:HD23	1:C:213:LEU:N	2.36	0.40	



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:2239:HOH:O	5:B:2430:HOH:O[4_446]	2.18	0.02
5:B:2399:HOH:O	5:B:2434:HOH:O[4_446]	2.19	0.01

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	211/213~(99%)	207~(98%)	4 (2%)	0	100 100
1	В	211/213~(99%)	210 (100%)	1 (0%)	0	100 100
1	С	211/213~(99%)	205~(97%)	6 (3%)	0	100 100
All	All	633/639~(99%)	622~(98%)	11 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outlie		Perce	ntiles
1	А	163/163~(100%)	159~(98%)	4(2%)	42	34
1	В	163/163~(100%)	161~(99%)	2(1%)	67	65
1	С	163/163~(100%)	161 (99%)	2(1%)	67	65
All	All	489/489~(100%)	481 (98%)	8 (2%)	58	55



Mol	Chain	Res	Type
1	А	36	LEU
1	А	42	ARG
1	А	75	LEU
1	А	169	TYR
1	В	169	TYR
1	В	213	LEU
1	С	78	GLN
1	С	169	TYR

All (8) residues with a non-rotameric sidechain are listed below:

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	147	GLN
1	В	3	ASN
1	В	89	GLN
1	В	147	GLN
1	С	78	GLN
1	С	89	GLN
1	С	147	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and



the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain Deg Linh		Tink	Bond lengths			Bond angles		
INIOI	туре	Unain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
2	ACT	A	2001	-	3,3,3	1.11	0	3, 3, 3	0.39	0
3	SO4	А	2202	-	4,4,4	0.36	0	$6,\!6,\!6$	0.10	0
2	ACT	А	2002	-	3,3,3	1.10	0	3,3,3	0.36	0
4	PYR	А	2101	1	$5,\!5,\!5$	4.40	3 (60%)	$3,\!6,\!6$	3.10	1 (33%)
4	PYR	В	2102	1	$5,\!5,\!5$	4.47	2 (40%)	$3,\!6,\!6$	3.05	2 (66%)
4	PYR	С	2103	1	$5,\!5,\!5$	4.31	2 (40%)	$3,\!6,\!6$	3.18	2 (66%)
3	SO4	В	2201	-	4,4,4	0.34	0	6,6,6	0.09	0
3	SO4	С	2203	-	$4,\!4,\!4$	0.33	0	$6,\!6,\!6$	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PYR	А	2101	1	-	4/4/4/4	-
4	PYR	В	2102	1	-	2/4/4/4	-
4	PYR	С	2103	1	-	3/4/4/4	-

Mol	Chain	\mathbf{Res}	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
4	В	2102	PYR	O3-CA	9.27	1.44	1.23
4	А	2101	PYR	O3-CA	8.96	1.43	1.23
4	С	2103	PYR	O3-CA	8.89	1.43	1.23
4	В	2102	PYR	CA-C	-2.65	1.45	1.54
4	А	2101	PYR	CA-C	-2.60	1.45	1.54
4	С	2103	PYR	CA-C	-2.48	1.46	1.54
4	А	2101	PYR	CB-CA	-2.08	1.46	1.50

All (7) bond length outliers are listed below:

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	А	2101	PYR	O3-CA-CB	-4.82	108.97	119.77
4	С	2103	PYR	O3-CA-CB	-4.72	109.19	119.77



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	В	2102	PYR	O3-CA-CB	-4.52	109.65	119.77
4	В	2102	PYR	OXT-C-CA	2.34	120.07	113.59
4	С	2103	PYR	OXT-C-CA	2.32	120.03	113.59

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	А	2101	PYR	O-C-CA-O3
4	А	2101	PYR	OXT-C-CA-O3
4	А	2101	PYR	OXT-C-CA-CB
4	В	2102	PYR	O-C-CA-CB
4	В	2102	PYR	OXT-C-CA-CB
4	С	2103	PYR	O-C-CA-CB
4	С	2103	PYR	OXT-C-CA-CB
4	А	2101	PYR	O-C-CA-CB
4	С	2103	PYR	O-C-CA-O3

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2002	ACT	3	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

Warning: The R factor obtained from EDS is 0.2659, which does not match the depositor's R factor of 0.201. Please interpret the results in this section carefully.

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ	$>\!\!2$	$OWAB(Å^2)$	Q<0.9
1	А	213/213 (100%)	0.89	20 (9%) 15	19	17, 31, 50, 62	0
1	В	213/213~(100%)	1.64	69 (32%) 1	1	14, 23, 38, 55	0
1	С	213/213~(100%)	1.62	65 (30%) 1	1	17, 33, 51, 65	0
All	All	639/639~(100%)	1.38	154 (24%)	2 2	14, 29, 48, 65	0

All (154) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	213	LEU	4.7
1	В	26	LEU	4.6
1	С	213	LEU	4.5
1	С	105	THR	4.4
1	В	30	VAL	4.4
1	В	52	CYS	4.4
1	В	29	ALA	4.4
1	В	105	THR	4.1
1	С	211	ALA	3.9
1	В	192	LEU	3.9
1	С	8	ALA	3.9
1	В	56	ALA	3.8
1	А	27	GLU	3.7
1	В	54	VAL	3.7
1	С	23	VAL	3.7
1	С	6	THR	3.7
1	В	35	ALA	3.7
1	В	187	VAL	3.7
1	В	37	VAL	3.6
1	В	36	LEU	3.6



Mol	Chain	Res	Type	RSRZ
1	В	59	ALA	3.6
1	В	60	ILE	3.6
1	В	64	VAL	3.6
1	В	186	LEU	3.5
1	С	38	ALA	3.5
1	С	62	LYS	3.5
1	С	212	LYS	3.5
1	С	54	VAL	3.4
1	В	70	GLY	3.3
1	С	156	VAL	3.3
1	В	189	ALA	3.2
1	В	22	VAL	3.2
1	В	48	LEU	3.2
1	С	197	TYR	3.2
1	В	83	VAL	3.1
1	В	68	ILE	3.1
1	В	193	GLU	3.1
1	А	149	ILE	3.1
1	В	188	PRO	3.0
1	В	149	ILE	3.0
1	В	69	VAL	3.0
1	В	38	ALA	3.0
1	А	192	LEU	3.0
1	А	203	LEU	2.9
1	А	150	ALA	2.9
1	В	213	LEU	2.9
1	С	42	ARG	2.9
1	В	63	GLU	2.9
1	В	197	TYR	2.9
1	C	178	VAL	2.9
1	С	68	ILE	2.8
1	В	196	ASP	2.8
1	В	163	GLY	2.8
1	В	61	ALA	2.8
1	В	55	ASP	2.8
1	A	197	TYR	2.8
1	В	32	MET	2.8
1	В	25	LYS	2.8
1	С	10	SER	2.8
1	A	64	VAL	2.8
1	С	194	ALA	2.8
1	С	107	GLY	2.8



Mol	Chain	Res	Type	RSRZ
1	С	186	LEU	2.7
1	С	187	VAL	2.7
1	С	190	ASP	2.7
1	В	57	ILE	2.7
1	В	195	GLY	2.7
1	С	36	LEU	2.7
1	В	24	LYS	2.7
1	В	88	ALA	2.7
1	В	194	ALA	2.7
1	С	4	TRP	2.7
1	В	23	VAL	2.7
1	А	105	THR	2.6
1	С	37	VAL	2.6
1	С	191	ALA	2.6
1	С	203	LEU	2.6
1	В	208	VAL	2.6
1	С	41	VAL	2.6
1	С	198	ASP	2.6
1	С	192	LEU	2.6
1	В	53	ALA	2.6
1	С	53	ALA	2.6
1	А	151	GLY	2.5
1	В	40	GLY	2.5
1	С	1	MET	2.5
1	В	33	ALA	2.5
1	С	185	TRP	2.5
1	С	31	PRO	2.5
1	А	196	ASP	2.5
1	В	201	THR	2.5
1	В	81	ALA	2.5
1	С	65	PRO	2.5
1	В	4	TRP	2.5
1	С	14	THR	2.5
1	В	74	VAL	2.5
1	А	36	LEU	2.4
1	В	191	ALA	2.4
1	С	61	ALA	2.4
1	В	86	ALA	2.4
1	С	35	ALA	2.4
1	В	87	GLY	2.4
1	С	15	GLY	2.4
1	В	31	PRO	2.4



Mol	Chain	Res	Type	RSRZ
1	А	141	GLY	2.4
1	С	22	VAL	2.4
1	В	90	PHE	2.4
1	В	84	THR	2.4
1	С	3	ASN	2.3
1	В	211	ALA	2.3
1	С	209	GLU	2.3
1	В	41	VAL	2.3
1	С	208	VAL	2.3
1	А	191	ALA	2.3
1	С	210	GLY	2.3
1	С	205	ARG	2.3
1	С	24	LYS	2.3
1	С	32	MET	2.3
1	В	172	TYR	2.3
1	В	44	LEU	2.3
1	В	46	VAL	2.3
1	С	138	GLU	2.3
1	В	78	GLN	2.2
1	В	58	ARG	2.2
1	С	121	LEU	2.2
1	С	179	LEU	2.2
1	С	109	ILE	2.2
1	А	5	LYS	2.2
1	В	50	THR	2.2
1	С	30	VAL	2.2
1	В	185	TRP	2.2
1	С	158	PHE	2.2
1	С	155	GLN	2.2
1	В	47	THR	2.1
1	А	202	LYS	2.1
1	А	183	GLY	2.1
1	С	163	GLY	2.1
1	С	69	VAL	2.1
1	А	148	ALA	2.1
1	С	67	ALA	2.1
1	А	6	THR	2.1
1	С	52	CYS	2.1
1	В	198	ASP	2.1
1	С	150	ALA	2.1
1	С	189	ALA	2.1
1	В	73	THR	2.1



Mol	Chain	Res	Type	RSRZ
1	В	169	TYR	2.1
1	А	35	ALA	2.0
1	С	59	ALA	2.0
1	С	26	LEU	2.0
1	С	153	PHE	2.0
1	С	39	GLY	2.0
1	С	63	GLU	2.0
1	С	173	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
2	ACT	А	2001	4/4	0.68	0.20	41,43,44,44	0
2	ACT	А	2002	4/4	0.72	0.18	46,48,49,49	0
4	PYR	А	2101	6/6	0.76	0.15	30,33,36,36	0
3	SO4	А	2202	5/5	0.79	0.15	64,64,65,66	5
4	PYR	В	2102	6/6	0.80	0.14	25,27,31,32	0
4	PYR	С	2103	6/6	0.81	0.14	32,36,39,42	0
3	SO4	В	2201	5/5	0.84	0.18	47,49,50,50	5
3	SO4	С	2203	5/5	0.89	0.13	45,46,47,47	5

6.5 Other polymers (i)

There are no such residues in this entry.

