



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2024 – 05:23 PM EDT

PDB ID : 1GLH  
Title : CATION BINDING TO A BACILLUS (1,3-1,4)-BETA-GLUCANASE.  
GEOMETRY, AFFINITY AND EFFECT ON PROTEIN STABILITY  
Authors : Keitel, T.; Heinemann, U.  
Deposited on : 1994-11-25  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	214	 73% 19% 6%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1813 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1,3-1,4-BETA-GLUCANASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	214	1697	1090	273	329	5	0	0	0

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	115	Total	O	0	0
			115	115		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: 1,3-1,4-BETA-GLUCANASE



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.22Å 72.56Å 49.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.00	Depositor
% Data completeness (in resolution range)	(Not available) (8.00-2.00)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	PROFFT	Depositor
R, $R_{free}$	0.198 , 0.225	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1813	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.94	1/1753 (0.1%)	2.14	67/2384 (2.8%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	63	GLU	CD-OE2	5.28	1.31	1.25

All (67) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	65	ARG	NE-CZ-NH1	19.30	129.95	120.30
1	A	65	ARG	NE-CZ-NH2	-19.17	110.71	120.30
1	A	65	ARG	CD-NE-CZ	16.69	146.97	123.60
1	A	35	ARG	NE-CZ-NH1	13.52	127.06	120.30
1	A	187	THR	N-CA-CB	11.68	132.50	110.30
1	A	47	LEU	CA-CB-CG	11.55	141.86	115.30
1	A	52	THR	N-CA-CB	-11.52	88.41	110.30
1	A	35	ARG	NE-CZ-NH2	-11.39	114.61	120.30
1	A	139	ASP	CB-CG-OD1	11.04	128.24	118.30
1	A	21	ALA	CB-CA-C	10.50	125.85	110.10
1	A	190	ASP	CB-CG-OD1	10.47	127.72	118.30
1	A	77	VAL	CB-CA-C	-10.29	91.84	111.40
1	A	34	TRP	CB-CG-CD1	-9.02	115.27	127.00
1	A	207	ASP	CB-CG-OD1	8.95	126.36	118.30
1	A	118	VAL	CB-CA-C	-8.95	94.40	111.40
1	A	44	ASP	CB-CG-OD1	8.90	126.31	118.30
1	A	59	PHE	CB-CG-CD2	-8.83	114.62	120.80
1	A	18	TRP	CG-CD2-CE3	-8.63	126.13	133.90
1	A	34	TRP	CB-CG-CD2	8.51	137.66	126.60
1	A	139	ASP	CB-CG-OD2	-8.38	110.75	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	120	PHE	CB-CG-CD2	-7.43	115.60	120.80
1	A	89	SER	N-CA-CB	7.42	121.63	110.50
1	A	190	ASP	CB-CG-OD2	-7.42	111.62	118.30
1	A	164	LEU	CA-CB-CG	7.39	132.30	115.30
1	A	13	TYR	CB-CG-CD2	-7.08	116.75	121.00
1	A	149	PHE	N-CA-CB	6.97	123.14	110.60
1	A	65	ARG	O-C-N	6.87	133.68	122.70
1	A	52	THR	CA-CB-CG2	6.74	121.83	112.40
1	A	49	LEU	CB-CG-CD1	-6.59	99.80	111.00
1	A	51	LEU	C-N-CA	6.52	137.99	121.70
1	A	60	ASP	CB-CG-OD2	-6.46	112.49	118.30
1	A	118	VAL	O-C-N	6.26	132.72	122.70
1	A	109	GLU	CG-CD-OE2	-6.18	105.93	118.30
1	A	135	SER	N-CA-CB	6.12	119.67	110.50
1	A	47	LEU	N-CA-C	-5.97	94.88	111.00
1	A	187	THR	CB-CA-C	-5.94	95.56	111.60
1	A	207	ASP	CB-CG-OD2	-5.86	113.03	118.30
1	A	21	ALA	N-CA-CB	-5.82	101.95	110.10
1	A	47	LEU	N-CA-CB	5.76	121.93	110.40
1	A	13	TYR	CB-CG-CD1	5.76	124.46	121.00
1	A	18	TRP	CE2-CD2-CE3	5.76	125.61	118.70
1	A	156	ILE	N-CA-CB	5.76	124.04	110.80
1	A	65	ARG	CA-C-O	-5.75	108.02	120.10
1	A	65	ARG	CB-CA-C	-5.72	98.96	110.40
1	A	1	GLN	CB-CG-CD	5.72	126.47	111.60
1	A	196	TYR	CB-CG-CD2	-5.59	117.64	121.00
1	A	209	VAL	CA-CB-CG2	5.59	119.29	110.90
1	A	18	TRP	CA-CB-CG	5.58	124.30	113.70
1	A	209	VAL	N-CA-CB	5.50	123.61	111.50
1	A	118	VAL	CG1-CB-CG2	5.47	119.65	110.90
1	A	95	THR	N-CA-CB	5.44	120.64	110.30
1	A	193	LEU	N-CA-CB	-5.43	99.54	110.40
1	A	78	SER	N-CA-CB	5.43	118.64	110.50
1	A	194	GLY	N-CA-C	-5.41	99.57	113.10
1	A	77	VAL	CG1-CB-CG2	5.34	119.45	110.90
1	A	18	TRP	NE1-CE2-CZ2	5.33	136.26	130.40
1	A	150	ASP	CB-CG-OD1	5.31	123.08	118.30
1	A	44	ASP	CB-CG-OD2	-5.30	113.53	118.30
1	A	116	THR	CA-CB-CG2	5.28	119.80	112.40
1	A	108	ILE	N-CA-CB	5.28	122.94	110.80
1	A	1	GLN	CA-CB-CG	5.27	124.99	113.40
1	A	63	GLU	CG-CD-OE1	5.26	128.82	118.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	7	PHE	N-CA-CB	5.23	120.02	110.60
1	A	193	LEU	N-CA-C	5.21	125.06	111.00
1	A	127	VAL	CA-CB-CG2	5.20	118.70	110.90
1	A	180	MET	CG-SD-CE	5.12	108.39	100.20
1	A	88	VAL	CG1-CB-CG2	-5.05	102.83	110.90

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1697	0	1556	11	0
2	A	1	0	0	0	0
3	A	115	0	0	0	0
All	All	1813	0	1556	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:52:THR:HG22	1:A:60:ASP:HB2	1.73	0.71
1:A:32:CYS:HB3	1:A:185:ASN:HD21	1.70	0.56
1:A:77:VAL:HG13	1:A:209:VAL:HG12	1.91	0.52
1:A:114:ASP:OD1	1:A:116:THR:HB	2.10	0.52
1:A:38:ASN:HD21	1:A:52:THR:HB	1.76	0.50
1:A:69:ILE:HD12	1:A:176:PRO:HB2	1.97	0.47
1:A:87:ILE:HD13	1:A:202:LEU:HD13	1.99	0.45
1:A:175:THR:HA	1:A:176:PRO:HD3	1.81	0.44
1:A:110:PHE:CE2	1:A:118:VAL:HG13	2.54	0.42
1:A:21:ALA:HB3	1:A:63:GLU:HB3	2.03	0.41
1:A:61:CYS:HB2	1:A:183:LEU:O	2.21	0.40



There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	212/214 (99%)	201 (95%)	11 (5%)	0	100 100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	176/176 (100%)	160 (91%)	16 (9%)	9 5

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	GLN
1	A	14	ASN
1	A	17	THR
1	A	47	LEU
1	A	52	THR
1	A	58	LYS
1	A	65	ARG
1	A	77	VAL
1	A	116	THR
1	A	118	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	135	SER
1	A	164	LEU
1	A	180	MET
1	A	187	THR
1	A	191	ASP
1	A	193	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	ASN
1	A	38	ASN
1	A	43	ASN
1	A	99	HIS
1	A	102	GLN
1	A	152	GLN
1	A	185	ASN
1	A	214	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.