



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2024 – 10:47 AM EDT

PDB ID : 1HNG  
Title : CRYSTAL STRUCTURE AT 2.8 ANGSTROMS RESOLUTION OF A SOLUBLE FORM OF THE CELL ADHESION MOLECULE CD2  
Authors : Jones, E.Y.; Davis, S.J.; Williams, A.F.; Harlos, K.; Stuart, D.I.  
Deposited on : 1994-08-10  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	176	
1	B	176	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2796 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CD2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	175	1398	877	243	268	10	0	0	0
1	B	175	1398	877	243	268	10	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	90	ASN	ASP	conflict	UNP P08921
B	90	ASN	ASP	conflict	UNP P08921


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: CD2

Chain A:  72% 22% 5% ..



- Molecule 1: CD2

Chain B:  65% 26% 7% ..



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	111.40Å 111.40Å 86.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.80	Depositor
% Data completeness (in resolution range)	(Not available) (20.00-2.80)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.215 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2796	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.72	0/1422	1.47	10/1918 (0.5%)
1	B	0.72	0/1422	1.51	12/1918 (0.6%)
All	All	0.72	0/2844	1.49	22/3836 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	78	VAL	CG1-CB-CG2	7.49	122.89	110.90
1	B	78	VAL	CB-CA-C	-7.37	97.40	111.40
1	A	78	VAL	CB-CA-C	-6.80	98.48	111.40
1	B	137	ARG	NE-CZ-NH1	6.74	123.67	120.30
1	B	101	VAL	N-CA-CB	-6.54	97.12	111.50
1	A	162	ARG	NE-CZ-NH2	-6.34	117.13	120.30
1	B	140	ARG	NE-CZ-NH1	6.25	123.43	120.30
1	A	80	VAL	CG1-CB-CG2	-6.12	101.11	110.90
1	A	89	LEU	CA-CB-CG	6.12	129.38	115.30
1	A	44	ARG	CA-C-N	6.08	130.56	117.20
1	B	123	THR	N-CA-CB	-5.85	99.19	110.30
1	B	78	VAL	CA-CB-CG1	-5.84	102.15	110.90
1	A	117	CYS	CA-CB-SG	-5.80	103.56	114.00
1	A	70	ARG	NE-CZ-NH2	-5.69	117.46	120.30
1	B	96	ARG	NE-CZ-NH1	5.43	123.02	120.30
1	B	163	VAL	N-CA-CB	-5.40	99.62	111.50

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	34	ARG	NE-CZ-NH2	-5.35	117.63	120.30
1	B	89	LEU	CA-CB-CG	5.34	127.57	115.30
1	A	58	LEU	CB-CG-CD2	-5.17	102.22	111.00
1	A	50	LEU	CA-CB-CG	5.11	127.05	115.30
1	B	45	LYS	CA-CB-CG	5.08	124.57	113.40
1	A	176	GLU	N-CA-C	-5.00	97.49	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	42	PHE	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1398	0	1392	13	0
1	B	1398	0	1392	25	0
All	All	2796	0	2784	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3:SER:HA	1:B:92:ALA:O	1.79	0.83
1:B:30:VAL:HG13	1:B:42:PHE:HB3	1.71	0.71
1:B:102:SER:HB3	1:B:121:GLU:HB2	1.78	0.65
1:A:69:THR:HG22	1:A:71:ASP:H	1.63	0.64
1:B:128:LYS:HD3	1:B:138:SER:OG	2.01	0.61
1:A:110:CYS:SG	1:B:173:ASN:ND2	2.78	0.57
1:B:18:ILE:HD11	1:B:61:GLY:O	2.05	0.57
1:B:98:LEU:HD22	1:B:123:THR:HG22	1.87	0.56

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:104:PRO:HG3	1:B:159:ALA:HB2	1.85	0.56
1:A:102:SER:HB3	1:A:121:GLU:HB2	1.89	0.54
1:B:25:ASP:HA	1:B:44:ARG:NH1	2.22	0.54
1:A:80:VAL:HG12	1:A:89:LEU:HD22	1.90	0.52
1:B:69:THR:HG22	1:B:71:ASP:H	1.74	0.52
1:B:25:ASP:HA	1:B:44:ARG:HH12	1.75	0.51
1:B:28:ASP:HB2	1:B:83:THR:HA	1.93	0.51
1:A:30:VAL:HG13	1:A:42:PHE:HB3	1.96	0.48
1:A:128:LYS:HD3	1:A:138:SER:OG	2.14	0.47
1:B:10:LEU:HD13	1:B:97:ILE:HD11	1.96	0.47
1:B:109:GLU:HB3	1:B:112:ASN:OD1	2.14	0.47
1:B:101:VAL:HB	1:B:164:SER:OG	2.16	0.45
1:A:136:LEU:HG	1:A:146:TYR:CZ	2.52	0.45
1:B:154:PRO:HG3	1:B:171:VAL:HG12	1.99	0.45
1:B:73:SER:HB3	1:B:97:ILE:HG23	1.99	0.44
1:B:80:VAL:CG1	1:B:89:LEU:HB3	2.48	0.44
1:A:3:SER:HA	1:A:92:ALA:O	2.19	0.43
1:B:79:THR:HA	1:B:89:LEU:O	2.18	0.43
1:B:80:VAL:HG12	1:B:89:LEU:HD22	2.00	0.43
1:A:139:LEU:HA	1:A:139:LEU:HD23	1.77	0.43
1:B:46:MET:HG2	1:B:47:LYS:O	2.19	0.42
1:B:82:SER:CB	1:B:88:ILE:HD11	2.49	0.42
1:A:28:ASP:HB2	1:A:83:THR:HA	2.03	0.41
1:B:18:ILE:HA	1:B:19:PRO:HD3	1.80	0.41
1:A:30:VAL:CG1	1:A:42:PHE:HB3	2.51	0.40
1:A:108:TRP:O	1:B:171:VAL:CG2	2.69	0.40
1:A:131:GLN:HG2	1:A:153:ALA:HB3	2.03	0.40
1:B:63:LEU:HD23	1:B:63:LEU:HA	1.93	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	173/176 (98%)	168 (97%)	5 (3%)	0	100	100
1	B	173/176 (98%)	166 (96%)	4 (2%)	3 (2%)	9	29
All	All	346/352 (98%)	334 (96%)	9 (3%)	3 (1%)	17	46

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	44	ARG
1	B	85	GLY
1	B	20	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/157 (99%)	128 (82%)	28 (18%)	2	5
1	B	156/157 (99%)	133 (85%)	23 (15%)	3	9
All	All	312/314 (99%)	261 (84%)	51 (16%)	2	7

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	SER
1	A	20	ASN
1	A	30	VAL
1	A	31	ARG
1	A	38	LEU
1	A	48	PRO
1	A	50	LEU
1	A	57	ILE
1	A	58	LEU
1	A	63	LEU
1	A	78	VAL
1	A	79	THR
1	A	83	THR

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	89	LEU
1	A	90	ASN
1	A	95	LEU
1	A	97	ILE
1	A	101	VAL
1	A	111	SER
1	A	115	LEU
1	A	120	LEU
1	A	121	GLU
1	A	123	THR
1	A	129	LEU
1	A	136	LEU
1	A	158	LYS
1	A	167	SER
1	A	170	GLU
1	B	24	THR
1	B	30	VAL
1	B	45	LYS
1	B	57	ILE
1	B	58	LEU
1	B	63	LEU
1	B	67	ASN
1	B	78	VAL
1	B	79	THR
1	B	83	THR
1	B	86	THR
1	B	89	LEU
1	B	90	ASN
1	B	94	ASP
1	B	95	LEU
1	B	97	ILE
1	B	108	TRP
1	B	111	SER
1	B	121	GLU
1	B	123	THR
1	B	129	LEU
1	B	136	LEU
1	B	163	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	ASN
1	A	77	ASN
1	A	90	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.