

Full wwPDB X-ray Structure Validation Report (i)

Nov 3, 2024 – 08:50 PM EST

PDB ID	:	1JJH
Title	:	E2 DNA-binding Domain from Bovine Papillomavirus Type 1
Authors	:	Hegde, R.S.; Wang, A.F.; Kim, S.S.; Schapira, M.
Deposited on		
Resolution	:	2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# {\rm Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
Clashscore	180529	6282(2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chair	1	
1	А	86	66%	23%	10%
1	В	86	70%	19%	• 9%
1	С	86	52%	33%	15%



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 1896 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Λ	77	Total	С	Ν	0	S	0	0	0
	А	11	619	395	112	108	4	0	0	0
1	В	78	Total	С	Ν	0	S	0	0	0
	D	10	624	398	113	109	4	0	0	0
1	С	73	Total	С	Ν	0	S	0	0	0
	U	10	591	378	107	102	4	0	0	0

• Molecule 1 is a protein called REGULATORY PROTEIN E2.

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	325	ALA	-	cloning artifact	UNP P03122
В	325	ALA	-	cloning artifact	UNP P03122
С	325	ALA	-	cloning artifact	UNP P03122

• Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	29	TotalO2929	0	0
2	В	18	Total O 18 18	0	0
2	С	15	Total O 15 15	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: REGULATORY PROTEIN E2





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants	122.29Å 122.29Å 85.02Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 - 2.50	Depositor
% Data completeness	75.4 (50.00-2.50)	Depositor
(in resolution range)	10.4 (00.00 2.00)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	$CNS \ 0.5$	Depositor
R, R_{free}	0.225 , 0.266	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1896	wwPDB-VP
Average B, all atoms $(Å^2)$	35.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
	Ullalli	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.45	0/635	0.59	0/856
1	В	0.43	0/640	0.58	0/863
1	С	0.42	0/607	0.58	0/817
All	All	0.43	0/1882	0.58	0/2536

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	619	0	608	13	1
1	В	624	0	613	16	0
1	С	591	0	579	23	0
2	А	29	0	0	0	0
2	В	18	0	0	1	1
2	С	15	0	0	1	0
All	All	1896	0	1800	49	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:349:HIS:HD2	1:A:352:ARG:HE	1.15	0.92
1:A:349:HIS:CD2	1:A:352:ARG:HE	2.04	0.75
1:C:346:LYS:HE3	1:C:347:LYS:HE2	1.72	0.72
1:B:387:GLN:HG2	2:B:145:HOH:O	1.92	0.69
1:A:349:HIS:HD2	1:A:352:ARG:NE	1.90	0.68
1:B:344:ARG:HH11	1:B:344:ARG:HG3	1.62	0.64
1:C:328:PHE:HD2	1:C:379:THR:HG22	1.64	0.62
1:C:328:PHE:CD2	1:C:379:THR:HG22	2.35	0.62
1:C:332:SER:HB3	1:C:375:GLN:HG2	1.84	0.60
1:A:394:PRO:O	1:A:396:PRO:HD3	2.02	0.59
1:B:328:PHE:CE1	1:B:379:THR:HG22	2.38	0.57
1:C:382:SER:HB2	1:C:383:PRO:HD2	1.87	0.57
1:C:334:THR:OG1	1:C:337:GLN:HG3	2.07	0.54
1:C:341:TYR:O	1:C:345:VAL:HG23	2.08	0.53
1:A:328:PHE:CE2	1:A:379:THR:HG22	2.45	0.52
1:A:403:GLY:O	1:B:408:LEU:HD12	2.09	0.52
1:C:328:PHE:HE1	1:C:406:ALA:HB3	1.76	0.51
1:B:347:LYS:NZ	1:C:337:GLN:HE21	2.11	0.47
1:C:339:LYS:HB2	2:C:110:HOH:O	2.14	0.47
1:B:344:ARG:HG3	1:B:344:ARG:NH1	2.28	0.47
1:B:336:ASN:HB3	1:C:343:PHE:CE2	2.52	0.45
1:C:333:GLY:O	1:C:374:ALA:HB3	2.16	0.45
1:C:357:THR:CG2	1:C:377:LEU:HB3	2.46	0.45
1:C:387:GLN:O	1:C:391:LYS:HG3	2.17	0.45
1:B:350:ARG:HA	1:B:350:ARG:HE	1.82	0.45
1:C:328:PHE:CE1	1:C:406:ALA:HB3	2.51	0.45
1:A:335:ALA:HA	1:A:374:ALA:HB2	1.99	0.45
1:B:344:ARG:HH21	1:B:394:PRO:HG3	1.82	0.44
1:A:338:VAL:CG1	1:A:376:ILE:HG23	2.48	0.44
1:B:350:ARG:HA	1:B:350:ARG:NE	2.33	0.44
1:A:334:THR:OG1	1:A:337:GLN:HG3	2.16	0.44
1:B:331:ILE:HB	1:B:376:ILE:HG13	2.00	0.44
1:C:339:LYS:HD3	1:C:339:LYS:C	2.39	0.44
1:B:350:ARG:NH2	1:B:353:TYR:CE1	2.86	0.43
1:C:393:VAL:HA	1:C:394:PRO:HD3	1.88	0.43
1:C:357:THR:HG21	1:C:377:LEU:HB3	1.99	0.43
1:B:328:PHE:CD1	1:B:379:THR:HG22	2.54	0.43
1:A:386:ARG:NH2	1:A:404:PHE:O	2.52	0.43
1:C:332:SER:CB	1:C:375:GLN:HG2	2.49	0.42
1:C:374:ALA:C	1:C:375:GLN:HG3	2.40	0.42
1:A:393:VAL:HA	1:A:394:PRO:HD3	1.86	0.42
1:A:345:VAL:HG12	1:A:353:TYR:CE2	2.55	0.42

Continued on next page...



ous page		
Atom 2	Interatomic	Clash
Atom-2	distance (\AA)	overlap (Å)
1:B:402:SER:O	2.20	0.41
1:C:399:MET:SD	2.59	0.41
1:A:331:ILE:HD12	2.36	0.41

2.56

1.92

2.51

2.57

0.41

0.41

0.40

0.40

Continued from previous page...

Atom-1

1:B:329:ALA:HA

1:C:337:GLN:HB3

1:A:331:ILE:N

1:B:331:ILE:HG12

1:C:396:PRO:HA

1:C:338:VAL:CG1

1:B:360:TRP:CZ3

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:158:HOH:O	2:B:158:HOH:O[11_555]	1.70	0.50
1:A:340:CYS:SG	1:A:340:CYS:SG[9_555]	2.15	0.05

1:B:341:TYR:CE1

1:C:397:PRO:HD3

1:C:376:ILE:HG23

1:B:375:GLN:HG3

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percenti	les
1	А	73/86~(85%)	71 (97%)	2 (3%)	0	100 10)0
1	В	74/86~(86%)	72 (97%)	2(3%)	0	100 10)0
1	С	69/86~(80%)	68 (99%)	1 (1%)	0	100 10)0
All	All	216/258~(84%)	211 (98%)	5 (2%)	0	100 10)0

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	68/73~(93%)	68~(100%)	0	100 100
1	В	68/73~(93%)	65~(96%)	3~(4%)	24 47
1	С	65/73~(89%)	65 (100%)	0	100 100
All	All	201/219~(92%)	198 (98%)	3 (2%)	60 82

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	350	ARG
1	В	375	GLN
1	В	377	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	348	ASN
1	А	349	HIS
1	А	400	ASN
1	В	336	ASN
1	В	400	ASN
1	С	336	ASN
1	С	337	GLN
1	С	348	ASN
1	С	375	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.



5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

