

# Full wwPDB NMR Structure Validation Report (i)

#### Aug 20, 2022 – 08:47 AM EDT

PDB ID : 1K8N

Title: NMR structure of the XBY2 DNA duplex, an analog of CK14 containing

phosphorodithioate groups at C22 and C24

Authors: Volk, D.E.; Yang, X.; Fennewald, S.M.; King, D.J.; Bassett, S.E.; Venkitacha-

lam, S.; Herzog, N.; Luxon, B.A.; Gorenstein, D.G.

Deposited on : 2001-10-24

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

ShiftChecker : 2.29

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.29

### 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $SOLUTION\ NMR$ 

The overall completeness of chemical shifts assignment was not calculated.

There are no overall percentile quality scores available for this entry.

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain						
1	A	14	100%						
2	В	14	100%						



## 2 Ensemble composition and analysis (i)

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.



### 3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 886 atoms, of which 318 are hydrogens and 0 are deuteriums.

• Molecule 1 is a DNA chain called FIRST STRAND OF CK14 DNA DUPLEX.

Mol	Chain	Residues		Atoms					Trace
1	Λ	1.4	Total	С	Н	N	О	Р	0
1	A	14	440	135	158	54	80	13	U

• Molecule 2 is a DNA chain called SECOND STRAND OF CK14 DNA DUPLEX.

Mol	Chain	Residues			Ato	ms				Trace
9	D	1.4	Total	С	Н	N	О	Р	S	0
	D	14	446	137	160	52	80	13	4	U



### 4 Residue-property plots (i)

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: FIRST STRAND OF CK14 DNA DUPLEX

Chain A:	100%	
C1 C2 A3 G5 G5 A6 G7 C11 C12 A13 A13		
• Molecule 2: SECOND STRAN	ND OF CK14 DNA DUPLEX	
Chain B:	100%	
116 117 118 119 120 120 120 120 120 120 120 120 120 120		



#### Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: molecular dynamics.

Of the 10 calculated structures, 1 were deposited, based on the following criterion: structures with the lowest energy.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	structure solution	5
Amber	refinement	5

No chemical shift data was provided.



### 6 Model quality (i)

#### 6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: C2S

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

M	ol	Chain	Non-H	H(model)	H(added)	Clashes
A	.11	All	0	0	0	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

#### 6.3 Torsion angles (i)

#### 6.3.1 Protein backbone (i)

There are no protein molecules in this entry.

#### 6.3.2 Protein sidechains (i)

There are no protein molecules in this entry.

#### 6.3.3 RNA (i)

There are no RNA molecules in this entry.



#### 6.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Tuno	Chain	in Res	Link	Bond lengths			
IVIOI	туре	Chain		LIIIK	Counts	RMSZ	#Z>2	
2	C2S	В	22	2,1	17,20,21	0.59	0 (0%)	
2	C2S	В	24	2,1	17,20,21	0.60	0 (0%)	

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mal	Tuno	Chain	Dec	Tiple	Bond angles				
IVIOI	туре	Cham	rtes	LIIIK	Counts	RMSZ	#Z>2		
2	C2S	В	22	2,1	24,28,31	0.87	1 (4%)		
2	C2S	В	24	2,1	24,28,31	1.04	1 (4%)		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

$\mathbf{Mol}$	Type	Chain	$\operatorname{Res}$	Link	Chirals	Torsions	Rings
2	C2S	В	22	2,1	=	0,7,21,22	0,2,2,2
2	C2S	В	24	2,1	-	0,7,21,22	0,2,2,2

There are no bond-length outliers.

All angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	В	24	C2S	O2-C2-N3	2.82	117.74	122.33

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
2	В	22	C2S	O2-C2-N3	2.81	117.75	122.33

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

#### 6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 6.6 Ligand geometry (i)

There are no ligands in this entry.

#### 6.7 Other polymers (i)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 7 Chemical shift validation (i)

No chemical shift data were provided

