



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 13, 2024 – 10:57 AM EDT

PDB ID : 1O5D
Title : Dissecting and Designing Inhibitor Selectivity Determinants at the S1 site Using an Artificial Ala190 Protease (Ala190 uPA)
Authors : Katz, B.A.; Luong, C.; Ho, J.D.; Somoza, J.R.; Gjerstad, E.; Tang, J.; Williams, S.R.; Verner, E.; Mackman, R.L.; Young, W.B.; Sprengeler, P.A.; Chan, H.; Mortara, K.; Janc, J.W.; McGrath, M.E.
Deposited on : 2003-09-09
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	152	 5% 45% 15% 37%
2	H	254	 4% 78% 17% 5%
3	T	218	 5% 51% 15% 31%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CR9	H	258	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9593 atoms, of which 5038 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Coagulation factor VII.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	L	96	1395	439	668	126	149	13	0	2	0

- Molecule 2 is a protein called Coagulation factor VII.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	H	254	3965	1263	1977	353	359	13	0	3	0

- Molecule 3 is a protein called Tissue factor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
3	T	151	2406	775	1186	198	245	2	0	1	0

- Molecule 4 is 2-{5-[AMINO(IMINIO)METHYL]-6-FLUORO-1H-BENZIMIDAZOL-2-YL}-6-[(2-METHYLCYCLOHEXYL)OXY]BENZENOLATE (three-letter code: CR9) (formula: C₂₁H₂₃FN₄O₂).

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	78.54Å 69.03Å 79.12Å 90.00° 89.99° 90.00°	Depositor
Resolution (Å)	7.00 – 2.05 7.00 – 2.05	Depositor EDS
% Data completeness (in resolution range)	77.3 (7.00-2.05) 77.3 (7.00-2.05)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.90 (at 2.05Å)	Xtrriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.226 , 0.262 0.203 , 0.200	Depositor DCC
R_{free} test set	4073 reflections (10.15%)	wwPDB-VP
Wilson B-factor (Å ²)	24.7	Xtrriage
Anisotropy	0.894	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 115.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.003 for -l,k,h 0.028 for h,-k,-l 0.022 for -l,-k,-h	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9593	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CR9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	L	1.33	0/748	1.40	4/1009 (0.4%)
2	H	1.44	5/2046 (0.2%)	1.63	41/2784 (1.5%)
3	T	1.29	0/1250	1.59	20/1701 (1.2%)
All	All	1.38	5/4044 (0.1%)	1.58	65/5494 (1.2%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	52	VAL	CA-CB	5.11	1.65	1.54
2	H	141	TRP	CG-CD2	-5.05	1.35	1.43
2	H	237	TRP	CG-CD2	-5.04	1.35	1.43
2	H	190[A]	SER	CA-CB	5.00	1.60	1.52
2	H	190[B]	SER	CA-CB	5.00	1.60	1.52

All (65) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	29	TRP	CD1-NE1-CE2	10.55	118.50	109.00
2	H	61	TRP	CD1-NE1-CE2	10.26	118.23	109.00
3	T	45	TRP	CD1-NE1-CE2	9.90	117.91	109.00
2	H	207	TRP	CD1-NE1-CE2	9.61	117.65	109.00
3	T	25	TRP	CD1-NE1-CE2	9.53	117.58	109.00
2	H	215	TRP	CD1-NE1-CE2	9.49	117.54	109.00
2	H	51	TRP	CD1-NE1-CE2	9.39	117.45	109.00
2	H	237	TRP	CD1-NE1-CE2	9.17	117.25	109.00
3	T	74	ARG	NE-CZ-NH2	-9.11	115.75	120.30
3	T	14	TRP	CD1-NE1-CE2	8.96	117.07	109.00
2	H	29	TRP	NE1-CE2-CZ2	8.81	140.09	130.40
2	H	141	TRP	CD1-NE1-CE2	8.79	116.91	109.00
1	L	123	ASP	CB-CG-OD2	-8.68	110.48	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	207	TRP	NE1-CE2-CZ2	8.51	139.76	130.40
2	H	237	TRP	NE1-CE2-CZ2	8.28	139.51	130.40
2	H	29	TRP	CG-CD1-NE1	-8.14	101.96	110.10
2	H	61	TRP	CG-CD1-NE1	-7.99	102.11	110.10
2	H	199	HIS	N-CA-C	-7.74	90.11	111.00
3	T	25	TRP	CG-CD1-NE1	-7.55	102.55	110.10
3	T	45	TRP	NE1-CE2-CZ2	7.50	138.65	130.40
3	T	14	TRP	NE1-CE2-CZ2	7.48	138.63	130.40
2	H	207	TRP	CG-CD1-NE1	-7.45	102.65	110.10
2	H	141	TRP	NE1-CE2-CZ2	7.41	138.55	130.40
2	H	243	ARG	NE-CZ-NH2	-7.33	116.64	120.30
2	H	129(G)	VAL	N-CA-C	-7.27	91.37	111.00
3	T	45	TRP	CG-CD1-NE1	-7.25	102.86	110.10
2	H	237	TRP	CG-CD1-NE1	-7.18	102.92	110.10
2	H	215	TRP	CG-CD1-NE1	-7.06	103.04	110.10
2	H	51	TRP	CG-CD1-NE1	-7.04	103.06	110.10
3	T	157	TYR	N-CA-C	-6.99	92.12	111.00
2	H	29	TRP	NE1-CE2-CD2	-6.98	100.32	107.30
2	H	134	ARG	NE-CZ-NH2	-6.91	116.84	120.30
2	H	253	ARG	NE-CZ-NH2	-6.91	116.84	120.30
3	T	14	TRP	CG-CD1-NE1	-6.79	103.31	110.10
2	H	61	TRP	NE1-CE2-CZ2	6.71	137.78	130.40
1	L	104	ASP	CB-CG-OD2	-6.68	112.29	118.30
2	H	141	TRP	CG-CD1-NE1	-6.66	103.44	110.10
2	H	237	TRP	NE1-CE2-CD2	-6.57	100.73	107.30
2	H	215	TRP	NE1-CE2-CZ2	6.47	137.51	130.40
3	T	40	THR	N-CA-C	-6.47	93.54	111.00
2	H	207	TRP	NE1-CE2-CD2	-6.45	100.85	107.30
2	H	147	ARG	NE-CZ-NH2	-6.41	117.09	120.30
3	T	25	TRP	NE1-CE2-CZ2	6.36	137.39	130.40
2	H	61	TRP	NE1-CE2-CD2	-6.16	101.14	107.30
2	H	141	TRP	NE1-CE2-CD2	-6.09	101.21	107.30
3	T	196	ARG	NE-CZ-NH2	-6.06	117.27	120.30
2	H	215	TRP	NE1-CE2-CD2	-6.02	101.28	107.30
3	T	136	ARG	NE-CZ-NH2	-5.91	117.35	120.30
3	T	28	LYS	N-CA-C	-5.79	95.36	111.00
3	T	45	TRP	NE1-CE2-CD2	-5.79	101.51	107.30
3	T	14	TRP	NE1-CE2-CD2	-5.78	101.53	107.30
2	H	129(B)	ARG	NE-CZ-NH2	-5.70	117.45	120.30
2	H	51	TRP	NE1-CE2-CD2	-5.67	101.63	107.30
2	H	170(D)	LYS	N-CA-C	5.66	126.28	111.00
2	H	29	TRP	CG-CD2-CE3	-5.66	128.81	133.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	T	156	TYR	N-CA-C	-5.62	95.83	111.00
2	H	51	TRP	NE1-CE2-CZ2	5.53	136.48	130.40
1	L	113	ARG	NE-CZ-NH2	-5.43	117.58	120.30
3	T	25	TRP	NE1-CE2-CD2	-5.43	101.87	107.30
2	H	134	ARG	CA-CB-CG	-5.41	101.49	113.40
1	L	101	TYR	CB-CG-CD1	5.35	124.21	121.00
2	H	170(F)	GLY	N-CA-C	-5.24	100.00	113.10
2	H	79	ASP	N-CA-C	5.14	124.89	111.00
3	T	86	THR	N-CA-CB	-5.13	100.56	110.30
2	H	62	ARG	NE-CZ-NH2	-5.03	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	727	668	663	15	1
2	H	1988	1977	1967	26	0
3	T	1220	1186	1182	18	0
4	H	28	23	23	2	0
5	H	295	590	0	1	0
5	L	110	220	0	0	0
5	T	187	374	0	0	1
All	All	4555	5038	3835	56	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:195:SER:OG	4:H:258:CR9:O6'	2.07	0.71
1:L:61:CYS:SG	1:L:68:TYR:HB2	2.36	0.65
1:L:71:PHE:CE2	3:T:131:ARG:HD3	2.31	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:236:GLU:H	2:H:236:GLU:CD	2.02	0.62
2:H:143:GLN:HE22	2:H:147:ARG:H	1.50	0.59
1:L:65:LEU:HD12	1:L:65:LEU:N	2.17	0.59
2:H:136:SER:CB	2:H:199:HIS:CE1	2.90	0.55
2:H:136:SER:HB2	2:H:199:HIS:CE1	2.44	0.53
1:L:49:GLN:O	1:L:51:ALA:N	2.42	0.52
1:L:65:LEU:N	1:L:65:LEU:CD1	2.74	0.51
3:T:155:LEU:O	3:T:188:SER:N	2.43	0.51
2:H:228:TYR:N	2:H:228:TYR:CD1	2.78	0.51
3:T:87:GLY:O	3:T:89:ALA:N	2.43	0.51
3:T:51:TYR:CE1	3:T:83:VAL:HG13	2.46	0.49
2:H:50:ILE:HG13	2:H:51:TRP:CD1	2.48	0.48
2:H:135:PHE:HB3	2:H:159:ASN:ND2	2.28	0.48
2:H:124:PRO:O	2:H:235:ILE:HD13	2.13	0.47
1:L:121[B]:LEU:HD11	1:L:128:THR:HB	1.95	0.47
3:T:194:PRO:HA	3:T:200:ARG:NH1	2.28	0.47
4:H:258:CR9:O6'	4:H:258:CR9:N3	2.46	0.47
1:L:49:GLN:N	1:L:49:GLN:OE1	2.49	0.46
2:H:180:MET:HE1	5:H:440:HOH:O	2.15	0.46
3:T:108:LEU:HB2	3:T:202:SER:HB3	1.96	0.46
1:L:62:LYS:O	1:L:68:TYR:HA	2.16	0.46
2:H:45:THR:OG1	2:H:198:PRO:HB3	2.17	0.45
1:L:71:PHE:CE2	3:T:131:ARG:CD	3.00	0.44
2:H:143:GLN:HE22	2:H:147:ARG:N	2.14	0.44
2:H:27:CYS:N	2:H:28:PRO:CD	2.81	0.43
2:H:245:GLU:O	2:H:247:ARG:CD	2.66	0.43
3:T:45:TRP:CE2	3:T:74:ARG:NH2	2.86	0.43
3:T:150:ASP:O	3:T:193:ILE:HA	2.19	0.43
1:L:65:LEU:HD11	3:T:203:THR:HG21	2.01	0.43
2:H:91:PRO:O	2:H:253:ARG:NH2	2.52	0.43
3:T:51:TYR:CD1	3:T:83:VAL:HG22	2.53	0.42
3:T:154:THR:O	3:T:189:VAL:HA	2.19	0.42
2:H:92:SER:OG	2:H:255:PRO:HA	2.19	0.42
1:L:90:ILE:HD12	1:L:92:VAL:HG12	2.02	0.42
2:H:161:PRO:O	2:H:183:ALA:HA	2.20	0.41
3:T:135:ARG:HA	3:T:139:THR:O	2.20	0.41
2:H:127:THR:O	2:H:129(B):ARG:HB3	2.21	0.41
1:L:61:CYS:SG	1:L:68:TYR:CB	3.06	0.41
2:H:245:GLU:O	2:H:247:ARG:HD3	2.19	0.41
3:T:144:ARG:HD2	3:T:148:GLY:O	2.20	0.41
1:L:129:PRO:HB3	1:L:134:PRO:HB3	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:76:HIS:ND1	2:H:77:ASP:O	2.54	0.41
2:H:121:LEU:HD11	2:H:207:TRP:HB3	2.03	0.41
2:H:166:GLN:HB2	3:T:94:TYR:OH	2.21	0.41
2:H:239:GLN:O	2:H:243:ARG:HG3	2.21	0.41
2:H:245:GLU:CD	2:H:245:GLU:N	2.74	0.41
1:L:65:LEU:CD1	1:L:65:LEU:H	2.34	0.40
3:T:51:TYR:CD1	3:T:83:VAL:CG2	3.05	0.40
3:T:142:SER:O	3:T:145:ASP:N	2.54	0.40
2:H:215:TRP:CE2	2:H:227:VAL:HG11	2.56	0.40
3:T:86:THR:OG1	3:T:90:GLY:C	2.60	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:118:TYR:O	5:T:303:HOH:H1[2_646]	1.53	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	96/152 (63%)	81 (84%)	9 (9%)	6 (6%)	1	0
2	H	255/254 (100%)	228 (89%)	23 (9%)	4 (2%)	9	2
3	T	146/218 (67%)	125 (86%)	17 (12%)	4 (3%)	5	0
All	All	497/624 (80%)	434 (87%)	49 (10%)	14 (3%)	4	0

All (14) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	50	CYS
2	H	79	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	T	88	SER
3	T	148	GLY
1	L	57	ASN
2	H	49	THR
3	T	156	TYR
1	L	66	GLN
1	L	51	ALA
2	H	170(I)	PRO
3	T	137	ASN
1	L	81	CYS
2	H	255	PRO
1	L	74	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	85/132 (64%)	78 (92%)	7 (8%)	11	5
2	H	219/216 (101%)	210 (96%)	9 (4%)	30	23
3	T	140/199 (70%)	134 (96%)	6 (4%)	29	22
All	All	444/547 (81%)	422 (95%)	22 (5%)	24	16

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	49	GLN
1	L	65	LEU
1	L	83	THR
1	L	88	GLN
1	L	92	VAL
1	L	103	SER
1	L	110	ARG
2	H	68	LEU
2	H	75	GLU
2	H	84	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	92	SER
2	H	100	ASN
2	H	159	ASN
2	H	233	GLN
2	H	236	GLU
2	H	255	PRO
3	T	83	VAL
3	T	85	SER
3	T	86	THR
3	T	131	ARG
3	T	156	TYR
3	T	195	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	64	GLN
2	H	100	ASN
2	H	143	GLN
2	H	159	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	CR9	H	258	-	30,31,31	2.17	11 (36%)	32,45,45	1.38	4 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CR9	H	258	-	2/2/3/3	1/7/23/23	0/4/4/4

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	258	CR9	C1-C7	-5.89	1.39	1.48
4	H	258	CR9	C1-C2	4.05	1.44	1.38
4	H	258	CR9	C4-N3	-3.15	1.28	1.38
4	H	258	CR9	C3-C4	-3.13	1.37	1.41
4	H	258	CR9	C6-C5	-3.04	1.37	1.41
4	H	258	CR9	C6-C1	2.87	1.41	1.37
4	H	258	CR9	C4B-C5B	2.71	1.60	1.53
4	H	258	CR9	C3-C2	2.67	1.39	1.35
4	H	258	CR9	C5B-C6B	2.66	1.57	1.52
4	H	258	CR9	C8-N3	-2.59	1.30	1.34
4	H	258	CR9	C5-C4	-2.18	1.35	1.42

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	258	CR9	C1A-C1B-C6B	-3.44	108.11	112.65
4	H	258	CR9	F2-C2-C3	-3.31	116.10	120.47
4	H	258	CR9	C2-C3-C4	-2.97	115.95	118.84
4	H	258	CR9	C6-C5-N4	-2.13	124.73	130.83

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	H	258	CR9	C6B
4	H	258	CR9	C1B

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	H	258	CR9	C6-C1-C7-N1

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	258	CR9	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	96/152 (63%)	0.00	8 (8%) 11 11	12, 34, 59, 66	17 (17%)
2	H	254/254 (100%)	-0.42	11 (4%) 35 38	8, 26, 64, 92	21 (8%)
3	T	151/218 (69%)	0.13	10 (6%) 18 19	12, 37, 76, 92	21 (13%)
All	All	501/624 (80%)	-0.17	29 (5%) 23 25	8, 30, 69, 92	59 (11%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	66	GLN	11.9
3	T	138	ASN	10.6
3	T	137	ASN	9.1
2	H	170(G)	ASP	6.7
2	H	170(H)	SER	5.8
3	T	157	TYR	5.7
1	L	65	LEU	5.6
3	T	156	TYR	5.5
2	H	170(F)	GLY	4.7
3	T	188	SER	4.5
2	H	170(E)	VAL	4.0
1	L	48	ASP	3.9
3	T	109	GLY	3.8
3	T	88	SER	3.4
2	H	60(C)	LYS	3.4
1	L	50	CYS	3.2
1	L	49	GLN	3.2
2	H	248	PRO	3.0
2	H	61	TRP	2.8
2	H	170(D)	LYS	2.8
1	L	67	SER	2.7
3	T	155	LEU	2.4
2	H	60(B)	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	L	53	SER	2.4
2	H	247	ARG	2.3
2	H	63	ASN	2.3
3	T	205	SER	2.2
1	L	51	ALA	2.1
3	T	129	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CR9	H	258	28/28	0.95	0.11	22,44,47,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.