



Full wwPDB X-ray Structure Validation Report

Oct 23, 2021 – 09:49 AM EDT

PDB ID : 1PRX
Title : HORF6 A NOVEL HUMAN PEROXIDASE ENZYME
Authors : Choi, H.-J.; Kang, S.W.; Yang, C.-H.; Rhee, S.G.; Ryu, S.-E.
Deposited on : 1998-04-03
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	224	
1	B	224	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3574 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HORF6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	220	1743	1125	292	322	4	0	0	0
1	B	218	1719	1111	288	316	4	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	CSO	CYS	modified residue	UNP P30041
A	91	SER	CYS	engineered mutation	UNP P30041
B	47	CSO	CYS	modified residue	UNP P30041
B	91	SER	CYS	engineered mutation	UNP P30041

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	60	Total	O	0	0
			60	60		
2	B	52	Total	O	0	0
			52	52		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: HORF6

Chain A: 



- Molecule 1: HORF6

Chain B: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	47.85Å 75.17Å 63.30Å 90.00° 110.21° 90.00°	Depositor
Resolution (Å)	8.00 – 2.00	Depositor
% Data completeness (in resolution range)	93.7 (8.00-2.00)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.191 , 0.259	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3574	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/1777	0.73	2/2412 (0.1%)
1	B	0.44	0/1753	0.74	2/2380 (0.1%)
All	All	0.44	0/3530	0.73	4/4792 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	145	LEU	CA-CB-CG	9.20	136.47	115.30
1	B	145	LEU	CA-CB-CG	7.04	131.50	115.30
1	A	179	VAL	CB-CA-C	-6.02	99.95	111.40
1	B	179	VAL	CB-CA-C	-5.33	101.26	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1743	0	1775	32	0
1	B	1719	0	1749	41	0
2	A	60	0	0	0	0
2	B	52	0	0	6	0
All	All	3574	0	3524	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:ILE:HD11	1:A:121:GLU:HB3	1.64	0.80
1:A:50:GLU:HG2	1:B:179:VAL:HG13	1.66	0.78
1:B:44:THR:HG23	1:B:47:CSO:HB3	1.68	0.73
1:B:107:ASN:HB2	2:B:2050:HOH:O	1.88	0.72
1:B:168:GLN:O	1:B:172:GLU:HG2	1.93	0.68
1:B:211:LEU:HD21	1:B:218:LEU:HD12	1.77	0.64
1:B:174:ARG:HD2	2:B:2043:HOH:O	2.00	0.61
1:B:44:THR:CG2	1:B:47:CSO:HB3	2.32	0.60
1:A:149:TYR:HA	1:B:145:LEU:HD22	1.85	0.58
1:A:179:VAL:HG22	1:B:49:THR:HB	1.86	0.58
1:A:50:GLU:CG	1:B:179:VAL:HG13	2.33	0.57
1:B:162:ARG:NH2	2:B:2001:HOH:O	2.38	0.56
1:A:179:VAL:HG13	1:B:50:GLU:HG2	1.87	0.55
1:A:112:ILE:CD1	1:A:121:GLU:HB3	2.35	0.54
1:A:118:ASP:O	1:A:120:ALA:N	2.41	0.53
1:B:5:LEU:HD12	1:B:5:LEU:N	2.23	0.53
1:B:22:ARG:NE	1:B:22:ARG:H	2.07	0.51
1:B:95:THR:HG22	1:B:96:GLU:N	2.24	0.51
1:B:108:ARG:O	1:B:112:ILE:HG13	2.10	0.51
1:A:69:ILE:HG13	1:A:101:PRO:HG2	1.92	0.51
1:A:145:LEU:HD22	1:B:149:TYR:HA	1.92	0.51
1:A:179:VAL:CG1	1:B:50:GLU:HA	2.40	0.50
1:A:33:TRP:HB2	1:A:66:VAL:HG22	1.93	0.49
1:A:50:GLU:HA	1:B:179:VAL:HG11	1.93	0.49
1:B:41:ARG:HD3	1:B:73:ILE:O	2.12	0.49
1:B:176:ALA:O	1:B:187:VAL:HG22	2.12	0.49
1:A:181:TRP:CD1	1:A:185:ASP:HB2	2.48	0.48
1:B:133:VAL:HG22	1:B:148:LEU:HG	1.95	0.48
1:B:188:MET:CE	1:B:218:LEU:HD23	2.43	0.47
1:A:125:LYS:C	1:A:127:MET:H	2.18	0.47
1:A:179:VAL:HG11	1:B:50:GLU:HA	1.96	0.47
1:A:179:VAL:HG13	1:B:50:GLU:CG	2.44	0.47
1:A:37:PHE:HE2	1:A:68:LEU:HG	1.81	0.46
1:A:72:SER:HB3	1:A:79:HIS:CE1	2.51	0.46
1:B:37:PHE:HE1	1:B:68:LEU:HG	1.80	0.46
1:A:73:ILE:HD12	1:A:128:PRO:CB	2.45	0.46
1:A:50:GLU:HA	1:B:179:VAL:CG1	2.46	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:15:GLU:HB2	1:A:24:ARG:HH11	1.81	0.46
1:A:126:GLY:O	1:A:127:MET:HB2	2.15	0.46
1:A:125:LYS:O	1:A:127:MET:N	2.50	0.45
1:A:179:VAL:HG22	1:B:49:THR:CB	2.46	0.45
1:A:153:THR:HG21	1:B:163:VAL:HG22	1.99	0.43
1:A:133:VAL:HG22	1:A:148:LEU:HG	2.00	0.43
1:A:181:TRP:CZ2	1:A:183:ASP:HA	2.53	0.43
1:B:26:HIS:HB3	2:B:2059:HOH:O	2.19	0.43
1:B:97:LYS:HG3	2:B:2047:HOH:O	2.18	0.43
1:B:145:LEU:HD13	1:B:146:SER:N	2.34	0.43
1:B:209:LYS:HB3	1:B:218:LEU:HB3	2.02	0.42
1:B:51:LEU:HD12	1:B:51:LEU:HA	1.86	0.42
1:B:58:ALA:HB3	1:B:59:PRO:HD3	2.01	0.42
1:B:11:ALA:HB2	1:B:137:PHE:CE1	2.55	0.42
1:B:56:LYS:HE3	1:B:56:LYS:HA	2.02	0.42
1:A:188:MET:CE	1:A:218:LEU:HD22	2.50	0.41
1:A:13:ASN:OD1	1:A:24:ARG:HG3	2.21	0.41
1:B:45:PRO:HG2	2:B:2112:HOH:O	2.20	0.41
1:A:45:PRO:O	1:A:49:THR:HG23	2.21	0.41
1:A:149:TYR:CD1	1:B:145:LEU:HD21	2.56	0.41
1:B:76:VAL:CG2	1:B:104:ASP:HB2	2.51	0.41
1:B:95:THR:HG22	1:B:96:GLU:H	1.85	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/224 (97%)	201 (93%)	13 (6%)	3 (1%)	11	5
1	B	213/224 (95%)	206 (97%)	6 (3%)	1 (0%)	29	23
All	All	430/448 (96%)	407 (95%)	19 (4%)	4 (1%)	17	11

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	119	PRO
1	B	45	PRO
1	A	127	MET
1	A	126	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/193 (99%)	175 (92%)	16 (8%)	11	7
1	B	188/193 (97%)	172 (92%)	16 (8%)	10	6
All	All	379/386 (98%)	347 (92%)	32 (8%)	11	7

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	LEU
1	A	68	LEU
1	A	92	GLU
1	A	97	LYS
1	A	98	LEU
1	A	113	LEU
1	A	143	LEU
1	A	145	LEU
1	A	148	LEU
1	A	161	LEU
1	A	167	LEU
1	A	169	LEU
1	A	179	VAL
1	A	189	VAL
1	A	201	LEU
1	A	211	LEU
1	B	5	LEU
1	B	22	ARG
1	B	51	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	56	LYS
1	B	57	LEU
1	B	68	LEU
1	B	92	GLU
1	B	106	ARG
1	B	110	LEU
1	B	121	GLU
1	B	145	LEU
1	B	148	LEU
1	B	167	LEU
1	B	179	VAL
1	B	201	LEU
1	B	211	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	ASN
1	A	87	ASN
1	B	87	ASN
1	B	90	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSO	B	47	1	3,6,7	0.65	0	0,6,8	-	-
1	CSO	A	47	1	3,6,7	0.60	0	0,6,8	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	B	47	1	-	0/1/5/7	-
1	CSO	A	47	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	47	CSO	2	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.