

Full wwPDB NMR Structure Validation Report (i)

Dec 24, 2024 – 11:28 AM EST

PDB ID	:	1QMW
BMRB ID	:	4503
Title	:	Solution structure of alpha-conotoxin SI
Authors	:	Benie, A.J.; Whitford, D.; Hargittai, B.; Barany, G.; Janes, R.W.
Deposited on	:	1999-10-08

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
BMRB Restraints Analysis	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 78%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f NMR} \ {f archive} \ (\#{f Entries})$
Clashscore	210492	14027
Ramachandran outliers	207382	12486
Sidechain outliers	206894	12463

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain				
1	А	14	43%	21%	7%	29%	



2 Ensemble composition and analysis (i)

This entry contains 36 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues					
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model					
1	A:4-A:13 (10)	0.13	1		

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 6 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 8, 9, 10, 11, 13, 14, 15, 16, 18, 26
2	7, 12, 17, 19, 20, 23
3	24, 25, 27, 32
4	29, 30, 31
5	22, 28
6	33, 36
Single-model clusters	21; 34; 35



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 177 atoms, of which 86 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called ALPHA-CONOTOXIN SI.

Mol	Chain	Residues		I	Aton	ıs			Trace
1	٨	1.4	Total	С	Η	Ν	Ο	S	1
	A	14	177	55	86	16	16	4	1




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: ALPHA-CONOTOXIN SI



4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1 (medoid)

• Molecule 1: ALPHA-CONOTOXIN SI



4.2.2 Score per residue for model 2


• Molecule 1: ALPHA-CONOTOXIN SI





4.2.3 Score per residue for model 3

• Molecule 1: ALPHA-CONOTOXIN SI



4.2.4 Score per residue for model 4

• Molecule 1: ALPHA-CONOTOXIN SI



4.2.5 Score per residue for model 5

\bullet Molecule 1: ALPHA-CONOTOXIN SI

Chain A:	43%	14%	14%	29%
11 C2 C2 C7 N4 N4 N4 N4 X10 X11 S12 C13 C13 C13 C13 C13 C13 C13 C13 C12 C1 C1 C1 C1 C1 C1 C2 C2 C2 C2 C2 C3 C3 C3 C3 C3 C3 C3 C3 C3 C3 C3 C3 C3				


4.2.6 Score per residue for model 6

• Molecule 1: ALPHA-CONOTOXIN SI

Chain A:	43%	14%	14%	29%
11 62 63 63 71 11 812 812 013 013 013				

4.2.7 Score per residue for model 7

\bullet Molecule 1: ALPHA-CONOTOXIN SI





4.2.8 Score per residue for model 8

• Molecule 1: ALPHA-CONOTOXIN SI



4.2.9 Score per residue for model 9

• Molecule 1: ALPHA-CONOTOXIN SI



4.2.10 Score per residue for model 10

\bullet Molecule 1: ALPHA-CONOTOXIN SI

Chain A:	43%	14%	14%	29%
11 12 13 13 14 14 14 14 14 14 14 14 14 14 14 14 14				


4.2.11 Score per residue for model 11

• Molecule 1: ALPHA-CONOTOXIN SI




4.2.12 Score per residue for model 12

• Molecule 1: ALPHA-CONOTOXIN SI




4.2.13 Score per residue for model 13

• Molecule 1: ALPHA-CONOTOXIN SI



4.2.14 Score per residue for model 14

• Molecule 1: ALPHA-CONOTOXIN SI



4.2.15 Score per residue for model 15

• Molecule 1: ALPHA-CONOTOXIN SI




4.2.16 Score per residue for model 16

• Molecule 1: ALPHA-CONOTOXIN SI

Chain A:	50%	7%	14%	29%
11 62 63 63 812 613 613 613 613 613 613 613 613 613 613				


4.2.17 Score per residue for model 17

\bullet Molecule 1: ALPHA-CONOTOXIN SI




4.2.18 Score per residue for model 18

• Molecule 1: ALPHA-CONOTOXIN SI



4.2.19 Score per residue for model 19

• Molecule 1: ALPHA-CONOTOXIN SI



4.2.20 Score per residue for model 20

• Molecule 1: ALPHA-CONOTOXIN SI

Chain A:	36%	36%	29%
11 C2 C2 C3 C3 C4 M4 M2 M2 M2 M10 M110 M124 M124 M124			


4.2.21 Score per residue for model 21

• Molecule 1: ALPHA-CONOTOXIN SI

Chain A:	43%	14%	14%	29%
11 62 63 63 63 711 711 711 711 711 711 711 711 711 71				

4.2.22 Score per residue for model 22


• Molecule 1: ALPHA-CONOTOXIN SI





4.2.23 Score per residue for model 23

• Molecule 1: ALPHA-CONOTOXIN SI



4.2.24 Score per residue for model 24

• Molecule 1: ALPHA-CONOTOXIN SI



4.2.25 Score per residue for model 25

• Molecule 1: ALPHA-CONOTOXIN SI

Chain A:	36%	21%	14%	29%
11 C2 C3 C3 C3 C3 C4 M10 N11 S12 C12 C12 C12 C12 C12 C12 C12 C12 C12 C				


4.2.26 Score per residue for model 26

• Molecule 1: ALPHA-CONOTOXIN SI

Chain A:	50%	7%	14%	29%
11 62 63 63 12 13 12 14 14 14 14 14 14 14 14 14 14 14 14 14				

4.2.27 Score per residue for model 27

\bullet Molecule 1: ALPHA-CONOTOXIN SI



4.2.28 Score per residue for model 28

• Molecule 1: ALPHA-CONOTOXIN SI



4.2.29 Score per residue for model 29

• Molecule 1: ALPHA-CONOTOXIN SI



4.2.30 Score per residue for model 30

• Molecule 1: ALPHA-CONOTOXIN SI




4.2.31 Score per residue for model 31

• Molecule 1: ALPHA-CONOTOXIN SI

Chain A: 64% 7% 29%

4.2.32 Score per residue for model 32

• Molecule 1: ALPHA-CONOTOXIN SI





4.2.33 Score per residue for model 33

• Molecule 1: ALPHA-CONOTOXIN SI

Chain A:	57%	7%	7%	29%
11 62 63 812 63 713 013 013 013				

4.2.34 Score per residue for model 34

• Molecule 1: ALPHA-CONOTOXIN SI

Chain A:	29%	36%	7%	29%
11 C2 C2 C3 C3 C3 F1 F1 F1 F1 F1 F1 F1 F1 F1 F1 F1 F1 F1				

4.2.35 Score per residue for model 35

• Molecule 1: ALPHA-CONOTOXIN SI

Chain A:	36%	21%	14%	29%
11 C2 C2 C3 A 6 B C3 C1 S12 C13 S12 C13 S12 N1214				

- 4.2.36 Score per residue for model 36
- \bullet Molecule 1: ALPHA-CONOTOXIN SI





5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: torsion angle dynamics.

Of the 50 calculated structures, 36 were deposited, based on the following criterion: *LOWEST ENERGY STRUCTURES*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	refinement	
DYANA	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	118
Number of shifts mapped to atoms	118
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	78%



6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NH2

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	А	70	63	63	$2{\pm}1$
All	All	2520	2268	2268	55

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Moo	lels
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:4:ASN:CB	1:A:7:CYS:SG	0.52	2.98	23	25
1:A:11:TYR:C	1:A:11:TYR:CD1	0.44	2.90	32	3
1:A:13:CYS:SG	1:A:13:CYS:O	0.44	2.75	19	4
1:A:7:CYS:O	1:A:10:LYS:N	0.44	2.49	19	20
1:A:7:CYS:O	1:A:9:PRO:HD2	0.41	2.15	34	1
1:A:13:CYS:O	1:A:13:CYS:SG	0.40	2.79	25	1
1:A:11:TYR:CD1	1:A:11:TYR:C	0.40	2.95	34	1

All unique clashes are listed below, sorted by their clash magnitude.



6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Pe	erce	entiles
1	А	10/14~(71%)	$7\pm0~(71\pm5\%)$	$3\pm1~(26\pm8\%)$	$0{\pm}0~(3{\pm}5\%)$		5	37
All	All	360/504~(71%)	256 (71%)	93~(26%)	11 (3%)		5	37

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	А	8	GLY	11

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the side chain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	nalysed Rotameric		Percentiles		
1	А	8/11 (73%)	6 ± 1 (77 $\pm8\%$)	$2\pm1~(23\pm8\%)$	2 26		
All	All	288/396~(73%)	221 (77%)	67 (23%)	2 26		

All 4 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	А	12	SER	36
1	А	4	ASN	17
1	А	10	LYS	13
1	А	13	CYS	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.



6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 78% for the well-defined parts and 78% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: assigned_chem_shift_list_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	118
Number of shifts mapped to atoms	118
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 78%, i.e. 86 atoms were assigned a chemical shift out of a possible 110. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}\mathbf{N}$
Backbone	29/47~(62%)	19/19~(100%)	10/20~(50%)	0/8~(0%)
Sidechain	51/54~(94%)	35/35~(100%)	16/17~(94%)	0/2~(0%)
Aromatic	6/9~(67%)	4/4~(100%)	2/5~(40%)	0/0 (%)
Overall	86/110~(78%)	58/58~(100%)	28/42~(67%)	0/10~(0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 78%, i.e. 113 atoms were assigned a chemical shift out of a possible 144. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.




	Total	$^{1}\mathbf{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Backbone	37/62~(60%)	24/25~(96%)	13/26~(50%)	0/11~(0%)
Sidechain	70/73~(96%)	48/48 (100%)	22/23~(96%)	0/2~(0%)
Aromatic	6/9~(67%)	4/4 (100%)	2/5~(40%)	0/0 (%)
Overall	113/144~(78%)	76/77~(99%)	37/54~(69%)	0/13~(0%)

7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.



Random coil index (RCI) for chain A:



8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	127
Intra-residue (i-j =0)	20
Sequential (i-j =1)	40
Medium range ($ i-j >1$ and $ i-j <5$)	41
Long range $(i-j \ge 5)$	26
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	9.1
Number of long range restraints per residue ¹	1.9

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	1.4	0.18
0.2-0.5 (Medium)	0.3	0.46
>0.5 (Large)	None	None



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations



9 Distance violation analysis (i)


9.1 Summary of distance violations (i)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Destructure torms	Count	$\%^1$	Vie	olated	3	Consis	tently	y Violated ⁴
Restraints type	Count	701	Count	$\%^2$	$\%^1$	Count	$ \%^2 $	$\%^1$
Intra-residue (i-j =0)	20	15.7	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	2	1.6	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	15	11.8	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	3	2.4	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	40	31.5	6	15.0	4.7	0	0.0	0.0
Backbone-Backbone	17	13.4	4	23.5	3.1	0	0.0	0.0
Backbone-Sidechain	20	15.7	2	10.0	1.6	0	0.0	0.0
Sidechain-Sidechain	3	2.4	0	0.0	0.0	0	0.0	0.0
Medium range ($ i-j > 1 \& i-j < 5$)	41	32.3	4	9.8	3.1	0	0.0	0.0
Backbone-Backbone	13	10.2	2	15.4	1.6	0	0.0	0.0
Backbone-Sidechain	21	16.5	2	9.5	1.6	0	0.0	0.0
Sidechain-Sidechain	7	5.5	0	0.0	0.0	0	0.0	0.0
Long range $(i-j \ge 5)$	26	20.5	1	3.8	0.8	0	0.0	0.0
Backbone-Backbone	4	3.1	1	25.0	0.8	0	0.0	0.0
Backbone-Sidechain	15	11.8	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	7	5.5	0	0.0	0.0	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	127	100.0	11	8.7	8.7	0	0.0	0.0
Backbone-Backbone	36	28.3	7	19.4	5.5	0	0.0	0.0
Backbone-Sidechain	71	55.9	4	5.6	3.1	0	0.0	0.0
Sidechain-Sidechain	20	15.7	0	0.0	0.0	0	0.0	0.0

 1 percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models





9.1.1 Bar chart : Distribution of distance restraints and violations (i)

Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å

are not included in the statistics.

Model ID		Nun	nber o	f viola	ations	5	Mean (Å)	Max (Å)	SD^{6} (Å)	Median (Å)
Model ID	IR^1	SQ^2	MR^3	LR ⁴	IC ⁵	Total	Mean (A)	Max (A)	$SD^{*}(A)$	Median (A)
1	0	0	0	0	0	0	0.0	0.0	0.0	0.0
2	0	0	0	0	0	0	0.0	0.0	0.0	0.0
3	0	0	0	0	0	0	0.0	0.0	0.0	0.0
4	0	0	0	0	0	0	0.0	0.0	0.0	0.0
5	0	0	0	0	0	0	0.0	0.0	0.0	0.0
6	0	0	0	0	0	0	0.0	0.0	0.0	0.0
7	0	0	1	0	0	1	0.11	0.11	0.0	0.11
8	0	1	0	0	0	1	0.11	0.11	0.0	0.11
9	0	1	0	0	0	1	0.11	0.11	0.0	0.11
10	0	1	0	0	0	1	0.11	0.11	0.0	0.11

Continued on next page...




Continued from previous page										
Model ID	IR^1		h ber o		ations IC ⁵		Mean (Å)	Max (Å)	SD^{6} (Å)	Median (Å)
		SQ^2	MR^3	LR^4		Total			, , , ,	
11	0	1	0	0	0	1	0.11	0.11	0.0	0.11
12	0	1	1	0	0	2	0.11	0.11	0.0	0.11
13	0	1	0	0	0	1	0.11	0.11	0.0	0.11
14	0	1	0	0	0	1	0.11	0.11	0.0	0.11
15	0	1	0	0	0	1	0.11	0.11	0.0	0.11
16	0	1	0	0	0	1	0.11	0.11	0.0	0.11
17	0	2	1	0	0	3	0.11	0.11	0.0	0.11
18	0	1	0	0	0	1	0.11	0.11	0.0	0.11
19	0	1	1	1	0	3	0.11	0.11	0.0	0.11
20	0	1	1	0	0	2	0.11	0.11	0.0	0.11
21	0	2	0	0	0	2	0.11	0.11	0.0	0.11
22	0	1	1	1	0	3	0.12	0.15	0.02	0.11
23	0	1	1	1	0	3	0.11	0.11	0.0	0.11
24	0	1	1	1	0	3	0.11	0.13	0.01	0.11
25	0	0	2	0	0	2	0.2	0.26	0.06	0.2
26	0	1	1	0	0	2	0.21	0.3	0.09	0.21
27	0	1	2	1	0	4	0.15	0.24	0.06	0.14
28	0	0	1	1	0	2	0.17	0.18	0.01	0.17
29	0	1	1	1	0	3	0.17	0.27	0.07	0.13
30	0	1	1	0	0	2	0.2	0.27	0.07	0.2
31	0	2	1	0	0	3	0.17	0.27	0.07	0.12
32	0	1	1	0	0	2	0.22	0.27	0.05	0.22
33	0	2	1	0	0	3	0.19	0.32	0.09	0.15
34	0	2	2	0	0	4	0.22	0.41	0.11	0.17
35	0	2	2	0	0	4	0.23	0.46	0.13	0.16
36	0	4	1	0	0	5	0.16	0.29	0.07	0.13

Continued from previous page...

 1 Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation





9.2.1 Bar graph : Distance Violation statistics for each model (i)

The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 116(IR:20, SQ:34, MR:37, LR:25, IC:0) restraints are not violated in the ensemble.

Nu	mber	of vio	lated	Fractio	n of the ensemble		
IR^{1}	SQ^2	MR^3	LR^4	IC ⁵	Total	Count^6	%
0	0	1	0	0	1	1	2.8
0	3	1	0	0	4	2	5.6
0	0	0	0	0	0	3	8.3
0	0	0	0	0	0	4	11.1
0	1	0	0	0	1	5	13.9
0	1	0	0	0	1	6	16.7

Continued on next page...




		of vio				Fractio	n of the ensemble
IR^1	SQ^2	MR^3	LR^4	IC ⁵	Total	Count^6	%
0	0	1	1	0	2	7	19.4
0	0	0	0	0	0	8	22.2
0	0	0	0	0	0	9	25.0
0	0	0	0	0	0	10	27.8
0	0	0	0	0	0	11	30.6
0	0	0	0	0	0	12	33.3
0	0	0	0	0	0	13	36.1
0	0	1	0	0	1	14	38.9
0	0	0	0	0	0	15	41.7
0	0	0	0	0	0	16	44.4
0	0	0	0	0	0	17	47.2
0	0	0	0	0	0	18	50.0
0	1	0	0	0	1	19	52.8
0	0	0	0	0	0	20	55.6
0	0	0	0	0	0	21	58.3
0	0	0	0	0	0	22	61.1
0	0	0	0	0	0	23	63.9
0	0	0	0	0	0	24	66.7
0	0	0	0	0	0	25	69.4
0	0	0	0	0	0	26	72.2
0	0	0	0	0	0	27	75.0
0	0	0	0	0	0	28	77.8
0	0	0	0	0	0	29	80.6
0	0	0	0	0	0	30	83.3
0	0	0	0	0	0	31	86.1
0	0	0	0	0	0	32	88.9
0	0	0	0	0	0	33	91.7
0	0	0	0	0	0	34	94.4
0	0	0	0	0	0	35	97.2
0	0	0	0	0	0	36	100.0

Continued from previous page...

 1 Intra-residue restraints,
 2 Sequential restraints, 3 Medium range restraints,
 4 Long range restraints, 5 Inter-chain restraints,
 6 Number of models with violations






9.3.1 Bar graph : Distance violation statistics for the ensemble (i)

9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram : Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	19	0.11	0.0	0.11
(1,50)	1:7:A:CYS:HB2	1:10:A:LYS:H	14	0.12	0.02	0.12
(1,82)	1:10:A:LYS:HA	1:12:A:SER:H	7	0.19	0.04	0.17
(1,22)	1:3:A:CYS:HA	1:12:A:SER:H	7	0.12	0.03	0.11
(1,7)	1:1:A:ILE:HB	1:2:A:CYS:H	6	0.28	0.02	0.27
(1,80)	1:10:A:LYS:H	1:11:A:TYR:HA	5	0.1	0.0	0.1
(1,95)	1:12:A:SER:H	1:13:A:CYS:H	2	0.44	0.03	0.44
(1,29)	1:4:A:ASN:H	1:8:A:GLY:H	2	0.16	0.01	0.16
(1,19)	1:3:A:CYS:H	1:4:A:ASN:H	2	0.14	0.0	0.14
(1,97)	1:12:A:SER:HA	1:13:A:CYS:H	2	0.12	0.02	0.12


¹Number of violated models, ²Standard deviation



9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,95)	1:12:A:SER:H	1:13:A:CYS:H	35	0.46
(1,95)	1:12:A:SER:H	1:13:A:CYS:H	34	0.41
(1,7)	1:1:A:ILE:HB	1:2:A:CYS:H	33	0.32
(1,98)	1:12:A:SER:HA	1:14:A:NH2:HN1	26	0.3
(1,7)	1:1:A:ILE:HB	1:2:A:CYS:H	36	0.29
(1,7)	1:1:A:ILE:HB	1:2:A:CYS:H	29	0.27
(1,7)	1:1:A:ILE:HB	1:2:A:CYS:H	30	0.27
(1,7)	1:1:A:ILE:HB	1:2:A:CYS:H	31	0.27
(1,7)	1:1:A:ILE:HB	1:2:A:CYS:H	32	0.27
(1,82)	1:10:A:LYS:HA	1:12:A:SER:H	25	0.26

Continued on next page...



10 MW	T
1 2111 11	

Continued from previous page							
Key	Atom-1	Atom-2	Model ID	Violation (Å)			
(1,82)	1:10:A:LYS:HA	1:12:A:SER:H	27	0.24			
(1,82)	1:10:A:LYS:HA	1:12:A:SER:H	32	0.18			
(1,22)	1:3:A:CYS:HA	1:12:A:SER:H	28	0.18			
(1,82)	1:10:A:LYS:HA	1:12:A:SER:H	34	0.17			
(1,29)	1:4:A:ASN:H	1:8:A:GLY:H	34	0.17			
(1,82)	1:10:A:LYS:HA	1:12:A:SER:H	28	0.16			
(1,82)	1:10:A:LYS:HA	1:12:A:SER:H	35	0.16			
(1,50)	1:7:A:CYS:HB2	1:10:A:LYS:H	27	0.16			
(1,50)	1:7:A:CYS:HB2	1:10:A:LYS:H	25	0.15			
(1,50)	1:7:A:CYS:HB2	1:10:A:LYS:H	33	0.15			
(1,29)	1:4:A:ASN:H	1:8:A:GLY:H	35	0.15			
(1,22)	1:3:A:CYS:HA	1:12:A:SER:H	22	0.15			
(1,97)	1:12:A:SER:HA	1:13:A:CYS:H	36	0.14			
(1,19)	1:3:A:CYS:H	1:4:A:ASN:H	34	0.14			
(1,19)	1:3:A:CYS:H	1:4:A:ASN:H	35	0.14			
(1,82)	1:10:A:LYS:HA	1:12:A:SER:H	24	0.13			
(1,50)	1:7:A:CYS:HB2	1:10:A:LYS:H	29	0.13			
(1,50)	1:7:A:CYS:HB2	1:10:A:LYS:H	30	0.13			
(1,50)	1:7:A:CYS:HB2	1:10:A:LYS:H	36	0.13			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	26	0.12			
(1,50)	1:7:A:CYS:HB2	1:10:A:LYS:H	31	0.12			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	8	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	9	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	10	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	11	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	13	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	14	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	15	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	16	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	17	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	18	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	20	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	21	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	22	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	23	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	24	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	27	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	31	0.11			
(1,102)	1:13:A:CYS:HA	1:14:A:NH2:HN1	36	0.11			
(1,97)	1:12:A:SER:HA	1:13:A:CYS:H	33	0.11			
(1,80)	1:10:A:LYS:H	1:11:A:TYR:HA	19	0.11			
(1,80)	1:10:A:LYS:H	1:11:A:TYR:HA	36	0.11			

Continued from previous page...

Continued on next page...



1	\cap	AW	
Т	ωr	VIVV	

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,50)	1:7:A:CYS:HB2	1:10:A:LYS:H	7	0.11
(1,50)	1:7:A:CYS:HB2	1:10:A:LYS:H	12	0.11
(1,50)	1:7:A:CYS:HB2	1:10:A:LYS:H	17	0.11
(1,50)	1:7:A:CYS:HB2	1:10:A:LYS:H	19	0.11
(1,50)	1:7:A:CYS:HB2	1:10:A:LYS:H	20	0.11
(1,50)	1:7:A:CYS:HB2	1:10:A:LYS:H	23	0.11
(1,22)	1:3:A:CYS:HA	1:12:A:SER:H	19	0.11
(1,22)	1:3:A:CYS:HA	1:12:A:SER:H	23	0.11
(1,80)	1:10:A:LYS:H	1:11:A:TYR:HA	12	0.1
(1,80)	1:10:A:LYS:H	1:11:A:TYR:HA	17	0.1
(1,80)	1:10:A:LYS:H	1:11:A:TYR:HA	21	0.1
(1,50)	1:7:A:CYS:HB2	1:10:A:LYS:H	22	0.1
(1,22)	1:3:A:CYS:HA	1:12:A:SER:H	24	0.1
(1,22)	1:3:A:CYS:HA	1:12:A:SER:H	27	0.1
(1,22)	1:3:A:CYS:HA	1:12:A:SER:H	29	0.1

Continued from previous page..



10 Dihedral-angle violation analysis (i)

No dihedral-angle restraints found

