



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 5, 2025 – 06:02 PM EDT

PDB ID : 1RMO / pdb\_00001rmo  
Title : Probing the Role of Tryptophans in Aequorea Victoria Green Fluorescent Proteins with an Expanded Genetic Code  
Authors : Budisa, N.; Pal, P.P.; Alefelder, S.; Birle, P.; Krywcun, T.; Rubini, M.; Wenger, W.; Bae, J.H.; Steiner, T.  
Deposited on : 2003-11-28  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.45.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	236	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1855 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called wunen-nonfunctional GFP fusion protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	224	Total	C	N	O	S	0	0	0
			1791	1140	301	343	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	57	32T	TRP	modified residue	UNP P42212
A	66	CRO	THR	chromophore	UNP P42212
A	66	CRO	TYR	chromophore	UNP P42212
A	66	CRO	GLY	chromophore	UNP P42212

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	64	Total	O	0	0
			64	64		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: wunen-nonfunctional GFP fusion protein

Chain A: 



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.79Å 62.77Å 70.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	17.69 – 1.80	Depositor
% Data completeness (in resolution range)	93.7 (17.69-1.80)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS	Depositor
R, $R_{free}$	0.185 , 0.221	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1855	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CRO, 32T

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.28	5/1793 (0.3%)	1.28	13/2419 (0.5%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	122	ARG	NE-CZ	6.89	1.40	1.33
1	A	218	MET	SD-CE	6.06	1.94	1.79
1	A	88	MET	SD-CE	5.80	1.94	1.79
1	A	150	VAL	CA-CB	5.22	1.60	1.53
1	A	11	VAL	C-N	5.12	1.37	1.32

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	197	ASP	N-CA-C	-7.11	101.36	110.53
1	A	196	PRO	N-CA-C	6.65	121.67	111.57
1	A	97	THR	N-CA-C	-6.42	98.94	109.40
1	A	190	ASP	N-CA-C	-5.98	105.17	112.88
1	A	216	ASP	N-CA-C	-5.84	101.01	109.59
1	A	34	GLU	CA-C-N	-5.61	117.84	121.65
1	A	34	GLU	C-N-CA	-5.61	117.84	121.65
1	A	116	GLY	CA-C-N	-5.41	114.07	122.67
1	A	116	GLY	C-N-CA	-5.41	114.07	122.67
1	A	210	ASP	N-CA-C	-5.31	100.97	109.04
1	A	84	PHE	N-CA-C	5.18	116.61	111.07
1	A	182	TYR	N-CA-C	-5.05	100.66	108.90
1	A	107	LYS	N-CA-C	-5.00	100.75	108.90

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1791	0	1734	21	0
2	A	64	0	0	5	1
All	All	1855	0	1734	21	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:92:TYR:CE1	1:A:112:VAL:HG21	2.06	0.90
1:A:92:TYR:CZ	1:A:112:VAL:HG21	2.13	0.83
1:A:132:GLU:H	1:A:132:GLU:CD	1.94	0.76
1:A:113:LYS:HE3	1:A:122:ARG:HH21	1.59	0.65
1:A:112:VAL:O	1:A:112:VAL:HG23	1.96	0.64
1:A:43:THR:O	1:A:44:LEU:HD22	2.01	0.60
1:A:80:GLN:HG2	2:A:299:HOH:O	2.01	0.60
1:A:92:TYR:CE1	1:A:112:VAL:CG2	2.83	0.58
1:A:18:LEU:HD23	1:A:18:LEU:C	2.29	0.57
1:A:47:ILE:HD13	1:A:215:ARG:CZ	2.35	0.56
1:A:113:LYS:HE3	1:A:122:ARG:NH2	2.20	0.56
1:A:80:GLN:CG	2:A:299:HOH:O	2.55	0.53
1:A:132:GLU:CD	1:A:132:GLU:N	2.67	0.52
1:A:112:VAL:O	1:A:112:VAL:CG2	2.57	0.52
1:A:163:VAL:HB	1:A:183:GLN:HB3	1.91	0.51
1:A:43:THR:C	1:A:44:LEU:HD22	2.38	0.47
1:A:107:LYS:HD3	2:A:288:HOH:O	2.16	0.45
1:A:166:LYS:HD3	1:A:178:LEU:HD11	1.99	0.44
1:A:80:GLN:CD	2:A:299:HOH:O	2.61	0.44
1:A:225:THR:HG23	2:A:275:HOH:O	2.18	0.43
1:A:92:TYR:CD1	1:A:112:VAL:CG2	3.04	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:270:HOH:O	2:A:302:HOH:O[4_556]	1.69	0.51

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	218/236 (92%)	216 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/205 (94%)	185 (96%)	8 (4%)	26	14

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	GLU
1	A	52	LYS
1	A	117	ASP
1	A	132	GLU
1	A	133	ASP
1	A	178	LEU
1	A	184	GLN
1	A	225	THR



Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	146	ASN
1	A	149	ASN
1	A	157	GLN
1	A	164	ASN
1	A	170	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CRO	A	66	1	22,23,24	2.03	8 (36%)	30,32,34	2.17	5 (16%)
1	32T	A	57	1	11,14,15	0.96	1 (9%)	4,19,21	7.12	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	A	66	1	-	0/12/31/32	0/2/2/2
1	32T	A	57	1	-	1/4/6/8	0/2/2/2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	CRO	OH-CZ	-4.43	1.27	1.37

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	CRO	CE2-CD2	4.09	1.45	1.38
1	A	66	CRO	CG2-CB2	-3.72	1.40	1.46
1	A	66	CRO	O3-C3	2.75	1.35	1.20
1	A	66	CRO	CD1-CG2	2.62	1.44	1.39
1	A	66	CRO	CA3-C3	-2.46	1.40	1.49
1	A	66	CRO	CE1-CZ	2.18	1.43	1.39
1	A	66	CRO	CE2-CZ	2.13	1.43	1.39
1	A	57	32T	CZ3-SUL	2.05	1.81	1.71

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	57	32T	CZ2-CE2-CD2	14.17	120.95	110.62
1	A	66	CRO	CA2-N2-C1	7.13	111.38	105.80
1	A	66	CRO	O2-C2-CA2	-6.07	127.15	131.02
1	A	66	CRO	C2-CA2-N2	-4.68	105.59	108.95
1	A	66	CRO	C3-CA3-N3	2.51	118.13	112.43
1	A	66	CRO	CB2-CA2-N2	2.11	131.62	128.76

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	57	32T	C-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.