



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 12:54 AM EDT

PDB ID : 1RPL
Title : 2.3 ANGSTROMS CRYSTAL STRUCTURE OF THE CATALYTIC DOMAIN OF DNA POLYMERASE BETA
Authors : Davies II, J.F.; Almassy, R.J.
Deposited on : 1994-10-25
Resolution : 2.30 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	251	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2024 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA POLYMERASE BETA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	241	1965	1236	345	376	8	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	59	Total	O	0	0
			59	59		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: DNA POLYMERASE BETA

Chain A:  57% 29% 6%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	120.80Å 63.40Å 38.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.30	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-2.30)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.220 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2024	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.17	2/2004 (0.1%)	1.79	45/2699 (1.7%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	296	TYR	CB-CG	-5.51	1.43	1.51
1	A	173	TYR	CB-CG	-5.48	1.43	1.51

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	254	ARG	NE-CZ-NH1	15.83	128.22	120.30
1	A	254	ARG	NE-CZ-NH2	-15.67	112.46	120.30
1	A	253	ARG	NE-CZ-NH2	-11.69	114.46	120.30
1	A	152	ARG	NE-CZ-NH1	10.04	125.32	120.30
1	A	253	ARG	NE-CZ-NH1	9.62	125.11	120.30
1	A	274	GLY	CA-C-N	-9.35	96.63	117.20
1	A	274	GLY	C-N-CA	9.00	144.19	121.70
1	A	299	ARG	NE-CZ-NH1	8.58	124.59	120.30
1	A	328	ARG	CA-CB-CG	-8.21	95.33	113.40
1	A	195	LEU	CA-CB-CG	7.92	133.52	115.30
1	A	274	GLY	O-C-N	7.68	134.98	122.70
1	A	258	ARG	NE-CZ-NH2	-7.54	116.53	120.30
1	A	173	TYR	CB-CG-CD2	-7.33	116.60	121.00
1	A	126	ARG	NE-CZ-NH1	7.21	123.90	120.30
1	A	326	ARG	NE-CZ-NH1	7.11	123.85	120.30
1	A	222	ARG	NE-CZ-NH1	7.07	123.83	120.30
1	A	206	LYS	N-CA-C	6.71	129.11	111.00
1	A	266	TYR	CA-CB-CG	-6.51	101.02	113.40
1	A	250	TYR	CA-CB-CG	-6.16	101.70	113.40
1	A	333	ARG	NE-CZ-NH2	-6.14	117.23	120.30
1	A	141	LYS	CA-CB-CG	6.03	126.66	113.40
1	A	158	MET	CG-SD-CE	6.00	109.80	100.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	254	ARG	CG-CD-NE	-5.84	99.54	111.80
1	A	172	GLU	CA-CB-CG	-5.74	100.77	113.40
1	A	275	SER	N-CA-CB	5.65	118.98	110.50
1	A	152	ARG	NE-CZ-NH2	-5.58	117.51	120.30
1	A	183	ARG	CB-CG-CD	-5.54	97.19	111.60
1	A	296	TYR	CB-CG-CD2	-5.46	117.72	121.00
1	A	244	GLU	N-CA-C	5.40	125.58	111.00
1	A	98	ASN	N-CA-CB	5.39	120.30	110.60
1	A	125	LEU	CA-CB-CG	5.38	127.67	115.30
1	A	280	LYS	CB-CG-CD	5.37	125.56	111.60
1	A	296	TYR	CD1-CG-CD2	5.37	123.80	117.90
1	A	102	ARG	NE-CZ-NH2	-5.32	117.64	120.30
1	A	275	SER	N-CA-C	-5.31	96.67	111.00
1	A	306	VAL	CA-C-N	-5.30	105.53	117.20
1	A	230	LYS	CA-CB-CG	5.30	125.06	113.40
1	A	182	ARG	NE-CZ-NH2	-5.25	117.67	120.30
1	A	234	LYS	CB-CG-CD	5.25	125.25	111.60
1	A	131	LYS	CA-CB-CG	-5.19	101.98	113.40
1	A	258	ARG	CB-CG-CD	-5.19	98.12	111.60
1	A	324	GLN	CA-CB-CG	-5.15	102.06	113.40
1	A	280	LYS	CG-CD-CE	5.13	127.30	111.90
1	A	98	ASN	CB-CA-C	-5.12	100.16	110.40
1	A	149	ARG	NE-CZ-NH2	-5.05	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1965	0	1944	44	0
2	A	59	0	0	4	0
All	All	2024	0	1944	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:99:PHE:HZ	1:A:122:LEU:HD13	1.58	0.68
1:A:99:PHE:CZ	1:A:122:LEU:HD13	2.30	0.66
1:A:303:VAL:HB	1:A:306:VAL:HG11	1.78	0.66
1:A:320:PHE:O	1:A:323:ILE:HG22	1.95	0.65
1:A:269:VAL:O	1:A:273:THR:HG22	1.99	0.62
1:A:328:ARG:NH1	1:A:335:GLU:HB3	2.17	0.60
1:A:206:LYS:HE3	1:A:208:PRO:HD2	1.85	0.59
1:A:303:VAL:HB	1:A:306:VAL:CG1	2.33	0.59
1:A:299:ARG:HG2	1:A:308:GLY:O	2.05	0.56
1:A:320:PHE:CD2	1:A:327:TYR:HD1	2.24	0.55
1:A:150:ILE:HD13	1:A:253:ARG:HD3	1.90	0.54
1:A:113:LYS:HE3	1:A:114:LEU:HG	1.90	0.54
1:A:273:THR:HG23	2:A:446:HOH:O	2.08	0.53
1:A:97:ILE:HG12	1:A:111:ALA:HB1	1.91	0.52
1:A:96:SER:HB2	1:A:120:LYS:HB2	1.92	0.52
1:A:112:ARG:O	1:A:115:VAL:HG12	2.11	0.51
1:A:285:HIS:CE1	1:A:289:LYS:HE2	2.45	0.51
1:A:252:HIS:HE1	2:A:454:HOH:O	1.94	0.50
1:A:249:GLU:O	1:A:251:PRO:HD2	2.12	0.49
1:A:122:LEU:O	1:A:126:ARG:HB2	2.13	0.49
1:A:266:TYR:HB2	1:A:313:VAL:HG12	1.95	0.49
1:A:323:ILE:HG23	1:A:325:TRP:HB2	1.95	0.48
1:A:275:SER:HB3	1:A:278:PHE:HB3	1.96	0.48
1:A:304:THR:H	1:A:306:VAL:HG12	1.79	0.46
1:A:174:ILE:HB	1:A:196:THR:HG22	1.96	0.46
1:A:279:ASN:HB2	1:A:280:LYS:HE2	1.97	0.46
1:A:97:ILE:O	1:A:97:ILE:HD13	2.16	0.46
1:A:328:ARG:HH12	1:A:335:GLU:HB3	1.79	0.45
1:A:99:PHE:O	1:A:102:ARG:HB2	2.17	0.45
1:A:252:HIS:CE1	2:A:454:HOH:O	2.68	0.45
1:A:305:GLY:HA2	2:A:451:HOH:O	2.17	0.45
1:A:195:LEU:HD13	1:A:259:LEU:HD13	1.99	0.44
1:A:323:ILE:HD13	1:A:323:ILE:HG21	1.66	0.44
1:A:225:THR:OG1	1:A:238:VAL:HG22	2.17	0.43
1:A:133:ASN:HD21	1:A:135:HIS:HB3	1.82	0.43
1:A:295:GLU:H	1:A:295:GLU:CD	2.22	0.43
1:A:287:LEU:HD22	1:A:287:LEU:HA	1.88	0.43
1:A:299:ARG:HB3	1:A:307:ALA:O	2.19	0.42
1:A:133:ASN:ND2	1:A:136:GLN:H	2.18	0.42
1:A:275:SER:HB3	1:A:278:PHE:CB	2.50	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:201:THR:O	1:A:204:SER:HB3	2.20	0.41
1:A:306:VAL:O	1:A:306:VAL:HG13	2.20	0.41
1:A:116:ASP:O	1:A:118:GLY:N	2.53	0.41
1:A:275:SER:HB2	1:A:333:ARG:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	239/251 (95%)	214 (90%)	17 (7%)	8 (3%)	4 2

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	275	SER
1	A	117	GLU
1	A	248	ASN
1	A	304	THR
1	A	206	LYS
1	A	97	ILE
1	A	244	GLU
1	A	247	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	219/229 (96%)	190 (87%)	29 (13%)	4 4

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	96	SER
1	A	97	ILE
1	A	100	LEU
1	A	103	VAL
1	A	116	ASP
1	A	126	ARG
1	A	127	LYS
1	A	129	GLU
1	A	133	ASN
1	A	140	LEU
1	A	150	ILE
1	A	164	ASN
1	A	168	LYS
1	A	187	SER
1	A	195	LEU
1	A	206	LYS
1	A	210	LEU
1	A	221	VAL
1	A	233	THR
1	A	253	ARG
1	A	276	ASP
1	A	280	LYS
1	A	283	ARG
1	A	287	LEU
1	A	300	PRO
1	A	304	THR
1	A	323	ILE
1	A	325	TRP
1	A	335	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	133	ASN
1	A	164	ASN
1	A	199	ASN
1	A	212	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	217	GLN
1	A	252	HIS
1	A	285	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.