



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2024 – 08:24 AM EDT

PDB ID : 1TRH
Title : TWO CONFORMATIONAL STATES OF CANDIDA RUGOSA LIPASE
Authors : Grochulski, P.; Cygler, M.
Deposited on : 1993-11-18
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	534	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4351 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LIPASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	534	4022	2556	659	788	19	0	0	0

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	301	Total	O	0	0
			301	301		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: LIPASE



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	105.00Å 106.70Å 59.80Å 90.00° 94.80° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.10	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.10)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.148 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	4351	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.74	0/4118	1.36	32/5601 (0.6%)

There are no bond length outliers.

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	235	ARG	NE-CZ-NH1	14.27	127.43	120.30
1	A	235	ARG	NE-CZ-NH2	-12.81	113.89	120.30
1	A	279	ARG	NE-CZ-NH1	8.40	124.50	120.30
1	A	161	TRP	CD1-CG-CD2	8.23	112.88	106.30
1	A	119	TRP	CD1-CG-CD2	8.09	112.77	106.30
1	A	161	TRP	CE2-CD2-CG	-7.95	100.94	107.30
1	A	119	TRP	CE2-CD2-CG	-7.56	101.25	107.30
1	A	221	TRP	CE2-CD2-CG	-7.43	101.36	107.30
1	A	188	TRP	CD1-CG-CD2	7.39	112.21	106.30
1	A	221	TRP	CD1-CG-CD2	7.29	112.14	106.30
1	A	188	TRP	CE2-CD2-CG	-7.19	101.55	107.30
1	A	320	TYR	CB-CG-CD2	-7.07	116.76	121.00
1	A	188	TRP	CB-CG-CD1	-6.95	117.97	127.00
1	A	324	ARG	NE-CZ-NH1	6.88	123.74	120.30
1	A	492	TYR	CB-CG-CD2	-6.71	116.97	121.00
1	A	489	TRP	CE2-CD2-CG	-6.71	101.94	107.30
1	A	161	TRP	CG-CD2-CE3	6.70	139.93	133.90
1	A	279	ARG	NE-CZ-NH2	-6.55	117.02	120.30
1	A	489	TRP	CD1-CG-CD2	6.55	111.54	106.30
1	A	522	TYR	CB-CG-CD2	-6.53	117.08	121.00
1	A	419	ARG	NE-CZ-NH1	6.32	123.46	120.30
1	A	161	TRP	CB-CG-CD1	-5.91	119.32	127.00
1	A	188	TRP	CG-CD2-CE3	5.79	139.11	133.90
1	A	419	ARG	NE-CZ-NH2	-5.68	117.46	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	119	TRP	CB-CG-CD1	-5.58	119.74	127.00
1	A	119	TRP	CG-CD2-CE3	5.57	138.91	133.90
1	A	318	ASP	CA-CB-CG	5.50	125.50	113.40
1	A	324	ARG	NE-CZ-NH2	-5.44	117.58	120.30
1	A	156	TYR	CB-CG-CD2	-5.33	117.80	121.00
1	A	405	ARG	CA-CB-CG	5.18	124.80	113.40
1	A	303	ARG	N-CA-C	-5.12	97.17	111.00
1	A	170	LYS	CA-CB-CG	-5.08	102.22	113.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4022	0	3893	28	0
2	A	28	0	26	0	0
3	A	301	0	0	0	0
All	All	4351	0	3919	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:202:LYS:HA	1:A:235:ARG:HD2	1.64	0.79
1:A:338:GLN:HE21	1:A:451:ASN:HD21	1.32	0.76
1:A:183:ARG:HH11	1:A:218:HIS:HD1	1.41	0.68
1:A:167:ASP:H	1:A:256:ASN:HD21	1.45	0.65
1:A:182:GLN:HE22	1:A:214:SER:HB3	1.61	0.65
1:A:20:ASN:HA	1:A:106:PRO:HD3	1.83	0.60
1:A:424:ASN:HD21	1:A:495:SER:H	1.49	0.59
1:A:168:GLU:HB3	1:A:275:LEU:HD22	1.91	0.51
1:A:37:ARG:HB2	1:A:279:ARG:HG2	1.93	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:130:THR:HG23	1:A:155:ASN:HD22	1.77	0.50
1:A:338:GLN:NE2	1:A:451:ASN:HD21	2.04	0.49
1:A:142:SER:HB2	1:A:147:LYS:O	2.14	0.48
1:A:338:GLN:HE21	1:A:451:ASN:ND2	2.08	0.47
1:A:77:ALA:HB1	1:A:445:LEU:HD12	1.98	0.45
1:A:25:GLY:H	1:A:101:ASN:HD22	1.65	0.45
1:A:402:GLN:OE1	1:A:405:ARG:HD2	2.17	0.44
1:A:437:LYS:HE3	1:A:504:MET:CE	2.46	0.44
1:A:253:ILE:HG13	1:A:254:TYR:N	2.32	0.44
1:A:68:THR:O	1:A:71:GLU:HG2	2.17	0.44
1:A:64:ASN:HA	1:A:65:PRO:HD3	1.85	0.43
1:A:36:LEU:HD23	1:A:36:LEU:HA	1.91	0.42
1:A:335:ILE:HG12	1:A:419:ARG:HG3	2.00	0.42
1:A:445:LEU:HB3	1:A:448:PHE:HB3	2.01	0.42
1:A:489:TRP:HA	1:A:490:PRO:HD2	1.81	0.41
1:A:389:SER:HA	1:A:390:PRO:C	2.41	0.41
1:A:386:THR:HA	1:A:394:GLY:O	2.20	0.41
1:A:335:ILE:O	1:A:433:SER:HA	2.21	0.40
1:A:455:PHE:HA	1:A:459:LEU:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	532/534 (100%)	506 (95%)	25 (5%)	1 (0%)	44 45

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	122	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	431/431 (100%)	418 (97%)	13 (3%)	36 40

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	ASN
1	A	68	THR
1	A	109	LYS
1	A	141	LYS
1	A	206	PHE
1	A	275	LEU
1	A	318	ASP
1	A	328	TYR
1	A	405	ARG
1	A	439	LEU
1	A	465	LEU
1	A	508	LEU
1	A	525	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	ASN
1	A	101	ASN
1	A	137	GLN
1	A	155	ASN
1	A	182	GLN
1	A	240	GLN
1	A	256	ASN
1	A	338	GLN
1	A	424	ASN
1	A	456	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	994	1	14,14,15	0.64	0	17,19,21	1.27	2 (11%)
2	NAG	A	990	1	14,14,15	0.67	0	17,19,21	1.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	994	1	-	0/6/23/26	0/1/1/1
2	NAG	A	990	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	994	NAG	C1-O5-C5	4.15	117.74	112.19
2	A	994	NAG	C6-C5-C4	-2.08	107.91	113.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.