



Full wwPDB NMR Structure Validation Report ⓘ

Feb 19, 2022 – 11:41 PM EST

PDB ID : 1TZ1
Title : Solution structure of the PB1 domain of CDC24P (short form)
Authors : Yoshinaga, S.; Terasawa, H.; Ogura, K.; Noda, Y.; Ito, T.; Sumimoto, H.;
Inagaki, F.
Deposited on : 2004-07-09

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.26
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.26


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	80	

2 Ensemble composition and analysis i

This entry contains 20 models. Model 12 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:780-A:803, A:812-A:851 (64)	0.26	12

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 6 clusters and 4 single-model clusters were found.

Cluster number	Models
1	2, 9, 11, 12, 18
2	8, 10, 14
3	16, 20
4	5, 15
5	3, 6
6	7, 19
Single-model clusters	1; 4; 13; 17

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1273 atoms, of which 629 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Cell division control protein 24.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	80	1273	409	629	104	128	3	0

There are 5 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	775	GLY	-	cloning artifact	UNP P11433
A	776	ALA	-	cloning artifact	UNP P11433
A	777	MET	-	cloning artifact	UNP P11433
A	778	GLY	-	cloning artifact	UNP P11433
A	779	SER	-	cloning artifact	UNP P11433

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Cell division control protein 24




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


- Molecule 1: Cell division control protein 24



4.2.2 Score per residue for model 2


- Molecule 1: Cell division control protein 24






4.2.3 Score per residue for model 3

- Molecule 1: Cell division control protein 24




4.2.4 Score per residue for model 4

- Molecule 1: Cell division control protein 24




4.2.5 Score per residue for model 5

- Molecule 1: Cell division control protein 24




4.2.6 Score per residue for model 6

- Molecule 1: Cell division control protein 24




4.2.7 Score per residue for model 7

- Molecule 1: Cell division control protein 24




4.2.8 Score per residue for model 8

- Molecule 1: Cell division control protein 24




4.2.9 Score per residue for model 9

- Molecule 1: Cell division control protein 24




4.2.10 Score per residue for model 10

- Molecule 1: Cell division control protein 24




4.2.11 Score per residue for model 11

- Molecule 1: Cell division control protein 24




4.2.12 Score per residue for model 12 (medoid)

- Molecule 1: Cell division control protein 24




4.2.13 Score per residue for model 13

- Molecule 1: Cell division control protein 24




4.2.14 Score per residue for model 14

- Molecule 1: Cell division control protein 24




4.2.15 Score per residue for model 15

- Molecule 1: Cell division control protein 24




4.2.16 Score per residue for model 16

- Molecule 1: Cell division control protein 24




4.2.17 Score per residue for model 17

- Molecule 1: Cell division control protein 24




4.2.18 Score per residue for model 18

- Molecule 1: Cell division control protein 24




4.2.19 Score per residue for model 19

- Molecule 1: Cell division control protein 24



4.2.20 Score per residue for model 20

- Molecule 1: Cell division control protein 24



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
ARIA	structure solution	
ARIA	refinement	
CNS	structure solution	

No chemical shift data was provided.

6 Model quality

6.1 Standard geometry

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	522	514	514	12±3
All	All	10440	10280	10280	247

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:781:PHE:HB3	1:A:851:ILE:HD12	0.88	1.46	18	7
1:A:841:LEU:HD11	1:A:849:LEU:HD23	0.73	1.61	20	18
1:A:781:PHE:HB3	1:A:851:ILE:HG13	0.64	1.69	13	8
1:A:790:ASN:HD21	1:A:792:ASP:HB3	0.63	1.54	16	4
1:A:799:ASN:HA	1:A:802:ILE:HG12	0.60	1.71	17	10
1:A:795:ILE:O	1:A:799:ASN:HB2	0.59	1.97	9	20
1:A:790:ASN:HD21	1:A:792:ASP:HB2	0.59	1.58	12	2
1:A:785:VAL:HG11	1:A:834:TRP:CZ2	0.58	2.33	13	5
1:A:782:THR:HG22	1:A:848:PHE:HB3	0.58	1.76	17	3
1:A:792:ASP:O	1:A:796:MET:HG3	0.57	1.99	16	20
1:A:841:LEU:HD23	1:A:846:GLU:HB3	0.56	1.77	7	2
1:A:781:PHE:HB2	1:A:851:ILE:HB	0.55	1.78	1	1
1:A:819:GLN:HB3	1:A:850:ASN:HB2	0.55	1.79	20	1
1:A:835:ASN:HA	1:A:838:LYS:HE3	0.55	1.79	6	11

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:818:TYR:CE1	1:A:826:VAL:HB	0.55	2.36	14	4
1:A:785:VAL:HG11	1:A:834:TRP:HZ2	0.55	1.62	13	2
1:A:813:ILE:HD13	1:A:813:ILE:H	0.54	1.62	19	1
1:A:818:TYR:CE1	1:A:826:VAL:HG23	0.53	2.38	6	13
1:A:802:ILE:HD13	1:A:813:ILE:HG12	0.52	1.79	18	3
1:A:783:LEU:HD22	1:A:801:LYS:HE2	0.52	1.80	1	1
1:A:794:LEU:HD22	1:A:834:TRP:CE2	0.51	2.40	9	15
1:A:817:LYS:HE2	1:A:827:VAL:HG13	0.51	1.81	20	2
1:A:821:GLU:HB3	1:A:840:MET:SD	0.50	2.47	6	1
1:A:799:ASN:HA	1:A:802:ILE:CG1	0.50	2.37	20	19
1:A:782:THR:HG23	1:A:850:ASN:ND2	0.50	2.21	13	1
1:A:799:ASN:HD21	1:A:812:PRO:HB3	0.49	1.67	6	1
1:A:790:ASN:ND2	1:A:793:ASP:H	0.48	2.07	16	6
1:A:818:TYR:O	1:A:825:PHE:HA	0.48	2.09	6	6
1:A:782:THR:HA	1:A:849:LEU:O	0.47	2.08	1	6
1:A:826:VAL:CG1	1:A:827:VAL:N	0.47	2.77	8	1
1:A:783:LEU:O	1:A:848:PHE:HA	0.47	2.09	1	5
1:A:832:GLU:O	1:A:836:VAL:HG23	0.47	2.09	8	7
1:A:780:ILE:HA	1:A:851:ILE:O	0.47	2.09	7	2
1:A:785:VAL:HG13	1:A:789:TRP:CE3	0.46	2.45	7	3
1:A:798:ILE:O	1:A:802:ILE:HG23	0.46	2.11	15	5
1:A:834:TRP:HA	1:A:834:TRP:CE3	0.44	2.47	5	3
1:A:831:ASP:O	1:A:835:ASN:HB3	0.44	2.13	7	1
1:A:790:ASN:ND2	1:A:792:ASP:HB2	0.43	2.24	12	2
1:A:784:LEU:O	1:A:784:LEU:HG	0.43	2.13	7	1
1:A:849:LEU:HD13	1:A:850:ASN:O	0.43	2.13	12	2
1:A:799:ASN:HA	1:A:802:ILE:HG13	0.43	1.89	2	1
1:A:786:GLU:HG2	1:A:788:VAL:HG12	0.43	1.91	4	2
1:A:794:LEU:HD13	1:A:834:TRP:CZ2	0.42	2.49	4	1
1:A:802:ILE:HG13	1:A:803:SER:N	0.42	2.29	18	1
1:A:816:ILE:H	1:A:816:ILE:HD13	0.42	1.74	18	1
1:A:834:TRP:O	1:A:838:LYS:HD3	0.42	2.14	19	3
1:A:834:TRP:CE3	1:A:834:TRP:HA	0.42	2.48	18	2
1:A:785:VAL:HG21	1:A:834:TRP:HH2	0.42	1.75	4	1
1:A:816:ILE:HD13	1:A:816:ILE:H	0.42	1.75	20	3
1:A:798:ILE:O	1:A:802:ILE:HG12	0.41	2.15	13	2
1:A:794:LEU:HD13	1:A:795:ILE:N	0.41	2.30	17	1
1:A:799:ASN:HD21	1:A:813:ILE:H	0.41	1.58	15	1
1:A:782:THR:HG23	1:A:850:ASN:OD1	0.41	2.15	3	1
1:A:819:GLN:HA	1:A:824:ASP:O	0.41	2.15	4	1
1:A:841:LEU:O	1:A:846:GLU:HB2	0.41	2.16	13	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	64/80 (80%)	50±2 (78±3%)	12±2 (19±3%)	2±1 (3±1%)	7	41
All	All	1280/1600 (80%)	997 (78%)	246 (19%)	37 (3%)	7	41

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	830	SER	18
1	A	788	VAL	11
1	A	845	ASN	4
1	A	812	PRO	3
1	A	829	GLY	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	59/72 (82%)	38±2 (64±3%)	21±2 (36±3%)	1	8
All	All	1180/1440 (82%)	756 (64%)	424 (36%)	1	8

All 45 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	781	PHE	20
1	A	790	ASN	20
1	A	849	LEU	20
1	A	782	THR	19
1	A	783	LEU	19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	780	ILE	18
1	A	784	LEU	16
1	A	787	LYS	16
1	A	827	VAL	15
1	A	801	LYS	13
1	A	800	SER	12
1	A	833	ASP	12
1	A	835	ASN	12
1	A	799	ASN	12
1	A	793	ASP	12
1	A	843	GLU	11
1	A	786	GLU	11
1	A	830	SER	11
1	A	847	LYS	11
1	A	821	GLU	10
1	A	846	GLU	10
1	A	851	ILE	10
1	A	816	ILE	9
1	A	822	ASP	9
1	A	815	LYS	9
1	A	824	ASP	9
1	A	819	GLN	8
1	A	826	VAL	8
1	A	828	LEU	7
1	A	814	THR	7
1	A	832	GLU	6
1	A	803	SER	5
1	A	839	GLU	5
1	A	844	ASN	5
1	A	831	ASP	4
1	A	802	ILE	4
1	A	845	ASN	4
1	A	834	TRP	4
1	A	820	ASP	3
1	A	838	LYS	2
1	A	836	VAL	2
1	A	848	PHE	1
1	A	840	MET	1
1	A	818	TYR	1
1	A	813	ILE	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided