



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 5, 2024 – 06:40 AM EST

PDB ID : 1XPJ  
Title : Crystal Structure of MCSG Target APC26283 from *Vibrio cholerae*  
Authors : Brunzelle, J.S.; Minasov, G.; Shuvalova, L.; Collart, F.R.; Anderson, W.F.;  
Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2004-10-08  
Resolution : 2.30 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	126	 67% 28% . .
1	B	126	 76% 21% .
1	C	126	 75% 22% .
1	D	126	 74% 20% . .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	TLA	A	303	-	X	-	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4686 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called hypothetical protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	124	1128	719	200	203	6	0	27	0
1	B	123	1093	702	189	196	6	0	20	0
1	C	123	1093	703	186	198	6	0	21	0
1	D	123	1085	697	189	194	5	0	17	0

- Molecule 2 is MERCURY (II) ION (three-letter code: HG) (formula: Hg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Hg	0	0
			1	1		
2	B	1	Total	Hg	0	0
			1	1		
2	C	1	Total	Hg	0	0
			1	1		
2	D	1	Total	Hg	0	0
			1	1		

- Molecule 3 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: C<sub>4</sub>H<sub>6</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 10 4 6	0	0
3	B	1	Total C O 10 4 6	0	0
3	C	1	Total C O 10 4 6	0	0
3	D	1	Total C O 10 4 6	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	66	Total O 66 66	0	0
4	B	49	Total O 49 49	0	0
4	C	59	Total O 59 59	0	0
4	D	69	Total O 69 69	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: hypothetical protein

Chain A: 




- Molecule 1: hypothetical protein

Chain B: 



- Molecule 1: hypothetical protein

Chain C: 



- Molecule 1: hypothetical protein

Chain D: 



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.73Å 52.89Å 72.22Å 90.00° 90.20° 90.00°	Depositor
Resolution (Å)	25.00 – 2.30 24.83 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.1 (25.00-2.30) 97.7 (24.83-2.15)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.79 (at 2.15Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.201 , 0.253 0.203 , 0.246	Depositor DCC
$R_{free}$ test set	1585 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.6	Xtrriage
Anisotropy	0.532	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 26.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.486 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4686	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 54.91 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.4110e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TLA, HG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/1290	0.82	4/1739 (0.2%)
1	B	0.47	0/1220	0.77	1/1649 (0.1%)
1	C	0.42	0/1226	0.75	1/1657 (0.1%)
1	D	0.51	0/1196	0.79	2/1617 (0.1%)
All	All	0.48	0/4932	0.78	8/6662 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	9	ASP	CB-CG-OD2	6.00	123.70	118.30
1	B	9	ASP	CB-CG-OD2	5.36	123.13	118.30
1	A	74	ASP	CB-CG-OD2	5.36	123.12	118.30
1	A	99	ASP	CB-CG-OD2	5.36	123.12	118.30
1	D	99	ASP	CB-CG-OD2	5.32	123.08	118.30
1	D	81	ASP	CB-CG-OD2	5.18	122.96	118.30
1	A	28	ASP	CB-CG-OD2	5.02	122.82	118.30
1	C	9	ASP	CB-CG-OD2	5.02	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1128	0	1166	29	0
1	B	1093	0	1118	26	0
1	C	1093	0	1117	26	0
1	D	1085	0	1108	17	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	10	0	4	3	0
3	B	10	0	4	0	0
3	C	10	0	4	1	0
3	D	10	0	4	0	0
4	A	66	0	0	1	0
4	B	49	0	0	1	0
4	C	59	0	0	3	0
4	D	69	0	0	0	0
All	All	4686	0	4525	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:96[B]:TYR:CD2	1:C:96[B]:TYR:HE2	1.60	1.19
1:B:96[B]:TYR:CE2	1:C:96[B]:TYR:CE2	2.31	1.18
1:B:96[B]:TYR:CD2	1:C:96[B]:TYR:CE2	2.34	1.16
1:B:96[B]:TYR:HE2	1:C:96[B]:TYR:CD2	1.77	1.02
1:A:90[B]:CYS:SG	1:A:96:TYR:CE2	2.58	0.96
1:B:96[B]:TYR:CE2	1:C:96[B]:TYR:CD2	2.54	0.94
1:B:51[A]:MET:HE3	1:B:61:ILE:CD1	2.00	0.92
1:A:90[B]:CYS:SG	1:A:96:TYR:CZ	2.64	0.91
1:C:51[A]:MET:HE3	1:C:61:ILE:CD1	2.04	0.87
1:B:96[B]:TYR:HD2	1:C:96[B]:TYR:HE2	1.21	0.83
1:A:12[B]:LEU:HD21	1:A:33[B]:LEU:HD11	1.61	0.81
1:A:1:MET:CE	1:A:3[A]:LYS:HG3	2.14	0.77
1:B:120[B]:GLU:OE1	1:B:123:LYS:HD2	1.86	0.76
1:A:1:MET:HE3	1:A:3[A]:LYS:HG3	1.69	0.73
1:A:1:MET:HE3	1:A:3[B]:LYS:HG3	1.74	0.70
1:A:87[B]:LYS:HE2	1:D:96[B]:TYR:OH	1.92	0.69
1:B:51[A]:MET:CE	1:B:61:ILE:CD1	2.70	0.69
1:B:51[A]:MET:HE3	1:B:61:ILE:HD12	1.75	0.69

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:ARG:NH1	1:A:71[B]:GLU:OE1	2.30	0.65
1:B:52:ARG:HH12	1:B:92[A]:HIS:CE1	2.16	0.64
1:D:1:MET:CE	1:D:3[A]:LYS:HG3	2.27	0.64
1:A:103[B]:ARG:NH2	1:D:106:GLU:OE1	2.32	0.62
1:C:51[A]:MET:CE	1:C:61:ILE:CD1	2.76	0.62
1:D:1:MET:HE1	1:D:3[A]:LYS:HG3	1.82	0.62
1:A:34[A]:ARG:HH21	1:A:77[A]:GLN:HB2	1.65	0.61
1:C:98:ASP:HB3	1:C:101:ALA:HB2	1.83	0.61
1:A:52:ARG:HG3	1:A:89[B]:TRP:CH2	2.36	0.61
1:A:12[B]:LEU:CD2	1:A:33[B]:LEU:HD11	2.29	0.60
1:A:1:MET:CE	1:A:3[B]:LYS:HG3	2.33	0.58
1:B:51[A]:MET:CE	1:B:61:ILE:HD11	2.33	0.58
1:D:110:MET:HB3	1:D:114:GLU:HG3	1.85	0.57
1:A:119:PHE:O	1:A:122[A]:GLU:HB2	2.05	0.56
1:A:50:ASN:H	3:A:303:TLA:H3	1.70	0.56
1:A:1:MET:HE1	1:A:3[A]:LYS:HG3	1.87	0.56
1:B:98:ASP:HB3	1:B:101:ALA:HB2	1.88	0.56
1:D:98:ASP:HB3	1:D:101:ALA:HB2	1.88	0.56
1:D:12[B]:LEU:HD21	1:D:33[B]:LEU:HD11	1.89	0.55
1:B:110:MET:HB3	1:B:114:GLU:HG3	1.87	0.55
1:C:33:LEU:HG	1:C:43:ILE:HD12	1.90	0.54
1:B:43:ILE:HG12	4:B:443:HOH:O	2.07	0.54
1:C:61:ILE:HG22	4:C:442:HOH:O	2.09	0.53
1:C:43:ILE:HG12	4:C:437:HOH:O	2.09	0.53
1:C:51[A]:MET:CE	1:C:61:ILE:HD11	2.38	0.53
1:C:51[A]:MET:HE3	1:C:61:ILE:HD12	1.86	0.53
1:C:114:GLU:HA	1:C:117:GLN:HE21	1.74	0.52
1:A:96:TYR:HB3	4:A:449:HOH:O	2.10	0.51
1:A:66:LEU:O	1:A:70[A]:THR:HG23	2.10	0.51
1:B:12:LEU:HD21	1:B:33[B]:LEU:HD11	1.92	0.51
1:A:98:ASP:HB3	1:A:101:ALA:HB2	1.93	0.51
1:A:21:ARG:NH1	1:A:71[A]:GLU:OE1	2.44	0.50
1:D:105:SER:O	1:D:109:SER:HB2	2.12	0.49
1:B:103[B]:ARG:NH2	1:C:106[B]:GLU:OE1	2.45	0.49
1:D:20:TYR:CZ	1:D:49[A]:ARG:HD2	2.48	0.48
1:D:1:MET:HE3	1:D:3[A]:LYS:HG3	1.96	0.48
1:B:33[B]:LEU:HD22	1:B:43:ILE:HD12	1.96	0.48
1:C:70:THR:HG22	1:C:83:ILE:CD1	2.44	0.48
1:C:50:ASN:HA	3:C:302:TLA:O11	2.14	0.47
1:B:51[A]:MET:HE2	1:B:61:ILE:HD11	1.95	0.47
1:D:66:LEU:O	1:D:70[B]:THR:HG23	2.15	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:106:GLU:OE1	1:D:103[B]:ARG:NH2	2.49	0.46
1:B:52:ARG:NH1	1:B:92[A]:HIS:CE1	2.82	0.46
1:C:43:ILE:HB	1:C:80:TYR:HA	1.99	0.45
1:A:52:ARG:HB3	3:A:303:TLA:C1	2.47	0.44
1:C:51[A]:MET:HE3	1:C:61:ILE:HD11	1.89	0.44
1:D:1:MET:HE3	1:D:3[B]:LYS:HG3	2.00	0.43
1:B:120[B]:GLU:CD	1:B:123:LYS:HD2	2.38	0.43
1:B:103[B]:ARG:HH11	1:B:103[B]:ARG:HD2	1.63	0.43
1:A:101:ALA:HB1	1:D:101:ALA:HB1	2.00	0.43
1:A:34[A]:ARG:NH2	1:A:77[A]:GLN:HB2	2.34	0.42
1:A:50:ASN:HA	3:A:303:TLA:O11	2.19	0.42
1:C:6:VAL:HG21	1:C:33:LEU:HD11	2.01	0.42
1:B:105:SER:HB3	1:C:122:GLU:OE2	2.19	0.42
1:B:30:ILE:O	1:B:33[B]:LEU:HB2	2.20	0.42
1:B:96[B]:TYR:HD2	1:C:96[B]:TYR:CE2	2.07	0.41
1:C:37:HIS:CE1	4:C:460:HOH:O	2.72	0.41
1:A:58:VAL:HA	1:A:61:ILE:HB	2.02	0.41
1:B:52:ARG:HH12	1:B:92[A]:HIS:HE1	1.66	0.41
1:A:3[A]:LYS:HG2	1:A:42:GLU:HB2	2.02	0.41
1:C:51[A]:MET:CE	1:C:61:ILE:HD12	2.48	0.41
1:D:9:ASP:OD2	1:D:49[A]:ARG:HD3	2.20	0.40
1:C:32:GLN:HA	1:C:32:GLN:OE1	2.21	0.40
1:D:3[B]:LYS:HG2	1:D:42:GLU:HB2	2.03	0.40
1:A:95:PHE:HB2	1:D:100:ARG:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	149/126 (118%)	146 (98%)	3 (2%)	0	<a href="#">100</a> <a href="#">100</a>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	141/126 (112%)	139 (99%)	2 (1%)	0	100	100
1	C	142/126 (113%)	140 (99%)	2 (1%)	0	100	100
1	D	138/126 (110%)	135 (98%)	3 (2%)	0	100	100
All	All	570/504 (113%)	560 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	140/115 (122%)	121 (86%)	19 (14%)	3	3
1	B	132/115 (115%)	119 (90%)	13 (10%)	8	9
1	C	133/115 (116%)	118 (89%)	15 (11%)	6	6
1	D	129/115 (112%)	109 (84%)	20 (16%)	2	2
All	All	534/460 (116%)	467 (88%)	67 (12%)	8	5

All (67) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	2[A]	LYS
1	A	2[B]	LYS
1	A	12[A]	LEU
1	A	12[B]	LEU
1	A	18	SER
1	A	38[A]	GLN
1	A	38[B]	GLN
1	A	39[A]	LEU
1	A	39[B]	LEU
1	A	52	ARG
1	A	75[A]	LYS
1	A	75[B]	LYS

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	77[A]	GLN
1	A	77[B]	GLN
1	A	112[A]	LEU
1	A	112[B]	LEU
1	A	121	LYS
1	A	124	SER
1	B	1	MET
1	B	2[A]	LYS
1	B	2[B]	LYS
1	B	4[A]	LEU
1	B	4[B]	LEU
1	B	18	SER
1	B	21[A]	ARG
1	B	21[B]	ARG
1	B	49[A]	ARG
1	B	49[B]	ARG
1	B	70	THR
1	B	99	ASP
1	B	112	LEU
1	C	1	MET
1	C	4[A]	LEU
1	C	4[B]	LEU
1	C	21[A]	ARG
1	C	21[B]	ARG
1	C	39[A]	LEU
1	C	39[B]	LEU
1	C	49[A]	ARG
1	C	49[B]	ARG
1	C	52	ARG
1	C	84[A]	LEU
1	C	84[B]	LEU
1	C	112[A]	LEU
1	C	112[B]	LEU
1	C	115	ILE
1	D	1	MET
1	D	4[A]	LEU
1	D	4[B]	LEU
1	D	12[A]	LEU
1	D	12[B]	LEU
1	D	18	SER
1	D	33[A]	LEU
1	D	33[B]	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	35	GLU
1	D	49[A]	ARG
1	D	49[B]	ARG
1	D	52[A]	ARG
1	D	52[B]	ARG
1	D	75[A]	LYS
1	D	75[B]	LYS
1	D	109	SER
1	D	112[A]	LEU
1	D	112[B]	LEU
1	D	121	LYS
1	D	123	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	32	GLN
1	B	38	GLN
1	B	117	GLN
1	C	117	GLN
1	D	32	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	TLA	B	301	-	9,9,9	0.76	0	12,12,12	1.35	3 (25%)
3	TLA	D	304	-	9,9,9	0.73	0	12,12,12	1.46	3 (25%)
3	TLA	C	302	-	9,9,9	0.55	0	12,12,12	1.58	3 (25%)
3	TLA	A	303	-	9,9,9	1.23	1 (11%)	12,12,12	1.82	4 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TLA	B	301	-	-	4/12/12/12	-
3	TLA	D	304	-	-	4/12/12/12	-
3	TLA	C	302	-	-	4/12/12/12	-
3	TLA	A	303	-	-	12/12/12/12	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	303	TLA	C2-C1	-3.08	1.48	1.52

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	303	TLA	O41-C4-C3	3.31	122.23	113.27
3	C	302	TLA	O11-C1-O1	-3.11	117.02	124.09
3	C	302	TLA	O11-C1-C2	2.91	121.14	113.27
3	D	304	TLA	O41-C4-C3	2.69	120.54	113.27
3	B	301	TLA	O41-C4-C3	2.64	120.42	113.27
3	D	304	TLA	O41-C4-O4	-2.63	118.13	124.09
3	A	303	TLA	O2-C2-C1	-2.58	105.26	110.66
3	B	301	TLA	O41-C4-O4	-2.57	118.25	124.09

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	304	TLA	O11-C1-C2	2.48	119.98	113.27
3	A	303	TLA	O11-C1-C2	2.36	119.66	113.27
3	C	302	TLA	O41-C4-C3	2.09	118.91	113.27
3	B	301	TLA	O11-C1-C2	2.08	118.88	113.27
3	A	303	TLA	O4-C4-C3	-2.03	116.30	121.63

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	303	TLA	C1-C2-C3-O3
3	A	303	TLA	C1-C2-C3-C4
3	A	303	TLA	O2-C2-C3-O3
3	A	303	TLA	O2-C2-C3-C4
3	A	303	TLA	O1-C1-C2-O2
3	A	303	TLA	O11-C1-C2-O2
3	A	303	TLA	O3-C3-C4-O4
3	A	303	TLA	O1-C1-C2-C3
3	A	303	TLA	O11-C1-C2-C3
3	A	303	TLA	O3-C3-C4-O41
3	C	302	TLA	O11-C1-C2-O2
3	C	302	TLA	O1-C1-C2-O2
3	D	304	TLA	O3-C3-C4-O4
3	D	304	TLA	C2-C3-C4-O4
3	B	301	TLA	C2-C3-C4-O4
3	A	303	TLA	C2-C3-C4-O41
3	A	303	TLA	C2-C3-C4-O4
3	C	302	TLA	O1-C1-C2-C3
3	C	302	TLA	O11-C1-C2-C3
3	D	304	TLA	O3-C3-C4-O41
3	D	304	TLA	C2-C3-C4-O41
3	B	301	TLA	O3-C3-C4-O4
3	B	301	TLA	C2-C3-C4-O41
3	B	301	TLA	O3-C3-C4-O41

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	302	TLA	1	0
3	A	303	TLA	3	0



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	124/126 (98%)	-0.49	0 100 100	18, 23, 26, 31	0
1	B	123/126 (97%)	-0.47	0 100 100	19, 23, 26, 29	0
1	C	123/126 (97%)	-0.53	1 (0%) 86 89	20, 23, 26, 30	0
1	D	123/126 (97%)	-0.48	0 100 100	17, 23, 26, 32	0
All	All	493/504 (97%)	-0.49	1 (0%) 95 96	17, 23, 26, 32	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	123	LYS	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	TLA	A	303	10/10	0.90	0.13	51,54,59,59	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	HG	D	404	1/1	0.91	0.08	65,65,65,65	1
2	HG	B	402	1/1	0.91	0.09	66,66,66,66	1
3	TLA	D	304	10/10	0.94	0.11	48,49,51,51	0
3	TLA	C	302	10/10	0.96	0.10	38,41,43,43	0
3	TLA	B	301	10/10	0.97	0.10	37,40,43,43	0
2	HG	A	401	1/1	0.98	0.09	64,64,64,64	1
2	HG	C	403	1/1	0.99	0.09	66,66,66,66	1

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.