



# Full wwPDB X-ray Structure Validation Report i

Oct 12, 2021 – 09:11 AM EDT

PDB ID : 1YQD  
Title : Sinapyl Alcohol Dehydrogenase complexed with NADP+  
Authors : Bomati, E.K.; Noel, J.P.  
Deposited on : 2005-02-01  
Resolution : 1.65 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	DTT	A	700	X	-	-	-
4	DTT	B	800	X	-	-	-
5	GOL	B	900	-	X	-	-

## 2 Entry composition [\(i\)](#)

There are 6 unique types of molecules in this entry. The entry contains 6020 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called sinapyl alcohol dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	359	Total	C 2717	N 1729	O 462	S 510	16	0	0
1	B	359	Total	C 2717	N 1729	O 462	S 510	16	0	0


There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	cloning artifact	UNP Q94G59
A	-2	SER	-	cloning artifact	UNP Q94G59
A	-1	HIS	-	cloning artifact	UNP Q94G59
A	0	GLY	-	cloning artifact	UNP Q94G59
A	140	ASN	ASP	engineered mutation	UNP Q94G59
B	-3	GLY	-	cloning artifact	UNP Q94G59
B	-2	SER	-	cloning artifact	UNP Q94G59
B	-1	HIS	-	cloning artifact	UNP Q94G59
B	0	GLY	-	cloning artifact	UNP Q94G59
B	140	ASN	ASP	engineered mutation	UNP Q94G59

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Zn 2 2	0	0
2	B	2	Total Zn 2 2	0	0

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C<sub>21</sub>H<sub>28</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total		C	N	O	P	
			48		21	7	17	3	
3	B	1	Total		C	N	O	P	
			48		21	7	17	3	

- Molecule 4 is 2,3-DIHYDROXY-1,4-DITHIOBUTANE (three-letter code: DTT) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>2</sub>S<sub>2</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total		C	O	S	
			8		4	2	2	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O S 8 4 2 2	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 6 3 3	0	0



- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	246	Total O 246 246	0	0
6	B	218	Total O 218 218	0	0



### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: sinapyl alcohol dehydrogenase



- Molecule 1: sinapyl alcohol dehydrogenase



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.48Å 137.80Å 69.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.32 – 1.65 38.32 – 1.65	Depositor EDS
% Data completeness (in resolution range)	94.0 (38.32-1.65) 94.0 (38.32-1.65)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	3.31 (at 1.65Å)	Xtriage
Refinement program	CNS 1.1	Depositor
$R$ , $R_{free}$	0.195 , 0.213 0.192 , 0.207	Depositor DCC
$R_{free}$ test set	4225 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.5	Xtriage
Anisotropy	0.168	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 44.5	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.49$ , $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6020	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: DTT, GOL, NAP, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.29	0/2778	0.60	0/3760
1	B	0.29	0/2778	0.59	0/3760
All	All	0.29	0/5556	0.60	0/7520

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2717	0	2698	39	0
1	B	2717	0	2698	41	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	48	0	25	4	0
3	B	48	0	25	4	0
4	A	8	0	10	2	0
4	B	8	0	10	1	0
5	B	6	0	4	0	0
6	A	246	0	0	3	0
6	B	218	0	0	5	0
All	All	6020	0	5470	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:500:NAP:O4B	3:A:500:NAP:C4B	1.64	1.45
3:B:600:NAP:O4B	3:B:600:NAP:C4B	1.64	1.38
1:B:115:ASN:H	1:B:115:ASN:HD22	1.23	0.85
1:A:281:GLU:HG2	1:A:282:LYS:HG3	1.70	0.71
1:B:271:HIS:HE1	1:B:295:ARG:HH21	1.38	0.70
1:B:115:ASN:HD22	1:B:115:ASN:N	1.85	0.69
1:A:271:HIS:CE1	1:A:295:ARG:HE	2.11	0.68
1:B:65:MET:HG2	6:B:4198:HOH:O	1.93	0.67
1:A:271:HIS:HE1	1:A:295:ARG:HH21	1.40	0.67
1:B:155:ASN:HD22	1:B:155:ASN:H	1.42	0.67
1:A:87:LYS:HG2	1:A:146:GLU:OE1	1.98	0.64
1:B:155:ASN:HD22	1:B:155:ASN:N	1.95	0.64
1:B:343:ASN:HA	6:B:4152:HOH:O	1.97	0.64
1:A:282:LYS:HB3	1:A:283:PRO:HD2	1.81	0.62
1:A:176:LYS:HD3	1:A:181:ASP:OD1	1.98	0.62
1:A:104:HIS:HE1	1:A:148:TYR:OH	1.83	0.62
1:B:271:HIS:CE1	1:B:295:ARG:HE	2.16	0.62
1:A:297:ILE:HG23	1:B:297:ILE:HG23	1.83	0.60
1:B:255:VAL:HG12	1:B:257:ALA:H	1.66	0.59
1:B:155:ASN:H	1:B:155:ASN:ND2	2.01	0.58
1:B:277:VAL:O	3:B:600:NAP:H2N	2.03	0.58
1:A:277:VAL:O	3:A:500:NAP:H2N	2.04	0.58
1:A:271:HIS:CE1	1:A:295:ARG:HH21	2.21	0.58
1:B:115:ASN:H	1:B:115:ASN:ND2	1.99	0.57
1:A:87:LYS:HG3	1:A:88:VAL:HG23	1.87	0.56
1:B:6:GLU:H	1:B:6:GLU:CD	2.08	0.56
1:A:255:VAL:HG12	1:A:257:ALA:H	1.71	0.56
1:A:262:LEU:HB3	1:A:263:PRO:HD3	1.87	0.56
1:B:262:LEU:HB3	1:B:263:PRO:HD3	1.87	0.55
1:A:222:GLU:HG3	6:A:2123:HOH:O	2.07	0.54
1:A:70:PRO:HG2	1:A:138:TYR:CE2	2.43	0.54
1:B:17:ALA:HB2	1:B:69:VAL:HG22	1.90	0.54
1:A:185:LYS:HB2	6:A:2097:HOH:O	2.09	0.53
1:B:125:ALA:HB1	6:B:4198:HOH:O	2.07	0.53
1:A:41:ARG:HH21	1:A:141:HIS:HE2	1.57	0.52
1:B:70:PRO:HG2	1:B:138:TYR:CE2	2.45	0.52
1:A:17:ALA:HB2	1:A:69:VAL:HG22	1.92	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:78:VAL:HG21	1:A:88:VAL:CG1	2.40	0.52
1:B:37:GLU:OE1	1:B:84:LYS:HE3	2.10	0.52
1:A:166:CYS:SG	3:A:500:NAP:H5N	2.51	0.51
1:B:333:LEU:HD21	1:B:351:ILE:HD11	1.93	0.51
1:B:115:ASN:N	1:B:115:ASN:ND2	2.57	0.50
1:B:271:HIS:HE1	1:B:295:ARG:NH2	2.09	0.50
1:B:271:HIS:CE1	1:B:295:ARG:HH21	2.25	0.50
1:B:162:ALA:HB3	1:B:163:PRO:HD3	1.94	0.49
1:B:120:MET:CE	1:B:122:LEU:HG	2.43	0.49
1:A:162:ALA:HB3	1:A:163:PRO:HD3	1.94	0.49
1:A:271:HIS:HE1	1:A:295:ARG:NH2	2.09	0.49
1:A:237:ASP:OD2	1:A:239:GLU:HB3	2.12	0.49
1:B:155:ASN:HD21	1:B:317:ALA:HB1	1.78	0.49
1:B:291:LEU:HD21	1:B:298:VAL:HG23	1.96	0.48
1:A:182:GLU:CD	1:A:185:LYS:HE3	2.34	0.47
1:B:262:LEU:C	1:B:262:LEU:HD13	2.35	0.47
1:B:270:SER:OG	1:B:271:HIS:HD2	1.98	0.47
1:A:281:GLU:OE1	1:A:281:GLU:N	2.48	0.46
1:A:271:HIS:HE1	1:A:295:ARG:HE	1.60	0.46
1:B:65:MET:HG3	6:B:4041:HOH:O	2.16	0.46
1:B:166:CYS:SG	3:B:600:NAP:H5N	2.56	0.46
3:B:600:NAP:N7N	4:B:800:DTT:S4	2.89	0.45
1:B:120:MET:HE1	1:B:122:LEU:HG	1.98	0.45
1:A:152:PHE:HA	1:A:153:PRO:HD3	1.89	0.45
1:A:345:VAL:HG12	6:A:2146:HOH:O	2.16	0.44
1:B:345:VAL:HG12	6:B:4081:HOH:O	2.16	0.44
1:B:334:ASN:O	1:B:338:GLU:HG2	2.17	0.44
1:A:233:LEU:HD22	1:A:240:GLN:HG2	1.99	0.44
1:A:87:LYS:HE2	1:A:146:GLU:HG2	2.00	0.43
1:B:41:ARG:HH21	1:B:141:HIS:HE2	1.66	0.43
1:A:297:ILE:HG12	1:B:299:ALA:HB2	2.00	0.43
1:B:281:GLU:N	1:B:281:GLU:OE1	2.49	0.43
1:A:333:LEU:HD21	1:A:351:ILE:HD11	2.01	0.42
1:B:222:GLU:HA	1:B:226:ASN:OD1	2.20	0.42
1:B:346:ARG:HA	1:B:347:TYR:HA	1.85	0.42
1:A:346:ARG:HA	1:A:347:TYR:HA	1.81	0.41
1:A:78:VAL:HG21	1:A:88:VAL:HG11	2.01	0.41
1:A:298:VAL:CG1	1:B:298:VAL:HB	2.50	0.41
1:A:52:SER:CB	4:A:700:DTT:O2	2.69	0.41
1:A:70:PRO:HG2	1:A:138:TYR:CD2	2.56	0.41
3:A:500:NAP:O4B	3:A:500:NAP:C5B	2.56	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:66:TYR:HA	1:B:67:PRO:C	2.42	0.41
1:A:52:SER:HB3	4:A:700:DTT:O2	2.22	0.40
1:A:182:GLU:HB2	1:A:185:LYS:HD2	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	357/366 (98%)	346 (97%)	10 (3%)	1 (0%)	41 22
1	B	357/366 (98%)	346 (97%)	10 (3%)	1 (0%)	41 22
All	All	714/732 (98%)	692 (97%)	20 (3%)	2 (0%)	41 22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	303	ILE
1	B	303	ILE

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	291/296 (98%)	287 (99%)	4 (1%)	67 46

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	291/296 (98%)	282 (97%)	9 (3%)	40 14
All	All	582/592 (98%)	569 (98%)	13 (2%)	52 27

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	72	HIS
1	A	281	GLU
1	A	343	ASN
1	A	348	ARG
1	B	6	GLU
1	B	59	ASN
1	B	72	HIS
1	B	115	ASN
1	B	120	MET
1	B	152	PHE
1	B	155	ASN
1	B	343	ASN
1	B	348	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	104	HIS
1	A	226	ASN
1	A	242	GLN
1	A	271	HIS
1	A	343	ASN
1	A	355	ASN
1	B	21	GLN
1	B	24	HIS
1	B	59	ASN
1	B	111	ASN
1	B	115	ASN
1	B	155	ASN
1	B	240	GLN
1	B	271	HIS
1	B	320	ASN
1	B	343	ASN
1	B	355	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	DTT	B	800	-	7,7,7	0.41	0	4,8,8	2.04	2 (50%)
5	GOL	B	900	-	5,5,5	4.38	4 (80%)	5,5,5	5.80	3 (60%)
3	NAP	A	500	-	45,52,52	3.49	18 (40%)	56,80,80	2.51	19 (33%)
4	DTT	A	700	-	7,7,7	0.39	0	4,8,8	2.03	2 (50%)
3	NAP	B	600	-	45,52,52	3.45	18 (40%)	56,80,80	2.51	19 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DTT	B	800	-	2/2/2/2	6/8/8/8	-
5	GOL	B	900	-	-	2/4/4/4	-
3	NAP	A	500	-	-	6/31/67/67	0/5/5/5

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DTT	A	700	-	2/2/2/2	6/8/8/8	-
3	NAP	B	600	-	-	6/31/67/67	0/5/5/5

All (40) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	500	NAP	O4B-C1B	-11.36	1.25	1.41
3	B	600	NAP	O4B-C1B	-11.33	1.25	1.41
3	B	600	NAP	O4B-C4B	8.91	1.64	1.45
3	A	500	NAP	O4B-C4B	8.83	1.64	1.45
5	B	900	GOL	C3-C2	-7.41	1.21	1.51
3	A	500	NAP	C3B-C4B	6.60	1.69	1.53
3	B	600	NAP	C3B-C4B	6.54	1.69	1.53
3	A	500	NAP	C2A-N3A	6.26	1.42	1.32
3	B	600	NAP	C2A-N3A	6.23	1.42	1.32
3	A	500	NAP	P2B-O2B	5.92	1.70	1.59
3	A	500	NAP	C2N-N1N	5.62	1.41	1.35
3	B	600	NAP	P2B-O2B	5.53	1.69	1.59
3	A	500	NAP	C2D-C1D	-5.20	1.45	1.53
3	A	500	NAP	C2A-N1A	5.19	1.43	1.33
3	B	600	NAP	C2N-N1N	5.16	1.41	1.35
3	B	600	NAP	C2A-N1A	5.13	1.43	1.33
3	B	600	NAP	C2D-C1D	-5.04	1.46	1.53
3	A	500	NAP	C4A-N3A	4.87	1.42	1.35
3	B	600	NAP	C4A-N3A	4.77	1.42	1.35
5	B	900	GOL	O1-C1	4.72	1.62	1.42
3	B	600	NAP	C4N-C3N	4.48	1.47	1.39
3	A	500	NAP	C4N-C3N	4.37	1.46	1.39
3	A	500	NAP	O3B-C3B	3.98	1.52	1.43
3	B	600	NAP	O3B-C3B	3.98	1.52	1.43
5	B	900	GOL	O3-C3	3.38	1.56	1.42
3	B	600	NAP	C6N-N1N	3.36	1.43	1.35
3	A	500	NAP	C6N-N1N	3.32	1.43	1.35
3	B	600	NAP	C3B-C2B	-2.91	1.46	1.52
3	A	500	NAP	C3B-C2B	-2.88	1.46	1.52
3	B	600	NAP	PA-O1A	-2.81	1.41	1.50
3	A	500	NAP	PA-O1A	-2.80	1.41	1.50
3	B	600	NAP	O4D-C1D	2.77	1.44	1.41
5	B	900	GOL	O2-C2	-2.70	1.35	1.43
3	A	500	NAP	O4D-C1D	2.60	1.44	1.41
3	A	500	NAP	C6N-C5N	2.40	1.43	1.38
3	B	600	NAP	C5N-C4N	2.27	1.43	1.38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	500	NAP	C5N-C4N	2.26	1.43	1.38
3	A	500	NAP	C3D-C4D	2.22	1.58	1.53
3	B	600	NAP	C3D-C4D	2.17	1.58	1.53
3	B	600	NAP	C6N-C5N	2.15	1.43	1.38

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	500	NAP	O4B-C4B-C3B	-10.66	84.01	105.11
3	B	600	NAP	O4B-C4B-C3B	-10.62	84.10	105.11
5	B	900	GOL	O3-C3-C2	10.53	160.70	110.20
5	B	900	GOL	O2-C2-C3	6.69	138.61	109.12
3	B	600	NAP	O3B-C3B-C4B	-4.82	97.11	111.05
3	A	500	NAP	O3B-C3B-C4B	-4.78	97.22	111.05
3	B	600	NAP	O5D-PN-O1N	-4.65	90.91	109.07
3	B	600	NAP	C5A-C6A-N6A	4.58	127.32	120.35
3	A	500	NAP	O5D-PN-O1N	-4.55	91.29	109.07
3	A	500	NAP	C5A-C6A-N6A	4.51	127.20	120.35
3	A	500	NAP	C2B-C3B-C4B	4.42	111.59	101.99
3	B	600	NAP	C2B-C3B-C4B	4.37	111.48	101.99
3	A	500	NAP	N3A-C2A-N1A	-3.98	122.45	128.68
3	B	600	NAP	N3A-C2A-N1A	-3.93	122.53	128.68
3	B	600	NAP	O2N-PN-O5D	-3.82	90.00	107.75
3	A	500	NAP	O2N-PN-O5D	-3.77	90.23	107.75
3	B	600	NAP	O3B-C3B-C2B	-3.73	100.57	111.17
3	A	500	NAP	O3B-C3B-C2B	-3.72	100.59	111.17
3	A	500	NAP	O5B-C5B-C4B	-3.61	96.58	108.99
3	B	600	NAP	O5B-C5B-C4B	-3.60	96.61	108.99
5	B	900	GOL	O1-C1-C2	3.42	126.58	110.20
3	A	500	NAP	C3B-C2B-C1B	-3.24	96.80	102.89
3	B	600	NAP	C3B-C2B-C1B	-3.21	96.86	102.89
4	B	800	DTT	C2-C1-S1	-2.88	106.10	114.47
4	A	700	DTT	C2-C1-S1	-2.87	106.11	114.47
4	B	800	DTT	C3-C4-S4	-2.80	106.33	114.47
3	B	600	NAP	C1B-N9A-C4A	-2.80	121.73	126.64
3	A	500	NAP	C1B-N9A-C4A	-2.80	121.73	126.64
4	A	700	DTT	C3-C4-S4	-2.79	106.35	114.47
3	A	500	NAP	O2N-PN-O1N	2.79	126.03	112.24
3	B	600	NAP	O2N-PN-O1N	2.79	126.02	112.24
3	A	500	NAP	C2N-C3N-C4N	2.73	121.35	118.26
3	B	600	NAP	C2N-C3N-C4N	2.69	121.30	118.26
3	A	500	NAP	O2A-PA-O1A	2.64	125.31	112.24

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	600	NAP	O2A-PA-O1A	2.64	125.28	112.24
3	A	500	NAP	C5N-C6N-N1N	-2.35	117.03	120.40
3	A	500	NAP	O2B-C2B-C1B	-2.33	101.70	110.10
3	B	600	NAP	C5N-C6N-N1N	-2.32	117.07	120.40
3	B	600	NAP	O2B-C2B-C1B	-2.31	101.78	110.10
3	B	600	NAP	O3X-P2B-O2X	2.31	116.45	107.64
3	A	500	NAP	O3X-P2B-O2X	2.24	116.18	107.64
3	A	500	NAP	C5N-C4N-C3N	-2.23	117.71	120.34
3	B	600	NAP	C5N-C4N-C3N	-2.23	117.71	120.34
3	A	500	NAP	C3N-C7N-N7N	2.19	120.37	117.75
3	B	600	NAP	C3N-C7N-N7N	2.14	120.32	117.75

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	700	DTT	C2
4	A	700	DTT	C3
4	B	800	DTT	C2
4	B	800	DTT	C3

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	500	NAP	O4D-C1D-N1N-C2N
3	A	500	NAP	O4D-C1D-N1N-C6N
3	A	500	NAP	C2D-C1D-N1N-C2N
3	A	500	NAP	C2D-C1D-N1N-C6N
3	B	600	NAP	O4D-C1D-N1N-C2N
3	B	600	NAP	O4D-C1D-N1N-C6N
3	B	600	NAP	C2D-C1D-N1N-C2N
3	B	600	NAP	C2D-C1D-N1N-C6N
4	A	700	DTT	C1-C2-C3-O3
4	A	700	DTT	C1-C2-C3-C4
4	A	700	DTT	O2-C2-C3-O3
4	A	700	DTT	O2-C2-C3-C4
4	A	700	DTT	C2-C3-C4-S4
4	A	700	DTT	O3-C3-C4-S4
4	B	800	DTT	C1-C2-C3-O3
4	B	800	DTT	C1-C2-C3-C4
4	B	800	DTT	O2-C2-C3-O3
4	B	800	DTT	O2-C2-C3-C4
4	B	800	DTT	C2-C3-C4-S4

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
4	B	800	DTT	O3-C3-C4-S4
5	B	900	GOL	O1-C1-C2-C3
5	B	900	GOL	C1-C2-C3-O3
3	A	500	NAP	C3B-C4B-C5B-O5B
3	B	600	NAP	C3B-C4B-C5B-O5B
3	A	500	NAP	O4B-C4B-C5B-O5B
3	B	600	NAP	O4B-C4B-C5B-O5B


There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	800	DTT	1	0
3	A	500	NAP	4	0
4	A	700	DTT	2	0
3	B	600	NAP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	359/366 (98%)	-0.07	13 (3%) 42 43	10, 16, 28, 37	0
1	B	359/366 (98%)	-0.03	12 (3%) 46 47	10, 17, 29, 37	0
All	All	718/732 (98%)	-0.05	25 (3%) 44 45	10, 16, 29, 37	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	362	PRO	5.8
1	A	63	PHE	5.5
1	B	63	PHE	5.2
1	A	239	GLU	3.9
1	B	184	GLY	3.1
1	B	238	GLN	3.1
1	A	281	GLU	3.1
1	A	289	PHE	2.8
1	B	155	ASN	2.6
1	A	184	GLY	2.5
1	B	289	PHE	2.5
1	B	182	GLU	2.4
1	A	238	GLN	2.4
1	B	7	GLU	2.3
1	A	242	GLN	2.3
1	B	65	MET	2.2
1	B	154	ASP	2.2
1	A	105	SER	2.2
1	A	7	GLU	2.1
1	B	87	LYS	2.1
1	A	236	ARG	2.1
1	B	293	ALA	2.1
1	B	281	GLU	2.0
1	A	65	MET	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	155	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	B	900	6/6	0.56	0.24	27,27,28,30	0
4	DTT	B	800	8/8	0.73	0.16	22,26,31,36	0
4	DTT	A	700	8/8	0.78	0.15	18,23,30,36	0
3	NAP	B	600	48/48	0.92	0.11	1,13,18,21	0
3	NAP	A	500	48/48	0.92	0.11	1,12,15,17	0
2	ZN	A	2000	1/1	0.98	0.03	17,17,17,17	0
2	ZN	B	3000	1/1	0.99	0.03	17,17,17,17	0
2	ZN	B	4000	1/1	1.00	0.05	14,14,14,14	0
2	ZN	A	1000	1/1	1.00	0.03	13,13,13,13	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.