



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 28, 2024 – 09:27 pm BST

PDB ID : 2CCF
Title : Antiparallel Configuration of pLI E20S
Authors : Yadav, M.K.; Leman, L.J.; Price, D.J.; Brooks 3rd, C.L.; Stout, C.D.; Ghadiri, M.R.
Deposited on : 2006-01-16
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	34	 32% 47% 12% 9%
1	B	34	 32% 59% 6%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 558 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called General control protein GCN4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	31	253	163	43	46	1	0	0	0
1	B	32	255	163	43	49		0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	ACE	-	acetylation	UNP P03069
A	5	ILE	LEU	engineered mutation	UNP P03069
A	9	LEU	VAL	engineered mutation	UNP P03069
A	12	ILE	LEU	engineered mutation	UNP P03069
A	16	LEU	ASN	engineered mutation	UNP P03069
A	19	ILE	LEU	engineered mutation	UNP P03069
A	20	SER	GLU	engineered mutation	UNP P03069
A	23	LEU	VAL	engineered mutation	UNP P03069
A	26	ILE	LEU	engineered mutation	UNP P03069
A	30	LEU	VAL	engineered mutation	UNP P03069
B	0	ACE	-	acetylation	UNP P03069
B	5	ILE	LEU	engineered mutation	UNP P03069
B	9	LEU	VAL	engineered mutation	UNP P03069
B	12	ILE	LEU	engineered mutation	UNP P03069
B	16	LEU	ASN	engineered mutation	UNP P03069
B	19	ILE	LEU	engineered mutation	UNP P03069
B	20	SER	GLU	engineered mutation	UNP P03069
B	23	LEU	VAL	engineered mutation	UNP P03069
B	26	ILE	LEU	engineered mutation	UNP P03069
B	30	LEU	VAL	engineered mutation	UNP P03069

- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	19	Total O 19 19	0	0
2	B	31	Total O 31 31	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: General control protein GCN4

Chain A: 



- Molecule 1: General control protein GCN4

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	35.39Å 35.39Å 104.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.13 – 1.70 29.28 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.2 (52.13-1.70) 98.2 (29.28-1.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	12.67 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.245 , 0.316 0.242 , 0.314	Depositor DCC
R_{free} test set	355 reflections (4.59%)	wwPDB-VP
Wilson B-factor (Å ²)	23.6	Xtrriage
Anisotropy	0.644	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 57.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	558	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.48 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.6072e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	3.10	22/254 (8.7%)	2.93	20/336 (6.0%)
1	B	3.37	34/256 (13.3%)	2.14	6/340 (1.8%)
All	All	3.24	56/510 (11.0%)	2.56	26/676 (3.8%)

All (56) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	6	GLU	CG-CD	13.01	1.71	1.51
1	B	17	TYR	CG-CD2	12.38	1.55	1.39
1	A	4	GLN	CG-CD	10.49	1.75	1.51
1	B	10	GLU	CD-OE1	9.97	1.36	1.25
1	B	1	ARG	N-CA	9.71	1.65	1.46
1	B	32	GLU	CG-CD	9.68	1.66	1.51
1	B	14	SER	N-CA	9.43	1.65	1.46
1	B	25	ARG	CZ-NH1	9.41	1.45	1.33
1	B	10	GLU	CD-OE2	9.00	1.35	1.25
1	B	25	ARG	CZ-NH2	8.87	1.44	1.33
1	B	22	GLU	CD-OE2	8.74	1.35	1.25
1	B	17	TYR	CD1-CE1	8.65	1.52	1.39
1	A	24	ALA	CA-CB	8.61	1.70	1.52
1	B	28	LYS	CD-CE	8.35	1.72	1.51
1	B	11	GLU	CD-OE2	8.30	1.34	1.25
1	A	25	ARG	CD-NE	8.18	1.60	1.46
1	A	14	SER	CB-OG	7.97	1.52	1.42
1	B	10	GLU	CB-CG	7.93	1.67	1.52
1	A	17	TYR	CZ-OH	7.84	1.51	1.37
1	A	25	ARG	NE-CZ	7.79	1.43	1.33
1	A	17	TYR	CE2-CZ	-7.71	1.28	1.38
1	A	6	GLU	CD-OE1	-7.63	1.17	1.25
1	B	28	LYS	CB-CG	7.51	1.72	1.52
1	B	3	LYS	C-O	7.45	1.37	1.23
1	A	31	GLY	N-CA	7.20	1.56	1.46
1	B	22	GLU	CG-CD	7.08	1.62	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	27	LYS	CD-CE	6.70	1.68	1.51
1	B	4	GLN	CA-C	6.28	1.69	1.52
1	A	21	ASN	C-O	6.24	1.35	1.23
1	B	6	GLU	CB-CG	-6.07	1.40	1.52
1	A	11	GLU	CG-CD	6.07	1.61	1.51
1	B	21	ASN	C-O	5.98	1.34	1.23
1	B	24	ALA	C-O	5.88	1.34	1.23
1	B	11	GLU	CA-CB	-5.87	1.41	1.53
1	B	6	GLU	CD-OE2	5.76	1.31	1.25
1	B	15	LYS	C-O	5.75	1.34	1.23
1	A	28	LYS	CB-CG	5.62	1.67	1.52
1	A	15	LYS	CD-CE	5.60	1.65	1.51
1	B	30	LEU	CG-CD2	5.60	1.72	1.51
1	B	25	ARG	CA-CB	-5.56	1.41	1.53
1	B	27	LYS	CD-CE	-5.53	1.37	1.51
1	A	28	LYS	CD-CE	-5.49	1.37	1.51
1	B	27	LYS	CG-CD	5.30	1.70	1.52
1	A	17	TYR	CD1-CE1	5.28	1.47	1.39
1	A	7	ASP	CG-OD2	5.26	1.37	1.25
1	A	27	LYS	CE-NZ	5.20	1.62	1.49
1	B	30	LEU	C-O	5.19	1.33	1.23
1	B	11	GLU	CB-CG	5.18	1.61	1.52
1	A	11	GLU	C-O	5.14	1.33	1.23
1	B	6	GLU	CD-OE1	5.14	1.31	1.25
1	A	4	GLN	CD-OE1	5.08	1.35	1.24
1	B	15	LYS	CD-CE	5.05	1.63	1.51
1	A	10	GLU	C-O	5.04	1.32	1.23
1	B	19	ILE	C-O	5.04	1.32	1.23
1	B	5	ILE	CB-CG2	5.03	1.68	1.52
1	B	30	LEU	CG-CD1	5.02	1.70	1.51

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	25	ARG	NE-CZ-NH1	18.39	129.49	120.30
1	A	7	ASP	CB-CG-OD2	-13.91	105.78	118.30
1	A	17	TYR	CB-CG-CD2	-12.40	113.56	121.00
1	A	31	GLY	N-CA-C	-12.29	82.39	113.10
1	A	28	LYS	CD-CE-NZ	-10.54	87.46	111.70
1	A	17	TYR	CD1-CE1-CZ	-8.48	112.17	119.80
1	A	25	ARG	NH1-CZ-NH2	-8.42	110.14	119.40
1	A	25	ARG	CD-NE-CZ	6.96	133.34	123.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	17	TYR	CG-CD2-CE2	-6.59	116.03	121.30
1	A	16	LEU	CA-CB-CG	-6.45	100.46	115.30
1	A	26	ILE	O-C-N	-6.38	112.49	122.70
1	A	10	GLU	OE1-CD-OE2	6.26	130.81	123.30
1	A	7	ASP	OD1-CG-OD2	6.24	135.16	123.30
1	A	17	TYR	CE1-CZ-CE2	6.23	129.76	119.80
1	A	14	SER	CA-CB-OG	-6.02	94.96	111.20
1	B	27	LYS	CG-CD-CE	-5.84	94.38	111.90
1	B	30	LEU	CB-CG-CD1	-5.63	101.42	111.00
1	B	5	ILE	CA-CB-CG1	-5.58	100.39	111.00
1	A	12	ILE	CA-CB-CG1	-5.57	100.41	111.00
1	B	30	LEU	CB-CG-CD2	5.56	120.45	111.00
1	A	17	TYR	CE1-CZ-OH	-5.49	105.29	120.10
1	A	16	LEU	N-CA-CB	5.16	120.72	110.40
1	A	19	ILE	CB-CG1-CD1	-5.14	99.52	113.90
1	A	29	LEU	CD1-CG-CD2	5.09	125.77	110.50
1	B	32	GLU	OE1-CD-OE2	-5.09	117.20	123.30
1	A	9	LEU	CB-CG-CD1	-5.04	102.44	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	253	0	277	7	0
1	B	255	0	266	3	0
2	A	19	0	0	5	0
2	B	31	0	0	2	0
All	All	558	0	543	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:4:GLN:CD	1:A:4:GLN:CG	1.75	1.50
1:A:27:LYS:HE2	2:A:2019:HOH:O	1.71	0.89
1:A:27:LYS:CE	2:A:2019:HOH:O	2.24	0.84
1:B:22:GLU:OE1	2:B:2024:HOH:O	2.04	0.76
1:A:27:LYS:NZ	2:A:2019:HOH:O	2.26	0.69
1:A:17:TYR:OH	2:A:2012:HOH:O	2.11	0.68
1:B:18:HIS:NE2	2:B:2018:HOH:O	2.28	0.66
1:A:6:GLU:OE1	2:A:2003:HOH:O	2.17	0.58
1:A:4:GLN:CD	1:A:4:GLN:CB	2.71	0.55
1:B:18:HIS:O	1:B:22:GLU:HG3	2.16	0.46

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	29/34 (85%)	29 (100%)	0	0	100	100
1	B	30/34 (88%)	30 (100%)	0	0	100	100
All	All	59/68 (87%)	59 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	28/31 (90%)	28 (100%)	0	100	100
1	B	27/31 (87%)	26 (96%)	1 (4%)	34	15
All	All	55/62 (89%)	54 (98%)	1 (2%)	59	43

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	8	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	31/34 (91%)	0.15	0 100 100	18, 26, 39, 42	0
1	B	32/34 (94%)	0.18	0 100 100	18, 27, 37, 44	0
All	All	63/68 (92%)	0.17	0 100 100	18, 27, 39, 44	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.