

# Full wwPDB X-ray Structure Validation Report (i)

### Nov 11, 2024 - 09:53 PM EST

PDB ID	:	2GEF
Title	:	Crystal structure of a Novel viral protease with a serine/lysine catalytic dyad
		mechanism
Authors	:	Paetzel, M.; Feldman, A.R.; Lee, J.; Delmas, B.
Deposited on	:	2006-03-20
Resolution	:	2.20  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of	of chain		
1	А	217	54%	30%	7%	9%
1	В	217	53%	32%	6%	6%



# 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 3285 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
1	Λ	108	Total	С	Ν	0	S	Se	0	0	0
1		190	1488	935	258	289	2	4	0	0	0
1	В	202	Total	С	Ν	0	S	Se	0	0	0
	ГБ	203	1519	954	263	296	2	4	0	0	0

• Molecule 1 is a protein called Protease VP4.

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	557	MET	-	initiating methionine	UNP Q8AZM0
А	615	MSE	MET	modified residue	UNP Q8AZM0
А	685	MSE	MET	modified residue	UNP Q8AZM0
А	743	MSE	MET	modified residue	UNP Q8AZM0
А	767	MSE	MET	modified residue	UNP Q8AZM0
В	557	MET	-	initiating methionine	UNP Q8AZM0
В	615	MSE	MET	modified residue	UNP Q8AZM0
В	685	MSE	MET	modified residue	UNP Q8AZM0
В	743	MSE	MET	modified residue	UNP Q8AZM0
В	767	MSE	MET	modified residue	UNP Q8AZM0

• Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	162	Total O 162 162	0	0
2	В	116	Total O 116 116	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Protease VP4



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants	144.07Å 144.07Å 144.07Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
$\mathbf{P}_{\text{acclution}}(\hat{\mathbf{A}})$	29.41 - 2.20	Depositor
Resolution (A)	29.41 - 2.20	EDS
% Data completeness	99.8 (29.41-2.20)	Depositor
(in resolution range)	99.9 (29.41-2.20)	EDS
R <sub>merge</sub>	0.06	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	14.15 (at 2.20Å)	Xtriage
Refinement program	CNS 1.1	Depositor
B B.	0.222 , $0.267$	Depositor
$\Pi, \Pi_{free}$	0.236 , (Not available)	DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor $(Å^2)$	33.9	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}(e/A^3), B_{sol}(A^2)$	0.35 , $38.2$	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.40, < L^2 > = 0.22$	Xtriage
Estimated twinning fraction	0.329 for -l,-k,-h	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3285	wwPDB-VP
Average B, all atoms $(Å^2)$	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.87% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	1.71	19/1516~(1.3%)	1.48	18/2060~(0.9%)	
1	В	1.43	10/1547~(0.6%)	1.42	21/2104 (1.0%)	
All	All	1.58	29/3063~(0.9%)	1.45	39/4164~(0.9%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	2

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
1	А	570	TYR	CZ-OH	8.27	1.51	1.37
1	В	597	ASN	N-CA	7.15	1.60	1.46
1	А	602	GLU	CB-CG	-7.09	1.38	1.52
1	А	591	VAL	CB-CG1	6.89	1.67	1.52
1	В	770	THR	CB-OG1	6.38	1.56	1.43
1	В	735	LYS	CD-CE	6.17	1.66	1.51
1	В	601	SER	CB-OG	-6.13	1.34	1.42
1	В	770	THR	C-O	6.08	1.34	1.23
1	А	695	ILE	CA-CB	-6.03	1.41	1.54
1	В	604	TYR	CZ-OH	5.88	1.47	1.37
1	А	711	TYR	CD2-CE2	5.63	1.47	1.39
1	А	601	SER	CB-OG	-5.60	1.34	1.42
1	А	669	PHE	CE2-CZ	5.51	1.47	1.37
1	В	763	ALA	CA-CB	-5.46	1.41	1.52
1	A	705	PHE	CD2-CE2	5.45	1.50	1.39
1	A	607	VAL	CB-CG1	5.38	1.64	1.52
1	А	581	VAL	CB-CG2	5.36	1.64	1.52
1	А	690	GLY	C-O	5.32	1.32	1.23



Mol	Chain	Res	Type	Atoms	Ζ	$\operatorname{Observed}(\operatorname{\AA})$	$\mathrm{Ideal}(\mathrm{\AA})$
1	В	587	ALA	CA-CB	5.31	1.63	1.52
1	А	720	VAL	CB-CG2	5.26	1.63	1.52
1	А	604	TYR	CZ-OH	5.25	1.46	1.37
1	А	656	THR	CB-CG2	5.23	1.69	1.52
1	А	632	VAL	CB-CG1	5.23	1.63	1.52
1	А	760	ALA	CA-CB	5.19	1.63	1.52
1	А	716	SER	CB-OG	-5.18	1.35	1.42
1	А	623	TYR	CG-CD2	-5.12	1.32	1.39
1	В	589	PHE	CE1-CZ	5.07	1.47	1.37
1	B	602	GLU	CB-CG	-5.05	1.42	1.52
1	А	600	THR	CA-CB	5.05	1.66	1.53

Continued from previous page...

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	620	ARG	NE-CZ-NH2	10.24	125.42	120.30
1	А	671	ARG	NE-CZ-NH2	-8.80	115.90	120.30
1	В	752	ARG	NE-CZ-NH2	-8.59	116.00	120.30
1	А	574	LEU	CA-CB-CG	8.29	134.36	115.30
1	А	731	ARG	NE-CZ-NH2	-7.99	116.31	120.30
1	В	608	ARG	NE-CZ-NH1	7.88	124.24	120.30
1	В	671	ARG	NE-CZ-NH1	7.84	124.22	120.30
1	В	608	ARG	NE-CZ-NH2	-7.56	116.52	120.30
1	А	608	ARG	NE-CZ-NH1	-7.47	116.56	120.30
1	В	769	SER	N-CA-C	7.37	130.91	111.00
1	В	770	THR	N-CA-C	7.08	130.13	111.00
1	А	608	ARG	NE-CZ-NH2	6.93	123.77	120.30
1	А	585	ASP	CB-CG-OD2	6.89	124.50	118.30
1	В	560	LEU	CB-CG-CD2	6.68	122.36	111.00
1	В	671	ARG	NE-CZ-NH2	-6.20	117.20	120.30
1	В	596	ASP	C-N-CA	-6.17	106.28	121.70
1	В	715	LEU	CB-CG-CD2	5.94	121.10	111.00
1	В	565	LEU	CB-CG-CD2	5.89	121.01	111.00
1	А	683	ASN	N-CA-C	-5.84	95.24	111.00
1	В	560	LEU	N-CA-C	-5.81	95.30	111.00
1	В	596	ASP	CB-CG-OD1	5.81	123.53	118.30
1	В	628	ASP	CB-CG-OD1	5.79	123.51	118.30
1	А	746	ASP	CB-CG-OD2	5.76	123.48	118.30
1	А	671	ARG	NE-CZ-NH1	5.63	123.12	120.30
1	В	619	VAL	CG1-CB-CG2	-5.63	101.90	110.90
1	А	758	ASP	CB-CG-OD2	-5.62	113.24	118.30
1	A	574	LEU	CB-CG-CD2	-5.60	101.47	111.00



Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	А	620	ARG	NE-CZ-NH1	-5.56	117.52	120.30
1	В	596	ASP	O-C-N	-5.50	113.90	122.70
1	А	750	ASP	CB-CG-OD2	5.49	123.24	118.30
1	А	564	LEU	CB-CG-CD2	-5.44	101.75	111.00
1	В	560	LEU	CB-CA-C	5.36	120.39	110.20
1	В	596	ASP	CB-CG-OD2	-5.28	113.55	118.30
1	А	754	CYS	CA-CB-SG	-5.23	104.59	114.00
1	В	660	LEU	CA-CB-CG	5.14	127.13	115.30
1	В	757	ASP	CB-CG-OD1	5.12	122.91	118.30
1	А	612	ARG	NE-CZ-NH1	-5.12	117.74	120.30
1	В	697	LEU	CA-CB-CG	5.07	126.95	115.30
1	А	562	ILE	N-CA-C	-5.05	97.36	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	636	THR	Peptide
1	В	769	SER	Peptide

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1488	0	1434	50	0
1	В	1519	0	1461	70	0
2	А	162	0	0	8	0
2	В	116	0	0	9	0
All	All	3285	0	2895	110	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All (110) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



20FF
2 GET

Atom 1	Atom 2	Interatomic	Clash	
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)	
1:B:573:PRO:HG2	1:B:602:GLU:HG3	1.31	1.12	
1:B:722:PRO:HB3	1:B:743:MSE:HE2	1.31	1.10	
1:A:565:LEU:HD22	1:B:565:LEU:CD2	1.86	1.05	
1:A:565:LEU:HD22	1:B:565:LEU:HD22	1.38	1.05	
1:A:752:ARG:NH2	2:A:216:HOH:O	1.89	1.05	
1:B:611:HIS:HB3	1:B:614:LEU:HD12	1.34	1.04	
1:B:584:THR:HA	1:B:675:SER:HB2	1.40	1.02	
1:A:596:ASP:O	1:A:597:ASN:HB3	1.74	0.88	
1:B:656:THR:O	1:B:657:ARG:HG2	1.77	0.83	
1:B:727:ASN:OD1	2:B:235:HOH:O	1.97	0.81	
1:B:559:ASP:HB3	1:B:560:LEU:HD23	1.62	0.80	
1:B:596:ASP:O	1:B:597:ASN:CB	2.21	0.79	
1:A:565:LEU:HD22	1:B:565:LEU:HD21	1.67	0.75	
1:B:668:VAL:HG13	1:B:678:VAL:HG22	1.67	0.75	
1:B:671:ARG:NH2	1:B:677:ASP:OD1	2.20	0.74	
1:B:573:PRO:HG2	1:B:602:GLU:CG	2.14	0.74	
1:B:596:ASP:O	1:B:597:ASN:HB3	1.87	0.73	
1:A:745:GLN:HE22	1:B:597:ASN:HA	1.54	0.72	
1:A:596:ASP:O	1:A:597:ASN:CB	2.32	0.72	
1:A:560:LEU:HB2	1:A:561:PRO:HD2	1.70	0.71	
1:A:580:ILE:HD13	1:A:580:ILE:N	2.05	0.71	
1:B:584:THR:HA	1:B:675:SER:CB	2.19	0.71	
1:B:672:ALA:O	2:B:234:HOH:O	2.09	0.69	
1:B:629:ASP:O	2:B:259:HOH:O	2.11	0.68	
1:A:651:LYS:CB	2:A:52:HOH:O	2.43	0.67	
1:A:615:MSE:HE2	1:A:619:VAL:HG12	1.75	0.67	
1:A:615:MSE:CE	1:A:619:VAL:HG12	2.25	0.65	
1:A:560:LEU:CD2	2:A:202:HOH:O	2.44	0.65	
1:B:722:PRO:CB	1:B:743:MSE:HE2	2.19	0.64	
1:B:665:VAL:HG11	1:B:678:VAL:HG11	1.79	0.64	
1:A:670:TRP:CE2	1:A:708:LYS:HG2	2.35	0.62	
1:A:684:ASP:N	2:A:237:HOH:O	2.07	0.62	
1:A:692:SER:HB3	1:A:723:VAL:HG21	1.82	0.62	
1:B:614:LEU:HD11	2:B:146:HOH:O	2.00	0.61	
1:B:711:TYR:HA	1:B:741:VAL:O	2.00	0.60	
1:B:714:CYS:HB3	2:B:64:HOH:O	2.01	0.60	
1:A:565:LEU:CD2	1:B:565:LEU:HD22	2.25	0.60	
1:B:612:ARG:O	1:B:620:ARG:NE	2.33	0.60	
1:B:586:GLY:HA2	1:B:700:ALA:HB1	1.84	0.59	
1:B:584:THR:OG1	1:B:585:ASP:OD1	2.21	0.58	
1:B:611:HIS:CB	1:B:614:LEU:HD12	2.23	0.58	
1:B:772:THR:O	1:B:773:ARG:CB	2.51	0.58	



		Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:B:771:VAL:HG12	1:B:772:THR:H	1.69	0.58	
1:B:722:PRO:HB3	1:B:743:MSE:CE	2.21	0.57	
1:A:674:GLY:HA2	1:A:767:MSE:CE	2.34	0.56	
1:B:656:THR:C	1:B:658:THR:H	2.08	0.56	
1:B:574:LEU:O	2:B:62:HOH:O	2.17	0.56	
1:A:612:ARG:O	1:A:620:ARG:HD3	2.06	0.55	
1:B:630:HIS:HB3	1:B:658:THR:O	2.06	0.55	
1:A:741:VAL:HG21	1:A:762:HIS:CE1	2.41	0.55	
1:A:571:LYS:HB2	1:A:598:HIS:HD2	1.73	0.54	
1:B:570:TYR:CE1	1:B:601:SER:HB3	2.43	0.54	
1:B:612:ARG:HA	1:B:615:MSE:SE	2.58	0.54	
1:A:668:VAL:HG13	1:A:678:VAL:HG22	1.90	0.54	
1:A:566:GLN:HE22	1:B:728:LEU:H	1.54	0.54	
1:B:749:GLU:O	1:B:749:GLU:HG2	2.07	0.53	
1:B:604:TYR:OH	1:B:683:ASN:ND2	2.26	0.52	
1:B:771:VAL:HG21	2:B:234:HOH:O	2.10	0.52	
1:A:566:GLN:NE2	1:B:728:LEU:H	2.08	0.52	
1:B:617:PRO:HA	1:B:620:ARG:HG3	1.92	0.51	
1:B:670:TRP:NE1	1:B:706:VAL:O	2.39	0.51	
1:B:656:THR:C	1:B:658:THR:N	2.65	0.50	
1:B:570:TYR:HE1	1:B:601:SER:HB3	1.76	0.50	
1:A:573:PRO:HG2	1:A:602:GLU:HG3	1.93	0.49	
1:A:615:MSE:HB2	1:A:620:ARG:HG2	1.93	0.49	
1:B:605:ILE:HG23	1:B:605:ILE:O	2.13	0.49	
1:B:715:LEU:HD21	1:B:756:LEU:HD13	1.94	0.49	
1:B:657:ARG:HH22	1:B:702:SER:HB3	1.77	0.49	
1:B:727:ASN:HB2	1:B:746:ASP:OD2	2.12	0.49	
1:A:561:PRO:HD3	1:B:571:LYS:HE2	1.95	0.49	
1:B:767:MSE:O	1:B:768:GLU:C	2.51	0.49	
1:B:631:LYS:HD2	1:B:633:TRP:CZ2	2.48	0.48	
1:B:671:ARG:NH1	2:B:75:HOH:O	2.34	0.48	
1:B:665:VAL:CG1	1:B:678:VAL:HG11	2.42	0.48	
1:A:561:PRO:HG3	2:B:80:HOH:O	2.14	0.48	
1:A:619:VAL:O	1:A:619:VAL:CG1	2.61	0.47	
1:A:685:MSE:HE2	2:A:9:HOH:O	2.14	0.47	
1:B:586:GLY:HA2	1:B:700:ALA:CB	2.45	0.47	
1:A:573:PRO:HG2	1:A:602:GLU:CD	2.36	0.46	
1:B:722:PRO:HA	1:B:743:MSE:HB3	1.98	0.45	
1:B:616:SER:HA	1:B:617:PRO:HD2	1.77	0.45	
1:A:580:ILE:N	1:A:580:ILE:CD1	2.74	0.45	
1:A:589:PHE:HB3	1:A:696:ALA:HB2	1.99	0.45	



Atom 1	Atom 2	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:A:573:PRO:HG2	1:A:602:GLU:CG	2.46	0.44	
1:A:582:HIS:CE1	1:A:675:SER:HB2	2.51	0.44	
1:B:719:ASN:HD22	1:B:719:ASN:HA	1.57	0.43	
1:A:560:LEU:HD21	2:A:202:HOH:O	2.13	0.43	
1:A:735:LYS:NZ	2:A:190:HOH:O	2.50	0.43	
1:A:560:LEU:CB	1:A:561:PRO:HD2	2.44	0.43	
1:B:769:SER:C	1:B:770:THR:HG22	2.38	0.43	
1:A:746:ASP:OD2	1:B:566:GLN:OE1	2.37	0.43	
1:A:560:LEU:HD23	1:A:560:LEU:H	1.82	0.43	
1:A:610:ASP:O	1:A:610:ASP:OD1	2.36	0.43	
1:B:597:ASN:O	1:B:598:HIS:HB3	2.19	0.43	
1:A:626:THR:HA	1:A:652:PHE:HB2	2.01	0.42	
1:A:619:VAL:HG11	2:A:1:HOH:O	2.20	0.42	
1:A:745:GLN:NE2	1:B:597:ASN:HA	2.26	0.42	
1:B:585:ASP:OD1	1:B:585:ASP:N	2.52	0.42	
1:A:582:HIS:HE1	1:A:675:SER:OG	2.02	0.42	
1:B:630:HIS:CB	1:B:658:THR:O	2.67	0.41	
1:A:734:HIS:NE2	1:A:750:ASP:OD2	2.35	0.41	
1:A:683:ASN:OD1	1:A:683:ASN:C	2.58	0.41	
1:B:601:SER:OG	1:B:732:ALA:HB2	2.20	0.41	
1:B:655:LYS:HZ3	1:B:655:LYS:HG3	1.70	0.41	
1:A:590:PRO:HB2	1:A:729:LYS:HG2	2.02	0.41	
1:A:657:ARG:NH1	1:A:657:ARG:HG3	2.36	0.41	
1:B:703:LEU:HA	1:B:704:PRO:HD3	1.98	0.41	
1:B:605:ILE:O	1:B:605:ILE:CG2	2.69	0.40	
1:B:712:THR:O	1:B:742:GLY:HA3	2.22	0.40	
1:A:743:MSE:HE3	1:A:752:ARG:CZ	2.52	0.40	

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Pere	centil	$\mathbf{les}$
1	А	194/217~(89%)	182 (94%)	9~(5%)	3~(2%)	8	6	
1	В	199/217~(92%)	181 (91%)	13~(6%)	5(2%)	4	2	
All	All	393/434 (91%)	363 (92%)	22 (6%)	8 (2%)	6	4	

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	768	GLU
1	В	768	GLU
1	В	769	SER
1	В	771	VAL
1	А	597	ASN
1	В	597	ASN
1	А	672	ALA
1	В	612	ARG

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	158/173~(91%)	145~(92%)	13 (8%)	99		
1	В	161/173~(93%)	147 (91%)	14 (9%)	8 8		
All	All	319/346~(92%)	292 (92%)	27 (8%)	8 9		

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	560	LEU
1	А	563	SER
1	А	572	GLN
1	А	574	LEU
1	А	576	ARG
1	А	579	ARG
1	А	580	ILE
1	А	581	VAL



Mol	Chain	Res	Type
1	А	598	HIS
1	А	610	ASP
1	А	657	ARG
1	А	704	PRO
1	А	768	GLU
1	В	560	LEU
1	В	565	LEU
1	В	576	ARG
1	В	579	ARG
1	В	581	VAL
1	В	585	ASP
1	В	610	ASP
1	В	618	ASP
1	В	619	VAL
1	В	622	SER
1	В	657	ARG
1	В	660	LEU
1	В	724	GLN
1	В	769	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	566	GLN
1	А	582	HIS
1	А	598	HIS
1	А	719	ASN
1	А	745	GLN
1	В	566	GLN
1	В	572	GLN
1	В	582	HIS
1	В	598	HIS
1	В	719	ASN

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



# 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

# 5.6 Ligand geometry (i)

There are no ligands in this entry.

# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

# 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSR	Z>2	$OWAB(Å^2)$	Q<0.9
1	А	194/217~(89%)	-1.50	0 100	100	6, 12, 24, 35	0
1	В	199/217~(91%)	-1.24	0 100	100	7, 24, 42, 48	0
All	All	393/434~(90%)	-1.36	0 100	100	6, 19, 39, 48	0

There are no RSRZ outliers to report.

# 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

There are no ligands in this entry.

### 6.5 Other polymers (i)

There are no such residues in this entry.

