



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 2, 2024 – 11:56 AM EST

PDB ID : 2HEY  
Title : Crystal structure of murine OX40L bound to human OX40  
Authors : Hymowitz, S.G.  
Deposited on : 2006-06-22  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.21  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.004 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	F	152	
1	G	152	
2	R	146	
2	T	146	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4356 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tumor necrosis factor ligand superfamily member 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	F	135	1077	686	181	204	6	1	0	0
1	G	134	1071	683	180	202	6	10	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	47	GLY	-	cloning artifact	UNP P43488
F	48	SER	-	cloning artifact	UNP P43488
F	49	HIS	-	cloning artifact	UNP P43488
F	50	MET	-	cloning artifact	UNP P43488
G	47	GLY	-	cloning artifact	UNP P43488
G	48	SER	-	cloning artifact	UNP P43488
G	49	HIS	-	cloning artifact	UNP P43488
G	50	MET	-	cloning artifact	UNP P43488

- Molecule 2 is a protein called Tumor necrosis factor receptor superfamily member 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	R	138	1025	613	195	198	19	4	0	0
2	T	135	1002	599	189	195	19	18	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	25	GLY	-	cloning artifact	UNP P43489
R	26	SER	-	cloning artifact	UNP P43489
R	27	HIS	-	cloning artifact	UNP P43489
R	28	MET	-	cloning artifact	UNP P43489

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
T	25	GLY	-	cloning artifact	UNP P43489
T	26	SER	-	cloning artifact	UNP P43489
T	27	HIS	-	cloning artifact	UNP P43489
T	28	MET	-	cloning artifact	UNP P43489

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	R	1	5	4	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	F	74	Total	O	0	0
			74	74		
4	G	42	Total	O	0	0
			42	42		
4	R	43	Total	O	0	0
			43	43		
4	T	17	Total	O	0	0
			17	17		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tumor necrosis factor ligand superfamily member 4




- Molecule 1: Tumor necrosis factor ligand superfamily member 4




- Molecule 2: Tumor necrosis factor receptor superfamily member 4



- Molecule 2: Tumor necrosis factor receptor superfamily member 4





ASP

## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.73Å 104.73Å 478.16Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.00 30.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.8 (30.00-2.00) 98.8 (30.00-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.06 (at 2.01Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.221 , 0.247 0.224 , 0.252	Depositor DCC
$R_{free}$ test set	6853 reflections (10.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.6	Xtrriage
Anisotropy	0.063	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 46.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	4356	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	F	0.48	1/1100 (0.1%)	0.60	0/1493
1	G	0.36	0/1094	0.58	0/1485
2	R	0.38	0/1051	0.57	0/1430
2	T	0.36	0/1026	0.59	2/1396 (0.1%)
All	All	0.40	1/4271 (0.0%)	0.58	2/5804 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	191	SER	CB-OG	-8.54	1.31	1.42

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	T	138	ASN	N-CA-CB	8.90	126.62	110.60
2	T	139	GLN	N-CA-CB	5.18	119.93	110.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1077	0	1057	9	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	1071	0	1052	5	0
2	R	1025	0	953	2	0
2	T	1002	0	931	5	0
3	R	5	0	0	0	0
4	F	74	0	0	3	0
4	G	42	0	0	0	0
4	R	43	0	0	0	0
4	T	17	0	0	0	0
All	All	4356	0	3993	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:116:ILE:HD12	1:G:138:ILE:HD11	1.69	0.75
2:T:39:ASN:HD22	2:T:41:ARG:HH12	1.41	0.69
1:F:116:ILE:HD12	1:F:138:ILE:CD1	2.27	0.65
1:F:131:MET:HG3	4:F:244:HOH:O	2.00	0.60
1:F:75:LEU:HD11	1:F:116:ILE:HD13	1.84	0.60
1:F:137:ARG:HD3	4:F:219:HOH:O	2.08	0.54
2:R:84:CYS:SG	2:R:99:CYS:CB	2.97	0.52
2:T:84:CYS:HG	2:T:99:CYS:CB	2.23	0.52
1:G:75:LEU:HD11	1:G:116:ILE:HD13	1.93	0.51
2:R:84:CYS:HG	2:R:99:CYS:CB	2.17	0.50
2:T:39:ASN:HD22	2:T:41:ARG:NH1	2.08	0.50
1:F:137:ARG:CD	4:F:219:HOH:O	2.60	0.49
2:T:147:CYS:HG	2:T:166:CYS:HG	1.64	0.46
1:F:103:ILE:HG13	1:F:178:LEU:HD11	1.99	0.45
1:F:116:ILE:HD12	1:F:138:ILE:HD11	1.97	0.44
1:F:148:PHE:O	1:F:149:LYS:HB2	2.18	0.44
1:G:63:LEU:HD11	1:G:89:VAL:HG23	2.00	0.43
1:G:148:PHE:O	1:G:149:LYS:HB2	2.20	0.42
1:G:116:ILE:HD12	1:G:138:ILE:CD1	2.47	0.40
2:T:132:HIS:ND1	2:T:143:PRO:HA	2.36	0.40
1:F:57:ASP:HB3	1:F:58:PRO:HD3	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	133/152 (88%)	130 (98%)	3 (2%)	0	100	100
1	G	132/152 (87%)	130 (98%)	2 (2%)	0	100	100
2	R	136/146 (93%)	133 (98%)	2 (2%)	1 (1%)	19	14
2	T	131/146 (90%)	128 (98%)	2 (2%)	1 (1%)	16	12
All	All	532/596 (89%)	521 (98%)	9 (2%)	2 (0%)	30	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	R	149	LEU
2	T	149	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	122/137 (89%)	116 (95%)	6 (5%)	21	18
1	G	121/137 (88%)	118 (98%)	3 (2%)	42	45
2	R	120/127 (94%)	118 (98%)	2 (2%)	56	61
2	T	118/127 (93%)	116 (98%)	2 (2%)	56	61
All	All	481/528 (91%)	468 (97%)	13 (3%)	40	42

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	63	LEU
1	F	69	ARG
1	F	131	MET
1	F	134	ASP
1	F	137	ARG
1	F	157	ASN
1	G	62	ARG
1	G	125	ASN
1	G	178	LEU
2	R	58	ARG
2	R	149	LEU
2	T	41	ARG
2	T	88	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	74	GLN
1	F	157	ASN
1	G	74	GLN
1	G	119	HIS
1	G	167	GLN
2	T	39	ASN
2	T	88	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	R	171	-	4,4,4	0.29	0	6,6,6	0.12	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

**Warning:** The R factor obtained from EDS is 0.2708, which does not match the depositor's R factor of 0.22054. Please interpret the results in this section carefully.

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	F	135/152 (88%)	1.04	27 (20%) <b>3</b> <b>3</b>	27, 34, 52, 65	1 (0%)
1	G	134/152 (88%)	1.43	31 (23%) <b>2</b> <b>2</b>	16, 36, 52, 64	2 (1%)
2	R	138/146 (94%)	1.35	37 (26%) <b>2</b> <b>2</b>	22, 36, 45, 51	1 (0%)
2	T	135/146 (92%)	2.66	83 (61%) <b>0</b> <b>1</b>	14, 37, 69, 76	4 (2%)
All	All	542/596 (90%)	1.62	178 (32%) <b>1</b> <b>1</b>	14, 36, 55, 76	8 (1%)

All (178) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	T	123	VAL	10.2
2	T	122	GLY	7.9
2	T	119	TYR	7.6
1	G	161	THR	7.2
2	T	135	PRO	6.8
2	T	120	LYS	6.3
2	R	117	ASP	6.2
2	T	29	LEU	5.8
2	T	136	GLY	5.5
1	G	162	LEU	5.2
2	T	148	THR	5.2
2	R	40	ASP	5.2
2	T	111	ALA	5.2
2	T	116	LEU	5.1
2	T	97	GLN	5.0
1	F	135	GLY	4.9
1	F	57	ASP	4.9
2	T	110	ARG	4.9
2	T	118	SER	4.7

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	F	161	THR	4.7
1	F	134	ASP	4.6
1	G	160	ASP	4.6
2	T	138	ASN	4.6
2	T	99	CYS	4.5
2	R	148	THR	4.4
2	T	130	PRO	4.4
2	R	111	ALA	4.4
2	T	133	PHE	4.4
2	T	109	CYS	4.3
2	R	109	CYS	4.3
2	T	121	PRO	4.2
1	G	157	ASN	4.2
2	T	106	VAL	4.2
1	F	190	HIS	4.1
1	F	191	SER	4.1
2	R	112	GLY	4.0
2	T	139	GLN	3.9
2	T	98	LEU	3.9
2	T	149	LEU	3.9
2	T	67	CYS	3.9
1	G	159	PRO	3.9
2	T	108	ARG	3.9
2	T	124	ASP	3.8
1	G	122	GLU	3.8
2	R	140	ALA	3.8
2	T	47	ARG	3.7
1	F	132	LEU	3.7
2	T	87	CYS	3.7
2	T	92	GLY	3.7
2	T	112	GLY	3.7
2	R	130	PRO	3.7
2	R	119	TYR	3.7
2	T	86	TRP	3.7
2	R	146	ASN	3.7
2	T	127	PRO	3.6
1	G	165	HIS	3.6
2	T	125	CYS	3.6
1	G	84	TYR	3.6
2	T	117	ASP	3.6
2	R	127	PRO	3.6
2	R	123	VAL	3.6

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	G	70	CYS	3.6
2	T	113	THR	3.6
2	T	57	SER	3.5
2	T	141	CYS	3.5
1	G	163	CYS	3.5
2	T	90	ARG	3.5
2	R	122	GLY	3.4
2	R	90	ARG	3.4
2	T	89	LEU	3.4
2	T	165	ILE	3.3
2	R	135	PRO	3.3
2	T	150	ALA	3.3
1	F	189	TYR	3.3
2	T	81	CYS	3.3
2	T	101	ALA	3.3
1	G	57	ASP	3.3
2	R	110	ARG	3.3
2	R	128	CYS	3.2
2	T	75	VAL	3.2
2	R	129	PRO	3.2
2	T	36	TYR	3.2
1	G	132	LEU	3.2
1	G	113	GLU	3.2
1	F	160	ASP	3.1
1	F	162	LEU	3.1
2	T	58	ARG	3.1
1	G	58	PRO	3.1
1	F	170	ASP	3.1
2	R	134	SER	3.1
2	T	59	SER	3.1
2	R	131	GLY	3.0
1	F	157	ASN	3.0
1	G	134	ASP	3.0
2	T	94	GLU	3.0
2	T	134	SER	3.0
2	R	133	PHE	3.0
2	T	100	THR	2.9
1	G	158	ALA	2.9
2	T	140	ALA	2.9
2	T	46	CYS	2.8
1	F	90	GLN	2.8
2	R	116	LEU	2.8

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	T	129	PRO	2.7
1	F	163	CYS	2.7
2	R	113	THR	2.7
1	F	188	SER	2.7
1	F	122	GLU	2.7
1	G	125	ASN	2.6
2	T	128	CYS	2.6
2	T	126	ALA	2.6
2	T	107	CYS	2.6
1	G	190	HIS	2.6
2	T	30	HIS	2.6
2	T	142	LYS	2.6
1	G	59	PRO	2.5
1	G	60	ILE	2.5
1	G	187	GLY	2.5
2	T	84	CYS	2.5
2	T	65	ARG	2.5
1	F	186	GLU	2.5
2	T	164	ALA	2.5
1	F	156	VAL	2.5
2	T	53	VAL	2.5
2	T	145	THR	2.5
2	T	160	ASN	2.5
1	G	136	ARG	2.4
1	F	68	THR	2.4
2	T	88	ASN	2.4
1	G	176	VAL	2.4
1	G	189	TYR	2.4
2	R	166	CYS	2.4
2	T	73	ASN	2.4
2	R	30	HIS	2.4
2	R	132	HIS	2.4
1	F	131	MET	2.4
1	F	187	GLY	2.4
2	R	141	CYS	2.4
2	T	144	TRP	2.4
1	G	180	PRO	2.4
2	T	93	SER	2.4
1	G	90	GLN	2.4
2	R	137	ASP	2.4
1	F	71	GLU	2.3
2	R	126	ALA	2.3

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	F	70	CYS	2.3
2	T	115	PRO	2.3
2	T	33	GLY	2.3
2	R	73	ASN	2.3
1	G	86	THR	2.3
2	T	56	CYS	2.3
1	F	72	ASP	2.3
2	T	69	PRO	2.2
2	T	70	GLY	2.2
1	F	133	ASN	2.2
2	R	139	GLN	2.2
2	T	105	THR	2.2
2	T	147	CYS	2.2
2	T	96	LYS	2.2
1	F	136	ARG	2.2
1	G	185	PRO	2.2
2	T	159	SER	2.1
1	G	105	TYR	2.1
2	R	65	ARG	2.1
1	F	185	PRO	2.1
2	R	138	ASN	2.1
1	G	183	CYS	2.1
2	T	103	GLN	2.1
2	T	166	CYS	2.1
1	G	188	SER	2.1
2	T	104	ASP	2.1
2	R	150	ALA	2.1
2	T	114	GLN	2.1
2	R	58	ARG	2.1
2	T	137	ASP	2.1
2	R	153	HIS	2.0
2	R	162	SER	2.0
2	T	31	CYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	R	171	5/5	0.98	0.14	27,34,37,39	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.