



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 25, 2024 – 04:33 PM EST

PDB ID : 2HOR  
Title : Crystal structure of alliinase from garlic- apo form  
Authors : Shimon, L.J.W.; Rabinkov, A.; Wilcheck, M.; Mirelman, D.; Frolow, F.  
Deposited on : 2006-07-16  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.21  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.004 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4274 (1.60-1.60)
Clashscore	180529	4682 (1.60-1.60)
Ramachandran outliers	177936	4583 (1.60-1.60)
Sidechain outliers	177891	4582 (1.60-1.60)
RSRZ outliers	164620	4272 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	427	
2	B	6	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BMA	B	3	X	-	-	-

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 4122 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alliin lyase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	425	3506	2242	581	657	26	6	15	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	176	ASP	ASN	SEE REMARK 999	UNP Q01594

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(2-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	6	71	40	2	29	0	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
3	A	1	14	8	1	5	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O S		
4	A	1	5	4 1	0	0
4	A	1	5	4 1	0	0
4	A	1	5	4 1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	2	Total Cl 2 2	0	0

- Molecule 7 is NITRATE ION (three-letter code: NO3) (formula:  $NO_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	N	O	0	0
			4	1	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	496	Total	O	0	0
			496	496		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alliin lyase 1



- Molecule 2: alpha-D-mannopyranose-(2-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.12Å 81.12Å 164.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.60 – 1.60 39.60 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (39.60-1.60) 99.8 (39.60-1.60)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.47 (at 1.60Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.163 , 0.179 0.234 , 0.243	Depositor DCC
$R_{free}$ test set	3663 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.4	Xtrriage
Anisotropy	0.361	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 47.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4122	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.14% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, CL, NO3, SO4, BMA, MAN, NAG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.60	2/3640 (0.1%)	0.69	5/4922 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	55[A]	GLN	N-CA	-13.55	1.19	1.46
1	A	55[B]	GLN	N-CA	-13.55	1.19	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	55[A]	GLN	CA-C-N	7.45	131.10	116.20
1	A	55[B]	GLN	CA-C-N	7.45	131.10	116.20
1	A	55[A]	GLN	CA-C-O	-7.26	104.85	120.10
1	A	55[B]	GLN	CA-C-O	-7.26	104.85	120.10
1	A	54	ILE	C-N-CA	5.53	135.52	121.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3506	0	3438	20	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	71	0	61	0	0
3	A	14	0	13	0	0
4	A	25	0	0	0	0
5	A	4	0	3	0	0
6	A	2	0	0	0	0
7	A	4	0	0	0	0
8	A	496	0	0	5	0
All	All	4122	0	3515	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:354[A]:ARG:NH2	8:A:1172:HOH:O	2.26	0.66
1:A:2[B]:MET:HA	1:A:2[B]:MET:HE3	1.78	0.63
1:A:294:LEU:HG	1:A:298:LYS:HE2	1.82	0.61
1:A:232:HIS:ND1	1:A:318:GLY:HA3	2.18	0.59
1:A:19:ASN:HD22	1:A:21[B]:SER:H	1.51	0.57
1:A:298:LYS:HE3	8:A:899:HOH:O	2.05	0.55
1:A:19:ASN:HD22	1:A:21[A]:SER:H	1.51	0.55
1:A:19:ASN:ND2	1:A:21[B]:SER:H	2.06	0.54
1:A:19:ASN:ND2	1:A:21[A]:SER:H	2.06	0.53
1:A:2[B]:MET:HA	1:A:2[B]:MET:CE	2.39	0.49
1:A:143:LEU:HD22	1:A:221:LYS:HD3	1.97	0.47
1:A:19:ASN:HD22	1:A:19:ASN:C	2.20	0.45
1:A:68:PHE:HB2	1:A:321[A]:LYS:HE2	2.00	0.43
1:A:74:LYS:NZ	8:A:1138:HOH:O	2.33	0.43
1:A:83:LEU:HD13	1:A:83:LEU:C	2.38	0.43
1:A:124:ARG:NH2	1:A:266:LYS:HE2	2.34	0.43
1:A:333:LEU:HD22	1:A:339:PHE:HB2	1.99	0.43
1:A:226[B]:MET:HE1	1:A:240:ALA:HB2	2.02	0.41
1:A:354[B]:ARG:NH1	8:A:946:HOH:O	2.34	0.41
1:A:11[B]:GLU:HG3	8:A:970:HOH:O	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/427 (103%)	426 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	388/375 (104%)	383 (99%)	5 (1%)	65	46

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	LYS
1	A	19	ASN
1	A	25	ARG
1	A	374	LYS
1	A	423	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	2,1	14,14,15	0.91	1 (7%)	17,19,21	0.79	0
2	NAG	B	2	2	14,14,15	0.58	0	17,19,21	0.95	1 (5%)
2	BMA	B	3	2	10,10,12	0.80	0	13,13,17	0.99	1 (7%)
2	MAN	B	4	2	12,12,12	0.54	0	17,17,17	1.41	4 (23%)
2	MAN	B	5	2	11,11,12	0.42	0	15,15,17	0.90	1 (6%)
2	FUC	B	6	2	10,10,11	0.36	0	14,14,16	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	1/1/3/5	0/2/15/22	0/1/1/1
2	MAN	B	4	2	-	2/2/22/22	0/1/1/1
2	MAN	B	5	2	-	0/2/19/22	0/1/1/1
2	FUC	B	6	2	-	-	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	NAG	O5-C1	-2.22	1.40	1.43

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4	MAN	C1-C2-C3	2.78	116.02	110.36
2	B	5	MAN	C1-O5-C5	2.76	115.89	112.19
2	B	4	MAN	C4-C3-C2	2.74	115.64	110.83
2	B	2	NAG	O4-C4-C3	-2.71	103.98	110.38
2	B	4	MAN	O5-C5-C6	2.40	112.39	106.44
2	B	4	MAN	C3-C4-C5	2.34	114.47	110.23
2	B	3	BMA	C6-C5-C4	-2.08	110.68	113.53

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	3	BMA	C3


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	4	MAN	O5-C5-C6-O6
2	B	4	MAN	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	SO4	A	601	-	4,4,4	0.47	0	6,6,6	0.41	0
4	SO4	A	702	-	4,4,4	0.36	0	6,6,6	0.18	0
4	SO4	A	703	-	4,4,4	0.27	0	6,6,6	0.49	0
4	SO4	A	600	-	4,4,4	0.41	0	6,6,6	0.58	0
7	NO3	A	707	-	1,3,3	<b>3.40</b>	<b>1 (100%)</b>	0,3,3	-	-
4	SO4	A	701	-	4,4,4	0.23	0	6,6,6	0.14	0
3	NAG	A	555	1	14,14,15	0.53	0	17,19,21	1.27	<b>1 (5%)</b>
5	ACT	A	602	-	3,3,3	0.88	0	3,3,3	1.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	555	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	707	NO3	O1-N	3.40	1.41	1.24

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	555	NAG	C1-O5-C5	3.63	117.05	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

**Warning:** The R factor obtained from EDS is 0.2407, which does not match the depositor's R factor of 0.163. Please interpret the results in this section carefully.

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	425/427 (99%)	1.38	77 (18%) <b>4</b> <b>3</b>	18, 31, 37, 48	15 (3%)

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	425	LYS	7.9
1	A	424	ALA	4.9
1	A	1	LYS	4.5
1	A	423	LYS	4.5
1	A	56	GLY	4.2
1	A	191	ASN	3.9
1	A	331	ALA	3.8
1	A	55[A]	GLN	3.7
1	A	422	VAL	3.6
1	A	54	ILE	3.4
1	A	356	MET	3.3
1	A	416	TYR	3.3
1	A	193	SER	3.3
1	A	332	LEU	3.2
1	A	355	ARG	3.2
1	A	22	GLU	3.1
1	A	334	ASP	3.1
1	A	336	SER	3.1
1	A	370	TRP	3.1
1	A	19	ASN	3.1
1	A	296	VAL	3.0
1	A	122	LYS	3.0
1	A	335	GLN	3.0
1	A	368	CYS	3.0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	333	LEU	2.9
1	A	152	ASP	2.9
1	A	52	GLU	2.9
1	A	339	PHE	2.9
1	A	57	CYS	2.8
1	A	190	VAL	2.8
1	A	341	TYR	2.8
1	A	123	ASP	2.7
1	A	419	LYS	2.7
1	A	349	TYR	2.6
1	A	337	ASP	2.5
1	A	327	VAL	2.5
1	A	53	LYS	2.5
1	A	218	LYS	2.5
1	A	187	ALA	2.5
1	A	75	GLN	2.5
1	A	371	GLU	2.5
1	A	277	TYR	2.5
1	A	192	VAL	2.4
1	A	344	LEU	2.4
1	A	82	VAL	2.4
1	A	100	PHE	2.3
1	A	149	ALA	2.3
1	A	351	ASN	2.3
1	A	291	LEU	2.3
1	A	294	LEU	2.3
1	A	385	ILE	2.3
1	A	324	GLU	2.3
1	A	6[A]	MET	2.2
1	A	320	LYS	2.2
1	A	61	VAL	2.2
1	A	127	VAL	2.2
1	A	329	ILE	2.2
1	A	374	LYS	2.2
1	A	128	PHE	2.1
1	A	421	MET	2.1
1	A	34	GLU	2.1
1	A	114[A]	GLU	2.1
1	A	133	THR	2.1
1	A	384	ARG	2.1
1	A	381	GLN	2.1
1	A	130	VAL	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	376	CYS	2.1
1	A	33	SER	2.1
1	A	297	LEU	2.1
1	A	238	TYR	2.1
1	A	398	ARG	2.0
1	A	2[A]	MET	2.0
1	A	286	PRO	2.0
1	A	358	PRO	2.0
1	A	330	THR	2.0
1	A	347	SER	2.0
1	A	101	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MAN	B	4	12/12	0.30	0.24	69,76,77,77	0
2	BMA	B	3	10/12	0.72	0.16	48,50,51,53	0
2	MAN	B	5	11/12	0.73	0.16	48,50,50,51	0
2	NAG	B	2	14/15	0.79	0.15	38,40,44,47	0
2	NAG	B	1	14/15	0.85	0.14	30,32,35,36	0
2	FUC	B	6	10/11	0.89	0.14	32,34,35,35	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	SO4	A	701	5/5	0.68	0.17	67,67,68,68	0
3	NAG	A	555	14/15	0.78	0.15	37,42,45,46	0
4	SO4	A	702	5/5	0.82	0.16	32,36,38,40	0
7	NO3	A	707	4/4	0.87	0.15	32,33,33,33	4
4	SO4	A	703	5/5	0.93	0.11	31,31,36,39	0
6	CL	A	706	1/1	0.95	0.13	47,47,47,47	0
4	SO4	A	600	5/5	0.98	0.11	24,24,27,28	0
5	ACT	A	602	4/4	0.98	0.08	19,20,20,21	0
4	SO4	A	601	5/5	0.99	0.07	16,17,18,20	0
6	CL	A	705	1/1	0.99	0.09	18,18,18,18	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.