



# Full wwPDB NMR Structure Validation Report ⓘ

Apr 21, 2024 – 08:42 AM EDT

PDB ID : 2MCI  
BMRB ID : 19440  
Title : NMR structure of DNA duplex  
Authors : Brzezinska, J.; Gdaniec, Z.; Popena, L.; Markiewicz, W.T.  
Deposited on : 2013-08-20

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
wwPDB-RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
wwPDB-ShiftChecker : v1.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2



## 2 Ensemble composition and analysis

This entry contains 10 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

### 3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 504 atoms, of which 182 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called DNA\_(5'-D(\*GP\*TP\*CP\*GP\*GP\*CP\*TP\*G)-3').

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
1	A	8	255	78	92	30	48	7	0

- Molecule 2 is a DNA chain called DNA\_(5'-D(\*CP\*AP\*GP\*CP\*CP\*GP\*AP\*C)-3').

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
2	B	8	249	76	90	32	44	7	0

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble


These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DNA\_(5'-D(\*GP\*TP\*CP\*GP\*GP\*CP\*TP\*G)-3')

Chain A: 



- Molecule 2: DNA\_(5'-D(\*CP\*AP\*GP\*CP\*CP\*GP\*AP\*C)-3')

Chain B: 



### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


#### 4.2.1 Score per residue for model 1

- Molecule 1: DNA\_(5'-D(\*GP\*TP\*CP\*GP\*GP\*CP\*TP\*G)-3')

Chain A: 




- Molecule 2: DNA\_(5'-D(\*CP\*AP\*GP\*CP\*CP\*GP\*AP\*C)-3')

Chain B: 




#### 4.2.2 Score per residue for model 2

- Molecule 1: DNA\_(5'-D(\*GP\*TP\*CP\*GP\*GP\*CP\*TP\*G)-3')

Chain A:  75% 25%




- Molecule 2: DNA\_(5'-D(\*CP\*AP\*GP\*CP\*CP\*GP\*AP\*C)-3')

Chain B:  88% 12%




#### 4.2.3 Score per residue for model 3

- Molecule 1: DNA\_(5'-D(\*GP\*TP\*CP\*GP\*GP\*CP\*TP\*G)-3')

Chain A:  75% 25%




- Molecule 2: DNA\_(5'-D(\*CP\*AP\*GP\*CP\*CP\*GP\*AP\*C)-3')

Chain B:  88% 12%



#### 4.2.4 Score per residue for model 4

- Molecule 1: DNA\_(5'-D(\*GP\*TP\*CP\*GP\*GP\*CP\*TP\*G)-3')

Chain A:  75% 25%




- Molecule 2: DNA\_(5'-D(\*CP\*AP\*GP\*CP\*CP\*GP\*AP\*C)-3')

Chain B:  88% 12%




#### 4.2.5 Score per residue for model 5

- Molecule 1: DNA\_(5'-D(\*GP\*TP\*CP\*GP\*GP\*CP\*TP\*G)-3')

Chain A:  75% 25%




- Molecule 2: DNA\_(5'-D(\*CP\*AP\*GP\*CP\*CP\*GP\*AP\*C)-3')

Chain B:  88% 12%




#### 4.2.6 Score per residue for model 6

- Molecule 1: DNA\_(5'-D(\*GP\*TP\*CP\*GP\*GP\*CP\*TP\*G)-3')

Chain A:  75% 25%




- Molecule 2: DNA\_(5'-D(\*CP\*AP\*GP\*CP\*CP\*GP\*AP\*C)-3')

Chain B:  88% 12%



#### 4.2.7 Score per residue for model 7

- Molecule 1: DNA\_(5'-D(\*GP\*TP\*CP\*GP\*GP\*CP\*TP\*G)-3')

Chain A:  75% 25%




- Molecule 2: DNA\_(5'-D(\*CP\*AP\*GP\*CP\*CP\*GP\*AP\*C)-3')

Chain B:  100%

There are no outlier residues in this chain.

#### 4.2.8 Score per residue for model 8

- Molecule 1: DNA\_(5'-D(\*GP\*TP\*CP\*GP\*GP\*CP\*TP\*G)-3')

Chain A:  75% 25%




- Molecule 2: DNA\_(5'-D(\*CP\*AP\*GP\*CP\*CP\*GP\*AP\*C)-3')

Chain B:  100%

There are no outlier residues in this chain.

#### 4.2.9 Score per residue for model 9

- Molecule 1: DNA\_(5'-D(\*GP\*TP\*CP\*GP\*GP\*CP\*TP\*G)-3')

Chain A:  75% 25%




- Molecule 2: DNA\_(5'-D(\*CP\*AP\*GP\*CP\*CP\*GP\*AP\*C)-3')

Chain B:  100%

There are no outlier residues in this chain.


#### 4.2.10 Score per residue for model 10

- Molecule 1: DNA\_(5'-D(\*GP\*TP\*CP\*GP\*GP\*CP\*TP\*G)-3')

Chain A:  75% 25%



- Molecule 2: DNA\_(5'-D(\*CP\*AP\*GP\*CP\*CP\*GP\*AP\*C)-3')

Chain B:  88% 12%





## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	v2.22
X-PLOR NIH	refinement	v2.22

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	2
Total number of shifts	174
Number of shifts mapped to atoms	174
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	48%

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.03±0.01	2±0/182 ( 1.0± 0.2%)	1.37±0.01	4±0/280 ( 1.3± 0.2%)
2	B	0.93±0.01	0±0/178 ( 0.0± 0.0%)	1.21±0.01	0±0/272 ( 0.0± 0.0%)
All	All	0.98	19/3600 ( 0.5%)	1.30	37/5520 ( 0.7%)

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	7	DT	C5-C7	5.63	1.53	1.50	4	10
1	A	2	DT	C5-C7	5.36	1.53	1.50	2	9

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	2	DT	C6-C5-C7	-5.61	119.53	122.90	6	10
1	A	7	DT	C6-C5-C7	-5.60	119.54	122.90	4	10
1	A	2	DT	C4-C5-C6	5.23	121.14	118.00	2	8
1	A	7	DT	C4-C5-C6	5.12	121.07	118.00	3	9

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
2	B	159	90	90	1±0
All	All	3220	1820	1820	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:1:DC:HO5'	2:B:1:DC:H6	0.90	1.05	2	6
2:B:1:DC:O5'	2:B:1:DC:H6	0.46	1.93	6	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 48% for the well-defined parts and 48% for the entire structure.

### 7.1 Chemical shift list 1

File name: working\_cs.cif

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	150
Number of shifts mapped to atoms	150
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

#### 7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

#### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 42%, i.e. 136 atoms were assigned a chemical shift out of a possible 322. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Sugar	112/192 (58%)	112/112 (100%)	0/80 (0%)	0/0 (—%)
Base	24/130 (18%)	24/82 (29%)	0/26 (0%)	0/22 (0%)
Overall	136/322 (42%)	136/194 (70%)	0/106 (0%)	0/22 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 42%, i.e. 136 atoms were assigned a chemical shift out of a possible 322. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Sugar	112/192 (58%)	112/112 (100%)	0/80 (0%)	0/0 (—%)
Base	24/130 (18%)	24/82 (29%)	0/26 (0%)	0/22 (0%)
Overall	136/322 (42%)	136/194 (70%)	0/106 (0%)	0/22 (0%)

#### 7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

No *random coil index*(RCI) plot could be generated from the current chemical shift list. RCI is only applicable to proteins

## 7.2 Chemical shift list 2

File name: working\_cs.cif

Chemical shift list name: *assigned\_chem\_shift\_list\_2*

#### 7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	24
Number of shifts mapped to atoms	24
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

#### 7.2.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

#### 7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 6%, i.e. 18 atoms were assigned a chemical shift out of a possible 322. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Sugar	0/192 (0%)	0/112 (0%)	0/80 (0%)	0/0 (—%)
Base	18/130 (14%)	18/82 (22%)	0/26 (0%)	0/22 (0%)
Overall	18/322 (6%)	18/194 (9%)	0/106 (0%)	0/22 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 6%, i.e. 18 atoms were assigned a chemical shift out of a possible 322. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Sugar	0/192 (0%)	0/112 (0%)	0/80 (0%)	0/0 (—%)
Base	18/130 (14%)	18/82 (22%)	0/26 (0%)	0/22 (0%)
Overall	18/322 (6%)	18/194 (9%)	0/106 (0%)	0/22 (0%)

#### 7.2.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 7.2.5 Random Coil Index (RCI) plots [i](#)

No *random coil index*(RCI) plot could be generated from the current chemical shift list. RCI is only applicable to proteins