



Full wwPDB NMR Structure Validation Report ⓘ

Apr 21, 2024 – 06:54 AM EDT

PDB ID : 2MEK
BMRB ID : 19523
Title : N-terminal domain of Bilbo1 from Trypanosoma brucei
Authors : Vidilaseris, K.; Morriswood, B.; Kontaxis, G.; Dong, G.
Deposited on : 2013-09-24

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 78%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	110	

2 Ensemble composition and analysis i

This entry contains 20 models. Model 20 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:2-A:10, A:16-A:43, A:57-A:77, A:83-A:90 (66)	0.50	20

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 3, 6, 7, 11, 12, 13, 14, 15, 16, 17, 18, 20
2	10, 19
3	4, 8
Single-model clusters	2; 5; 9

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1789 atoms, of which 882 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called N-TERMINAL DOMAIN OF BILBO1.


Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	110	1789	580	882	153	170	4	0

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1


- Molecule 1: N-TERMINAL DOMAIN OF BILBO1



4.2.2 Score per residue for model 2



- Molecule 1: N-TERMINAL DOMAIN OF BILBO1







4.2.3 Score per residue for model 3

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1





4.2.4 Score per residue for model 4

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1




4.2.5 Score per residue for model 5

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1




4.2.6 Score per residue for model 6

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1




4.2.7 Score per residue for model 7

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1




4.2.8 Score per residue for model 8

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1




4.2.9 Score per residue for model 9

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1




4.2.10 Score per residue for model 10

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1




4.2.11 Score per residue for model 11

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1




4.2.12 Score per residue for model 12

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1




4.2.13 Score per residue for model 13

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1




4.2.14 Score per residue for model 14

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1




4.2.15 Score per residue for model 15

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1




4.2.16 Score per residue for model 16

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1




4.2.17 Score per residue for model 17

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1




4.2.18 Score per residue for model 18

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1




4.2.19 Score per residue for model 19

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1



4.2.20 Score per residue for model 20 (medoid)

- Molecule 1: N-TERMINAL DOMAIN OF BILBO1



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing, molecular dynamics*.

Of the 500 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	
X-PLOR NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1111
Number of shifts mapped to atoms	1110
Number of unparsed shifts	0
Number of shifts with mapping errors	1
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	78%

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	2.65±0.01	24±2/556 (4.4± 0.4%)	1.03±0.01	1±1/755 (0.1± 0.1%)
All	All	2.65	490/11120 (4.4%)	1.03	22/15100 (0.1%)

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	87	TYR	CB-CG	6.98	1.62	1.51	1	20
1	A	4	LEU	CA-CB	6.36	1.68	1.53	2	20
1	A	32	THR	CA-CB	6.21	1.69	1.53	4	20
1	A	16	VAL	CA-CB	6.18	1.67	1.54	17	20
1	A	72	VAL	CA-CB	6.12	1.67	1.54	5	20
1	A	58	SER	CA-CB	6.12	1.62	1.52	18	20
1	A	42	GLU	CB-CG	6.07	1.63	1.52	13	20
1	A	74	LEU	CB-CG	5.95	1.69	1.52	4	20
1	A	36	GLU	CB-CG	5.94	1.63	1.52	5	19
1	A	64	TYR	CB-CG	5.88	1.60	1.51	14	20
1	A	17	ASN	CA-CB	5.87	1.68	1.53	18	20
1	A	66	GLU	CB-CG	5.82	1.63	1.52	16	19
1	A	24	SER	CA-CB	5.72	1.61	1.52	10	9
1	A	77	GLU	CB-CG	5.71	1.62	1.52	20	20
1	A	19	GLU	CB-CG	5.69	1.62	1.52	20	20
1	A	88	VAL	CA-CB	5.67	1.66	1.54	2	2
1	A	83	TYR	CB-CG	5.65	1.60	1.51	18	18
1	A	20	LEU	CA-CB	5.61	1.66	1.53	7	20
1	A	88	VAL	CB-CG2	5.52	1.64	1.52	1	18
1	A	5	VAL	CA-CB	5.50	1.66	1.54	9	11
1	A	39	PHE	CB-CG	5.49	1.60	1.51	15	14
1	A	22	PHE	CB-CG	5.43	1.60	1.51	5	18
1	A	31	LEU	CA-CB	5.43	1.66	1.53	19	13
1	A	39	PHE	CA-CB	5.40	1.65	1.53	1	5
1	A	71	TRP	CA-CB	5.33	1.65	1.53	16	18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	3	PHE	CB-CG	5.26	1.60	1.51	7	16
1	A	86	LEU	CA-CB	5.21	1.65	1.53	8	2
1	A	84	CYS	CA-CB	5.20	1.65	1.53	19	1
1	A	77	GLU	CA-CB	5.19	1.65	1.53	5	2
1	A	20	LEU	CB-CG	5.18	1.67	1.52	1	14
1	A	26	PRO	N-CA	5.18	1.56	1.47	17	5
1	A	31	LEU	CB-CG	5.17	1.67	1.52	3	5
1	A	68	LEU	CA-CB	5.17	1.65	1.53	13	8
1	A	7	VAL	CB-CG2	5.13	1.63	1.52	9	3
1	A	67	GLU	CB-CG	5.09	1.61	1.52	1	1
1	A	27	SER	CA-CB	5.08	1.60	1.52	4	2
1	A	59	SER	CA-CB	5.05	1.60	1.52	3	4
1	A	26	PRO	N-CD	5.03	1.54	1.47	16	2
1	A	61	ILE	CA-CB	5.02	1.66	1.54	7	1

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	22	PHE	CB-CG-CD1	6.64	125.45	120.80	19	19
1	A	33	ARG	NE-CZ-NH1	5.58	123.09	120.30	19	2
1	A	22	PHE	CB-CG-CD2	-5.30	117.09	120.80	19	1

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	543	525	525	16±6
All	All	10860	10500	10500	311

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:61:ILE:CD1	1:A:88:VAL:HG13	1.30	1.55	16	8
1:A:61:ILE:HD11	1:A:88:VAL:CG1	1.26	1.60	16	6
1:A:28:ILE:HD12	1:A:28:ILE:O	1.25	1.26	4	18
1:A:61:ILE:HD13	1:A:61:ILE:N	1.24	1.36	9	4
1:A:39:PHE:O	1:A:43:ILE:HD13	1.20	1.36	12	3
1:A:60:LYS:C	1:A:61:ILE:HD12	1.12	1.65	1	2
1:A:61:ILE:HD11	1:A:88:VAL:HG13	1.05	1.18	19	7
1:A:75:ILE:HD13	1:A:75:ILE:H	1.04	0.98	9	1
1:A:28:ILE:O	1:A:28:ILE:CD1	1.03	2.05	5	18
1:A:61:ILE:N	1:A:61:ILE:CD1	1.03	2.17	9	2
1:A:75:ILE:HD12	1:A:75:ILE:H	0.99	1.13	7	6
1:A:75:ILE:HD12	1:A:75:ILE:N	0.98	1.73	11	6
1:A:61:ILE:CD1	1:A:61:ILE:N	0.95	2.29	1	1
1:A:75:ILE:HD13	1:A:75:ILE:N	0.95	1.70	9	1
1:A:63:MET:SD	1:A:75:ILE:HD13	0.95	2.01	5	3
1:A:61:ILE:HD13	1:A:61:ILE:H	0.95	1.17	9	1
1:A:43:ILE:HD12	1:A:43:ILE:N	0.94	1.77	19	2
1:A:39:PHE:O	1:A:43:ILE:CD1	0.93	2.17	19	4
1:A:61:ILE:HD12	1:A:88:VAL:HG13	0.91	1.42	8	3
1:A:28:ILE:O	1:A:28:ILE:HD12	0.91	1.65	5	2
1:A:28:ILE:HD12	1:A:28:ILE:C	0.90	1.82	4	20
1:A:60:LYS:C	1:A:61:ILE:HD13	0.89	1.88	8	5
1:A:75:ILE:N	1:A:75:ILE:CD1	0.88	2.35	9	6
1:A:61:ILE:HD12	1:A:61:ILE:N	0.87	1.80	1	4
1:A:75:ILE:H	1:A:75:ILE:CD1	0.85	1.84	9	5
1:A:60:LYS:C	1:A:61:ILE:CD1	0.83	2.45	1	2
1:A:43:ILE:N	1:A:43:ILE:CD1	0.83	2.42	19	2
1:A:61:ILE:CD1	1:A:88:VAL:CG1	0.81	2.38	16	6
1:A:28:ILE:O	1:A:28:ILE:HD13	0.81	1.72	5	6
1:A:63:MET:SD	1:A:75:ILE:CD1	0.80	2.69	10	2
1:A:28:ILE:HD11	1:A:31:LEU:HD23	0.79	1.53	12	10
1:A:61:ILE:HD12	1:A:88:VAL:HG22	0.77	1.56	15	4
1:A:43:ILE:CD1	1:A:43:ILE:H	0.74	1.94	19	2
1:A:63:MET:CE	1:A:75:ILE:CD1	0.73	2.67	16	3
1:A:43:ILE:CD1	1:A:43:ILE:N	0.72	2.53	12	1
1:A:43:ILE:N	1:A:43:ILE:HD12	0.71	2.00	12	2
1:A:61:ILE:CD1	1:A:88:VAL:HA	0.71	2.15	15	3
1:A:61:ILE:CD1	1:A:88:VAL:HG22	0.70	2.16	12	4
1:A:28:ILE:C	1:A:28:ILE:CD1	0.70	2.60	4	3
1:A:60:LYS:O	1:A:61:ILE:HD13	0.69	1.87	19	7
1:A:63:MET:HE2	1:A:75:ILE:HD11	0.68	1.64	17	3

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:28:ILE:CD1	1:A:28:ILE:C	0.67	2.57	5	14
1:A:63:MET:CE	1:A:75:ILE:HD11	0.67	2.20	16	4
1:A:61:ILE:CD1	1:A:88:VAL:CB	0.66	2.72	15	3
1:A:61:ILE:HD12	1:A:88:VAL:CG2	0.63	2.24	15	3
1:A:61:ILE:HD12	1:A:88:VAL:HA	0.63	1.70	13	6
1:A:60:LYS:O	1:A:61:ILE:HD12	0.61	1.94	1	1
1:A:63:MET:HE3	1:A:75:ILE:HD11	0.61	1.71	11	1
1:A:61:ILE:HD12	1:A:88:VAL:CB	0.59	2.28	16	3
1:A:75:ILE:HD12	1:A:75:ILE:O	0.58	1.96	3	1
1:A:10:ASP:O	1:A:90:GLN:O	0.57	2.21	5	2
1:A:61:ILE:N	1:A:61:ILE:HD13	0.57	2.14	1	1
1:A:63:MET:HE2	1:A:75:ILE:CD1	0.56	2.30	7	1
1:A:43:ILE:HD12	1:A:43:ILE:H	0.56	1.58	2	2
1:A:61:ILE:CD1	1:A:88:VAL:CG2	0.56	2.83	15	3
1:A:75:ILE:C	1:A:75:ILE:CD1	0.56	2.74	3	1
1:A:63:MET:CE	1:A:75:ILE:HD13	0.56	2.31	16	2
1:A:61:ILE:HD11	1:A:88:VAL:CB	0.56	2.30	15	1
1:A:61:ILE:HD12	1:A:88:VAL:CG1	0.56	2.29	4	2
1:A:61:ILE:CD1	1:A:88:VAL:CA	0.55	2.84	15	3
1:A:63:MET:HE1	1:A:75:ILE:HD13	0.55	1.79	7	3
1:A:39:PHE:O	1:A:43:ILE:HD12	0.54	2.02	2	1
1:A:63:MET:HE1	1:A:75:ILE:CD1	0.53	2.34	16	3
1:A:61:ILE:HD12	1:A:88:VAL:CA	0.52	2.33	13	3
1:A:59:SER:OG	1:A:89:PHE:O	0.51	2.28	5	4
1:A:31:LEU:HD11	1:A:86:LEU:HD22	0.50	1.82	8	1
1:A:75:ILE:O	1:A:75:ILE:CD1	0.49	2.60	3	1
1:A:9:ALA:HB2	1:A:39:PHE:CE1	0.49	2.42	1	4
1:A:39:PHE:N	1:A:39:PHE:CD1	0.47	2.82	16	10
1:A:39:PHE:CD1	1:A:39:PHE:N	0.47	2.83	20	10
1:A:60:LYS:CA	1:A:61:ILE:HD13	0.47	2.40	9	1
1:A:43:ILE:HD13	1:A:43:ILE:H	0.46	1.70	19	1
1:A:68:LEU:H	1:A:68:LEU:HD23	0.44	1.71	5	11
1:A:33:ARG:O	1:A:37:THR:OG1	0.43	2.27	3	2
1:A:63:MET:O	1:A:72:VAL:HG13	0.42	2.14	2	1
1:A:59:SER:O	1:A:61:ILE:CD1	0.42	2.68	3	1
1:A:61:ILE:HG12	1:A:88:VAL:HG13	0.41	1.91	18	1
1:A:74:LEU:C	1:A:75:ILE:HD12	0.40	2.35	2	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	66/110 (60%)	60±1 (92±1%)	4±1 (6±1%)	2±1 (2±1%)	9	46
All	All	1320/2200 (60%)	1208 (92%)	80 (6%)	32 (2%)	9	46

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	84	CYS	16
1	A	23	PRO	15
1	A	77	GLU	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	61/104 (59%)	50±1 (82±2%)	11±1 (18±2%)	4	38
All	All	1220/2080 (59%)	1002 (82%)	218 (18%)	4	38

All 28 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	4	LEU	20
1	A	28	ILE	20
1	A	33	ARG	20
1	A	39	PHE	20
1	A	68	LEU	20
1	A	72	VAL	19
1	A	31	LEU	17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	41	ASN	15
1	A	32	THR	13
1	A	75	ILE	11
1	A	7	VAL	10
1	A	84	CYS	6
1	A	61	ILE	3
1	A	65	ASP	3
1	A	17	ASN	3
1	A	5	VAL	3
1	A	57	HIS	3
1	A	43	ILE	2
1	A	66	GLU	1
1	A	76	ARG	1
1	A	88	VAL	1
1	A	74	LEU	1
1	A	10	ASP	1
1	A	90	GLN	1
1	A	86	LEU	1
1	A	42	GLU	1
1	A	67	GLU	1
1	A	62	LYS	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 78% for the well-defined parts and 72% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1111
Number of shifts mapped to atoms	1110
Number of unparsed shifts	0
Number of shifts with mapping errors	1
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- No matching atom found in the structure. All 1 occurrences are reported below.

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
1	A	1	MET	H	8.321	.	.

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	104	0.86 ± 0.12	Should be applied
$^{13}\text{C}_\beta$	99	1.00 ± 0.20	Should be applied
$^{13}\text{C}'$	89	0.23 ± 0.10	None needed (< 0.5 ppm)
^{15}N	100	-0.23 ± 0.34	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 78%, i.e. 720 atoms were assigned a chemical shift out of a possible 924. 0 out of 11 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	302/326 (93%)	124/130 (95%)	116/132 (88%)	62/64 (97%)
Sidechain	343/502 (68%)	229/326 (70%)	111/158 (70%)	3/18 (17%)
Aromatic	75/96 (78%)	39/47 (83%)	35/47 (74%)	1/2 (50%)
Overall	720/924 (78%)	392/503 (78%)	262/337 (78%)	66/84 (79%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 72%, i.e. 1107 atoms were assigned a chemical shift out of a possible 1544. 0 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	486/530 (92%)	196/210 (93%)	193/220 (88%)	97/100 (97%)
Sidechain	530/872 (61%)	351/562 (62%)	174/275 (63%)	5/35 (14%)
Aromatic	91/142 (64%)	47/71 (66%)	42/66 (64%)	2/5 (40%)
Overall	1107/1544 (72%)	594/843 (70%)	409/561 (73%)	104/140 (74%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	64	TYR	HB2	0.72	1.09 – 4.72	-6.0
1	A	64	TYR	HB3	0.72	0.93 – 4.76	-5.5

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

