

# Full wwPDB X-ray Structure Validation Report (i)

#### Oct 11, 2021 – 05:49 AM EDT

PDB ID : 2Q6B

Title: Design and synthesis of novel, conformationally restricted HMG-COA reduc-

tase inhibitors

Authors: Pavlovsky, A.; Pfefferkorn, J.A.; Harris, M.S.; Finzel, B.C.

Deposited on : 2007-06-04

Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

 $Xtriage\ (Phenix) \quad : \quad 1.13$ 

EDS: 2.23.2

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.23.2

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
1.136113	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	441	90%	6%	-
1	В	441	12%	7% •	-
1	С	441	90%	5% •	_
1	D	441	91%	• 5	5%



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 13820 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 3-hydroxy-3-methylglutaryl-coenzyme A reductase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	425	Total	С	N	О	S	0	0	0
1	A	420	3167	1974	558	605	30	0	U	
1	В	423	Total	С	N	О	S	0	0	0
1			3148	1962	553	603	30		U	
1	C	422	Total	С	N	О	S	0	0	0
1		422	3139	1957	552	600	30	0	U	
1	D	110	Total	С	N	О	S	0	0	0
1		418	3113	1940	549	594	30	0	0	0

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	435	HIS	-	expression tag	UNP P04035
A	436	HIS	- expression tag		UNP P04035
A	437	HIS	-	expression tag	UNP P04035
A	438	HIS	-	expression tag	UNP P04035
A	439	HIS	-	expression tag	UNP P04035
A	440	HIS	-	expression tag	UNP P04035
A	485	ILE	MET	engineered mutation	UNP P04035
В	435	HIS	-	expression tag	UNP P04035
В	436	HIS	-	expression tag	UNP P04035
В	437	HIS	-	expression tag	UNP P04035
В	438	HIS	-	expression tag	UNP P04035
В	439	HIS	-	expression tag	UNP P04035
В	440	HIS	-	expression tag	UNP P04035
В	485	ILE	MET	engineered mutation	UNP P04035
С	435	HIS	-	expression tag	UNP P04035
С	436	HIS	-	expression tag	UNP P04035
С	437	HIS	-	expression tag	UNP P04035
С	438	HIS	-	expression tag	UNP P04035
С	439	HIS	-	expression tag	UNP P04035
С	440	HIS	-	expression tag	UNP P04035
С	485	ILE	MET	engineered mutation	UNP P04035



Continued from previous page...

Chain	Residue	Modelled	Actual Comment		Reference
D	435	HIS	-	expression tag	UNP P04035
D	436	HIS	-	expression tag	UNP P04035
D	437	HIS	-	expression tag	UNP P04035
D	438	HIS	-	expression tag	UNP P04035
D	439	HIS	-	expression tag	UNP P04035
D	440	HIS	-	expression tag	UNP P04035
D	485	ILE	MET	engineered mutation	UNP P04035


 $\bullet$  Molecule 2 is SULFATE ION (three-letter code: SO4) (formula:  $\mathrm{O_4S}).$ 



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	В	1	Total O S 5 4 1	0	0
2	С	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0

• Molecule 3 is (3R,5R)-7-[3-(4-FLUOROPHENYL)-1-ISOPROPYL-8-OXO-7-PHENYL-1,4 ,5,6,7,8-HEXAHYDROPYRROLO[2,3-C]AZEPIN-2-YL]-3,5-DIHYDROXYHEPTANOIC ACID (three-letter code: HR2) (formula:  $C_{30}H_{35}FN_2O_5$ ).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	Λ	1	Total	С	F	N	О	0	0
3	Α	1	38	30	1	2	5	0	0
3	Λ	1	Total	С	F	N	О	0	0
3	Α	1	38	30	1	2	5	0	
3	C	1	Total	С	F	N	О	0	0
3	C	1	38	30	1	2	5	0	U
3	D	1	Total	С	F	N	О	0	0
3	ש	1	38	30	1	2	5	U	U

#### • Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	260	Total O 260 260	0	0
4	В	251	Total O 251 251	0	0
4	С	285	Total O 285 285	0	0
4	D	285	Total O 285 285	0	0



# 3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 3-hydroxy-3-methylglutaryl-coenzyme A reductase











# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	78.72Å 133.19Å 82.67Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $92.63^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	30.00 - 2.00	Depositor
rtesolution (A)	34.89 - 2.00	EDS
% Data completeness	(Not available) $(30.00-2.00)$	Depositor
(in resolution range)	91.4 (34.89-2.00)	EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	0.98  (at  2.00Å)	Xtriage
Refinement program	REFMAC	Depositor
$R, R_{free}$	0.224 , $0.256$	Depositor
	0.244 , (Not available)	DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.3	Xtriage
Anisotropy	0.278	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	$0.39 \;, 48.1$	EDS
L-test for twinning <sup>2</sup>	$< L > = 0.47, < L^2> = 0.30$	Xtriage
	0.023 for l,k,-h	
Estimated twinning fraction	0.037  for h,-k,-l	Xtriage
	0.033 for l,-k,h	
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	13820	wwPDB-VP
Average B, all atoms $(\mathring{A}^2)$	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HR2, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain	Bond	lengths	Bond angles		
MIOI		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.28	0/3214	0.58	5/4345~(0.1%)	
1	В	0.28	0/3194	0.57	6/4319 (0.1%)	
1	С	0.27	0/3185	0.59	6/4306 (0.1%)	
1	D	0.28	0/3159	0.58	5/4269 (0.1%)	
All	All	0.28	0/12752	0.58	$22/17239 \ (0.1\%)$	

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	${\rm Observed}(^o)$	$Ideal(^{o})$
1	A	653	ASP	CB-CG-OD2	5.75	123.48	118.30
1	D	547	ASP	CB-CG-OD2	5.67	123.40	118.30
1	С	790	ASP	CB-CG-OD2	5.56	123.30	118.30
1	A	790	ASP	CB-CG-OD2	5.52	123.27	118.30
1	С	690	ASP	CB-CG-OD2	5.51	123.26	118.30
1	A	547	ASP	CB-CG-OD2	5.49	123.24	118.30
1	С	547	ASP	CB-CG-OD2	5.46	123.21	118.30
1	A	516	ASP	CB-CG-OD2	5.42	123.18	118.30
1	В	653	ASP	CB-CG-OD2	5.40	123.16	118.30
1	С	653	ASP	CB-CG-OD2	5.39	123.15	118.30
1	В	623	ASP	CB-CG-OD2	5.29	123.06	118.30
1	D	516	ASP	CB-CG-OD2	5.26	123.04	118.30
1	D	653	ASP	CB-CG-OD2	5.26	123.03	118.30
1	D	601	ASP	CB-CG-OD2	5.26	123.03	118.30
1	С	623	ASP	CB-CG-OD2	5.21	122.99	118.30
1	В	790	ASP	CB-CG-OD2	5.18	122.97	118.30
1	В	547	ASP	CB-CG-OD2	5.17	122.95	118.30
1	В	829	ASP	CB-CG-OD2	5.11	122.90	118.30
1	С	516	ASP	CB-CG-OD2	5.06	122.85	118.30
1	В	690	ASP	CB-CG-OD2	5.04	122.83	118.30



Continued from previous page...

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	623	ASP	CB-CG-OD2	5.03	122.83	118.30
1	D	623	ASP	CB-CG-OD2	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3167	0	3207	8	0
1	В	3148	0	3187	10	0
1	С	3139	0	3182	9	0
1	D	3113	0	3151	3	0
2	A	5	0	0	0	0
2	В	5	0	0	0	0
2	С	5	0	0	0	0
2	D	5	0	0	0	0
3	A	76	0	68	0	0
3	С	38	0	34	1	0
3	D	38	0	34	1	0
4	A	260	0	0	0	0
4	В	251	0	0	0	0
4	С	285	0	0	0	0
4	D	285	0	0	0	0
All	All	13820	0	12863	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ (\rm \mathring{A}) \end{array}$	Clash overlap (Å)
1:C:471:VAL:HG21	1:C:481:LEU:HD11	1.73	0.70
1:C:771:ASN:OD1	1:C:775:SER:OG	2.12	0.67
1:A:771:ASN:OD1	1:A:775:SER:OG	2.22	0.57



Continued from previous page...

A Land		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ ({\rm \AA})$	overlap (Å)
1:C:470:LEU:HB3	1:C:476:ILE:HG21	1.91	0.52
1:A:477:PRO:O	1:A:478:ALA:HB2	2.10	0.51
1:B:771:ASN:OD1	1:B:775:SER:OG	2.29	0.51
1:D:771:ASN:OD1	1:D:775:SER:OG	2.28	0.51
1:A:454:ASN:O	1:A:455:ALA:HB3	2.11	0.50
1:C:471:VAL:CG2	1:C:481:LEU:HD11	2.42	0.50
1:C:454:ASN:O	1:C:455:ALA:HB3	2.11	0.50
1:C:476:ILE:HD11	1:C:484:LEU:HD22	1.94	0.48
1:C:496:ARG:NH1	1:C:512:LEU:O	2.47	0.48
1:B:635:HIS:CD2	1:B:635:HIS:C	2.89	0.46
1:B:441:GLU:N	1:B:442:PRO:CD	2.79	0.45
3:C:3003:HR2:O14	3:C:3003:HR2:H113	2.16	0.45
1:B:555:MET:HE3	1:B:563:VAL:HG22	2.00	0.44
3:D:3004:HR2:O14	3:D:3004:HR2:H113	2.16	0.44
1:C:454:ASN:O	1:C:455:ALA:CB	2.64	0.44
1:B:782:GLU:OE2	1:C:641:ARG:HD3	2.18	0.44
1:D:560:GLY:O	1:D:561:CYS:HB2	2.18	0.44
1:A:477:PRO:O	1:A:478:ALA:CB	2.65	0.43
1:B:793:ILE:HD13	1:B:851:LEU:HG	2.00	0.43
1:D:774:SER:HA	1:D:799:SER:O	2.19	0.43
1:A:454:ASN:O	1:A:455:ALA:CB	2.67	0.43
1:A:702:ARG:O	1:A:799:SER:HA	2.19	0.43
1:A:793:ILE:HD13	1:A:851:LEU:HG	2.00	0.43
1:B:472:ASN:HD21	1:B:501:LYS:HZ3	1.67	0.42
1:B:702:ARG:O	1:B:799:SER:HA	2.19	0.42
1:B:635:HIS:CE1	1:B:696:ILE:HD11	2.55	0.41
1:B:541:ALA:HB2	1:B:555:MET:HE1	2.02	0.41
1:A:471:VAL:HG21	1:A:481:LEU:HD11	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Analysed Favoured Allowed		Outliers	Perce	ntiles
1	A	423/441 (96%)	402 (95%)	19 (4%)	2 (0%)	29	23
1	В	421/441 (96%)	404 (96%)	17 (4%)	0	100	100
1	C	420/441 (95%)	403 (96%)	15 (4%)	2 (0%)	29	23
1	D	414/441 (94%)	401 (97%)	13 (3%)	0	100	100
All	All	1678/1764 (95%)	1610 (96%)	64 (4%)	4 (0%)	47	44

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	478	ALA
1	A	455	ALA
1	С	455	ALA
1	С	479	TYR

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	339/355~(96%)	326 (96%)	13 (4%)	33 31		
1	В	337/355 (95%)	324 (96%)	13 (4%)	32 30		
1	C	336/355~(95%)	330 (98%)	6 (2%)	59 63		
1	D	334/355 (94%)	327 (98%)	7 (2%)	53 57		
All	All	1346/1420 (95%)	1307 (97%)	39 (3%)	42 43		

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	447	GLU
1	A	468	ILE
1	A	474	LYS
1	A	480	LYS
1	A	486	GLU
1	A	490	ARG
1	A	512	LEU



Continued from previous page...

Mol	Chain	Res	Type
1	A	568	ARG
1	A	571	ARG
1	A	630	ARG
1	A	787	THR
1	A	814	GLN
1	A	864	LYS
1	В	446	GLU
1	В	449	LEU
1	В	464	ASP
1	В	501	LYS
1	В	505	GLU
1	В	523	MET
1	В	568	ARG
1	В	571	ARG
1	В	630	ARG
1	В	635	HIS
1	В	752	HIS
1	В	784	SER
1	В	862	LEU
1	С	470	LEU
1	С	490	ARG
1	С	568	ARG
1	С	627	ARG
1	C C	752	HIS
1		862	LEU
1	D	466	GLU
1	D	475	HIS
1	D	482	GLU
1	D	487	THR
1	D	627	ARG
1	D	752	HIS
1	D	828	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	445	ASN
1	A	472	ASN
1	A	475	HIS
1	A	510	GLN
1	A	518	ASN
1	A	567	ASN



Continued from previous page...

Mol	Chain	Res	Type
1	A	632	GLN
1	A	861	HIS
1	В	445	ASN
1	В	472	ASN
1	В	632	GLN
1	С	472	ASN
1	D	518	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal True Chain		Dag	s Link	Bond lengths			Bond angles			
Mol	Type	Chain	Res	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	HR2	A	3001	-	35,41,41	0.88	2 (5%)	42,58,58	1.57	3 (7%)
2	SO4	A	2001	-	4,4,4	0.13	0	6,6,6	0.12	0
3	HR2	D	3004	-	35,41,41	0.89	2 (5%)	42,58,58	1.57	3 (7%)
2	SO4	В	2002	-	4,4,4	0.15	0	6,6,6	0.07	0
2	SO4	С	2003	-	4,4,4	0.14	0	6,6,6	0.16	0
2	SO4	D	2004	-	4,4,4	0.15	0	6,6,6	0.16	0



Mol	Type Chain		Chain Dea	Link	Bond lengths			Bond angles		
IVIOI	туре	Chain	Res	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	HR2	С	3003	-	35,41,41	0.90	2 (5%)	42,58,58	1.60	6 (14%)
3	HR2	A	3002	-	35,41,41	0.89	1 (2%)	42,58,58	1.55	4 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HR2	A	3002	-	-	4/23/39/39	0/4/4/4
3	HR2	С	3003	-	-	5/23/39/39	0/4/4/4
3	HR2	D	3004	-	-	5/23/39/39	0/4/4/4
3	HR2	A	3001	-	-	5/23/39/39	0/4/4/4

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\mathring{\mathrm{A}})$	Ideal(Å)
3	A	3002	HR2	C17-N15	-3.30	1.36	1.43
3	D	3004	HR2	C17-N15	-3.24	1.36	1.43
3	С	3003	HR2	C17-N15	-3.21	1.36	1.43
3	A	3001	HR2	C17-N15	-3.06	1.37	1.43
3	С	3003	HR2	C13-N15	-2.02	1.34	1.37
3	A	3001	HR2	C13-N15	-2.00	1.34	1.37
3	D	3004	HR2	C13-N15	-2.00	1.34	1.37

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	3001	HR2	C3-C5-C4	6.82	112.20	103.88
3	D	3004	HR2	C3-C5-C4	6.79	112.16	103.88
3	С	3003	HR2	C3-C5-C4	6.75	112.11	103.88
3	A	3002	HR2	C3-C5-C4	6.72	112.07	103.88
3	D	3004	HR2	C20-C19-C3	-2.82	106.28	112.66
3	С	3003	HR2	C19-C20-C21	-2.80	109.66	115.05
3	A	3002	HR2	C2-C6-C4	-2.80	109.54	113.83
3	A	3001	HR2	C2-C6-C4	-2.61	109.83	113.83
3	D	3004	HR2	C2-C6-C4	-2.56	109.91	113.83
3	С	3003	HR2	C2-C6-C4	-2.43	110.10	113.83
3	С	3003	HR2	C22-C23-C24	-2.25	108.70	113.19
3	A	3001	HR2	C18-C7-C15	-2.21	119.88	122.83
3	A	3002	HR2	C18-C7-C15	-2.19	119.92	122.83



Continued from previous page...

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	3002	HR2	C20-C19-C3	-2.07	107.98	112.66
3	С	3003	HR2	C18-C7-C15	-2.04	120.11	122.83
3	С	3003	HR2	C9-C5-C3	-2.04	122.49	127.85

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	С	3003	HR2	C19-C20-C21-O33
3	A	3001	HR2	C32-C17-N15-C16
3	A	3002	HR2	C28-C17-N15-C16
3	A	3002	HR2	C32-C17-N15-C16
3	С	3003	HR2	C32-C17-N15-C16
3	С	3003	HR2	C19-C20-C21-C22
3	A	3002	HR2	C11-C10-N1-C3
3	D	3004	HR2	C12-C10-N1-C3
3	A	3001	HR2	C28-C17-N15-C16
3	D	3004	HR2	C28-C17-N15-C16
3	D	3004	HR2	C32-C17-N15-C16
3	A	3001	HR2	C19-C20-C21-O33
3	D	3004	HR2	C19-C20-C21-O33
3	A	3001	HR2	C12-C10-N1-C3
3	A	3001	HR2	C11-C10-N1-C3
3	A	3002	HR2	C12-C10-N1-C3
3	С	3003	HR2	C12-C10-N1-C3
3	С	3003	HR2	C11-C10-N1-C3
3	D	3004	HR2	C11-C10-N1-C3

There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	3004	HR2	1	0
3	С	3003	HR2	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring




in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














# 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	425/441 (96%)	0.41	26 (6%) 21 20	20, 29, 52, 63	0
1	В	423/441 (95%)	0.68	53 (12%) 3 3	18, 28, 58, 60	0
1	С	422/441 (95%)	0.29	14 (3%) 46 45	19, 28, 40, 57	0
1	D	418/441 (94%)	0.56	40 (9%) 8 7	17, 28, 56, 61	0
All	All	1688/1764 (95%)	0.48	133 (7%) 12 11	17, 28, 55, 63	0

All (133) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	481	LEU	9.1
1	В	449	LEU	8.8
1	В	453	GLY	8.7
1	D	484	LEU	8.7
1	В	461	PHE	8.4
1	В	452	LEU	8.2
1	В	479	TYR	8.0
1	D	479	TYR	7.9
1	В	476	ILE	7.9
1	D	483	THR	7.5
1	D	475	HIS	7.2
1	В	451	ILE	6.7
1	D	476	ILE	6.6
1	В	483	THR	6.6
1	В	475	HIS	6.3
1	В	455	ALA	6.2
1	A	479	TYR	6.1
1	С	477	PRO	6.0
1	D	480	LYS	5.9
1	В	477	PRO	5.8
1	С	479	TYR	5.8



 $Continued\ from\ previous\ page...$ 

Mol	Chain	m Res	Type	RSRZ
1	D	473	ALA	5.8
1	D	461	PHE	5.7
1	D	470	LEU	5.6
1	D	449	LEU	5.6
1	D	458	GLY	5.5
1	С	863	VAL	5.5
1	D	442	PRO	5.4
1	В	463	SER	5.4
1	В	458	GLY	5.3
1	В	445	ASN	5.3
1	A	477	PRO	5.3
1	В	450	GLN	5.2
1	В	441	GLU	5.2
1	D	462	LEU	5.2
1	В	486	GLU	5.1
1	В	457	LYS	5.1
1	В	473	ALA	4.9
1	D	443	ARG	4.9
1	С	861	HIS	4.8
1	D	447	GLU	4.5
1	В	480	LYS	4.5
1	В	481	LEU	4.5
1	A	484	LEU	4.5
1	D	445	ASN	4.5
1	A	481	LEU	4.4
1	В	462	LEU	4.4
1	С	478	ALA	4.4
1	A	483	THR	4.3
1	A	480	LYS	4.2
1	D	448	CYS	4.2
1	В	447	GLU	4.2
1	В	443	ARG	4.1
1	В	484	LEU	4.1
1	D	446	GLU	4.1
1	В	448	CYS	4.0
1	В	446	GLU	4.0
1	D	477	PRO	3.9
1	В	472	ASN	3.9
1	В	470	LEU	3.8
1	В	454	ASN	3.8
1	D	469	GLN	3.8
1	В	485	ILE	3.7



Continued from previous page...

Mol	nued fron Chain	Res	Type	RSRZ
1	A	451	ILE	3.7
1	D	474	LYS	3.7
1	В	456	GLU	3.7
1	С	474	LYS	3.7
1	В	487	THR	3.7
1	D	486	GLU	3.6
1	В	482	GLU	3.6
1	D	467	ILE	3.6
1	В	444	PRO	3.5
1	С	476	ILE	3.5
1	В	787	THR	3.5
1	A	446	GLU	3.5
1	A	474	LYS	3.5
1	В	786	PRO	3.4
1	В	460	LYS	3.4
1	D	478	ALA	3.4
1	D	482	GLU	3.3
1	С	481	LEU	3.3
1	D	487	THR	3.2
1	A	459	ALA	3.2
1	A	786	PRO	3.2
1	D	450	GLN	3.2
1	A	476	ILE	3.2
1	A	472	ASN	3.2
1	D	463	SER	3.0
1	A	473	ALA	2.9
1	D	465	ALA	2.9
1	В	469	GLN	2.9
1	В	442	PRO	2.9
1	В	524	GLY	2.9
1	A	482	GLU	2.9
1	D	444	PRO	2.9
1	В	506	PRO	2.8
1	A	478	ALA	2.8
1	D	468	ILE	2.8
1	D	787	THR	2.8
1	D	471	VAL	2.7
1	В	511	TYR	2.7
1	В	465	ALA	2.7
1	D	524	GLY	2.7
1	В	863	VAL	2.6
1	A	457	LYS	2.6



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	С	475	HIS	2.6
1	В	861	HIS	2.5
1	D	866	HIS	2.5
1	A	444	PRO	2.5
1	В	474	LYS	2.5
1	A	787	THR	2.4
1	В	829	ASP	2.4
1	A	485	ILE	2.4
1	D	472	ASN	2.4
1	A	475	HIS	2.4
1	В	828	LYS	2.4
1	A	458	GLY	2.3
1	D	630	ARG	2.3
1	D	466	GLU	2.3
1	A	460	LYS	2.3
1	A	445	ASN	2.3
1	С	473	ALA	2.2
1	С	862	LEU	2.2
1	С	442	PRO	2.2
1	A	442	PRO	2.2
1	D	460	LYS	2.1
1	В	459	ALA	2.1
1	С	627	ARG	2.1
1	С	828	LYS	2.1
1	В	507	SER	2.1
1	В	468	ILE	2.1
1	A	452	LEU	2.1
1	В	464	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, $95^{th}$ percentile and maximum values of B factors of atoms in the group.	The column
labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.	

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B-factors}({f \AA}^2)$	Q < 0.9
3	HR2	A	3001	38/38	0.86	0.15	27,33,37,38	0
3	HR2	С	3003	38/38	0.86	0.15	28,37,40,42	0
2	SO4	В	2002	5/5	0.87	0.17	63,63,63,63	0
3	HR2	A	3002	38/38	0.90	0.16	28,32,35,37	0
2	SO4	С	2003	5/5	0.91	0.15	46,47,47,47	0
3	HR2	D	3004	38/38	0.91	0.15	24,29,34,36	0
2	SO4	A	2001	5/5	0.93	0.14	53,53,54,54	0
2	SO4	D	2004	5/5	0.96	0.13	44,45,45,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








# 





# 6.5 Other polymers (i)

There are no such residues in this entry.

