



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 3, 2023 – 09:39 pm GMT

PDB ID : 2VCY
Title : Crystal Structure of 2-Enoyl Thioester Reductase of Human FAS II
Authors : Haapalainen, A.M.; Pudas, R.; Smart, O.S.; Wierenga, R.K.
Deposited on : 2007-09-28
Resolution : 2.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4647 (2.44-2.40)
RSRZ outliers	127900	4543 (2.44-2.40)

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5401 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANS-2-ENOYL-COA REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	332	2528	1596	448	470	14	0	0	0
1	B	332	2528	1596	448	470	14	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	153	Total O 153 153	0	0
3	B	137	Total O 137 137	0	0

MolProbity failed to run properly - this section is therefore empty.

3 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	104.62Å 104.62Å 146.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.79 – 2.41 31.49 – 2.41	Depositor EDS
% Data completeness (in resolution range)	84.4 (33.79-2.41) 84.6 (31.49-2.41)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 2.42Å)	Xtrriage
Refinement program	BUSTER-TNT 2.1.1	Depositor
R, R_{free}	0.207 , 0.267 0.212 , 0.274	Depositor DCC
R_{free} test set	1391 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	49.9	Xtrriage
Anisotropy	0.208	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 60.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5401	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	1378	-	4,4,4	0.40	0	6,6,6	0.30	0
2	SO4	A	1374	-	4,4,4	0.30	0	6,6,6	0.21	0
2	SO4	A	1377	-	4,4,4	0.25	0	6,6,6	0.30	0
2	SO4	B	1376	-	4,4,4	0.18	0	6,6,6	0.08	0
2	SO4	B	1377	-	4,4,4	0.28	0	6,6,6	0.10	0
2	SO4	B	1379	-	4,4,4	0.27	0	6,6,6	0.20	0
2	SO4	B	1378	-	4,4,4	0.28	0	6,6,6	0.14	0
2	SO4	A	1375	-	4,4,4	0.22	0	6,6,6	0.21	0
2	SO4	B	1374	-	4,4,4	0.37	0	6,6,6	0.22	0
2	SO4	A	1376	-	4,4,4	0.38	0	6,6,6	0.15	0
2	SO4	B	1375	-	4,4,4	0.37	0	6,6,6	0.11	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	332/344 (96%)	0.02	12 (3%) 42 41	31, 51, 92, 129	0
1	B	332/344 (96%)	0.04	18 (5%) 25 23	30, 50, 91, 128	0
All	All	664/688 (96%)	0.03	30 (4%) 33 31	30, 51, 92, 129	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	251	PHE	8.2
1	A	247	MET	6.5
1	A	248	LYS	4.9
1	A	221	ILE	4.9
1	A	245	PRO	4.5
1	A	244	ARG	4.3
1	A	243	ARG	4.3
1	A	249	ASN	3.8
1	B	222	GLN	3.5
1	B	251	PHE	3.4
1	B	196	VAL	3.3
1	B	252	LYS	3.2
1	B	63	LEU	3.1
1	B	226	ASP	2.8
1	A	250	PHE	2.7
1	B	364	PHE	2.6
1	B	221	ILE	2.5
1	B	114	ALA	2.5
1	B	56	LYS	2.5
1	B	116	GLY	2.5
1	A	246	GLU	2.4
1	B	225	SER	2.4
1	B	42	ARG	2.4
1	B	57	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	290	LYS	2.1
1	B	64	GLU	2.1
1	A	289	ALA	2.1
1	B	289	ALA	2.0
1	A	49	GLY	2.0
1	B	250	PHE	2.0

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	1377	5/5	0.78	0.25	71,75,76,77	5
2	SO4	B	1376	5/5	0.88	0.20	93,98,99,99	5
2	SO4	B	1375	5/5	0.89	0.21	75,79,80,82	5
2	SO4	B	1379	5/5	0.94	0.13	57,61,63,64	0
2	SO4	A	1376	5/5	0.95	0.14	67,72,73,73	5
2	SO4	B	1374	5/5	0.95	0.11	70,74,75,76	5
2	SO4	A	1375	5/5	0.95	0.16	72,76,77,78	5
2	SO4	A	1374	5/5	0.96	0.13	64,67,68,69	5
2	SO4	B	1378	5/5	0.97	0.23	81,85,86,87	0
2	SO4	A	1377	5/5	0.97	0.14	65,68,70,70	0
2	SO4	A	1378	5/5	0.98	0.10	57,61,62,63	0

5.5 Other polymers [i](#)

There are no such residues in this entry.