



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 6, 2023 – 07:29 am GMT

PDB ID : 2WAP  
Title : 3D-crystal structure of humanized-rat fatty acid amide hydrolase (FAAH) conjugated with the drug-like urea inhibitor PF-3845  
Authors : Mileni, M.; Kamtekar, S.; Stevens, R.C.  
Deposited on : 2009-02-11  
Resolution : 2.80 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : **NOT EXECUTED**  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36

# 1 Overall quality at a glance



The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

There are no overall percentile quality scores available for this entry.

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	543	 % 79% 20% •
1	B	543	 % 82% 17% •

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8468 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FATTY-ACID AMIDE HYDROLASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	541	4174	2663	712	769	30	0	0	0
1	B	543	4195	2678	715	772	30	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	192	PHE	LEU	engineered mutation	UNP P97612
A	194	TYR	PHE	engineered mutation	UNP P97612
A	377	THR	ALA	engineered mutation	UNP P97612
A	435	ASN	SER	engineered mutation	UNP P97612
A	491	VAL	ILE	engineered mutation	UNP P97612
A	495	MET	VAL	engineered mutation	UNP P97612
B	192	PHE	LEU	engineered mutation	UNP P97612
B	194	TYR	PHE	engineered mutation	UNP P97612
B	377	THR	ALA	engineered mutation	UNP P97612
B	435	ASN	SER	engineered mutation	UNP P97612
B	491	VAL	ILE	engineered mutation	UNP P97612
B	495	MET	VAL	engineered mutation	UNP P97612

- Molecule 2 is 4-(3-{[5-(trifluoromethyl)pyridin-2-yl]oxy}benzyl)piperidine-1-carboxylic acid (three-letter code: PIX) (formula: C<sub>19</sub>H<sub>19</sub>F<sub>3</sub>N<sub>2</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total	C	F	N	O	0	0
			26	19	3	2	2		
2	B	1	Total	C	F	N	O	0	0
			26	19	3	2	2		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		
4	B	2	Total	Na	0	0
			2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	18	Total	O	0	0
			18	18		
5	B	25	Total	O	0	0
			25	25		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: FATTY-ACID AMIDE HYDROLASE 1



- Molecule 1: FATTY-ACID AMIDE HYDROLASE 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.77Å 105.20Å 148.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.01 – 2.80 37.01 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.9 (37.01-2.80) 98.9 (37.01-2.80)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.22 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.192 , 0.232 0.218 , 0.258	Depositor DCC
$R_{free}$ test set	1963 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.0	Xtrriage
Anisotropy	0.089	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 24.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.029 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	8468	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PIX, NA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	1/4267 (0.0%)	0.71	1/5787 (0.0%)
1	B	0.57	2/4290 (0.0%)	0.70	1/5820 (0.0%)
All	All	0.56	3/8557 (0.0%)	0.70	2/11607 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	252	CYS	CB-SG	-5.58	1.72	1.81
1	B	252	CYS	CB-SG	-5.58	1.72	1.81
1	B	499	CYS	CB-SG	-5.09	1.73	1.81

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	420	LEU	CA-CB-CG	6.92	131.22	115.30
1	B	74	LEU	CA-CB-CG	5.08	126.99	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4174	0	4225	40	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	4195	0	4242	34	0
2	A	26	0	18	0	0
2	B	26	0	18	1	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
4	B	2	0	0	0	0
5	A	18	0	0	2	0
5	B	25	0	0	1	0
All	All	8468	0	8503	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:486:ARG:HB3	1:A:534:ILE:HD13	1.75	0.69
1:B:34:ARG:HG2	1:B:396:PHE:CE2	2.33	0.64
1:A:472:THR:HG21	1:A:550:GLN:HE21	1.64	0.62
1:A:287:VAL:HG12	1:A:569:GLU:OE2	2.00	0.62
1:A:287:VAL:HG11	1:A:566:ARG:HA	1.82	0.61
1:A:559:GLU:HG2	5:A:2017:HOH:O	2.00	0.61
1:A:79:LEU:HD23	1:A:231:PRO:HB2	1.83	0.61
1:B:119:THR:H	1:B:150:HIS:CE1	2.21	0.59
1:A:463:LYS:HE2	1:B:304:THR:O	2.02	0.59
1:A:426:PHE:HB3	1:A:429:LEU:HD22	1.83	0.59
1:A:345:ALA:O	1:A:349:THR:HG23	2.04	0.57
1:A:472:THR:HG22	1:A:473:PRO:O	2.04	0.57
1:A:472:THR:CG2	1:A:550:GLN:HE21	2.17	0.57
1:B:470:LEU:HB3	1:B:552:VAL:HB	1.87	0.57
1:B:345:ALA:O	1:B:349:THR:HG23	2.04	0.56
1:A:52:LEU:HD21	1:A:108:ASN:ND2	2.21	0.55
1:B:341:ALA:HB1	1:B:547:VAL:HG21	1.89	0.55
1:B:51:SER:O	1:B:55:MET:HG2	2.07	0.54
1:A:269:CYS:HB2	1:A:391:ASN:ND2	2.23	0.54
1:A:173:VAL:O	1:A:177:GLN:HB2	2.10	0.51
1:B:119:THR:H	1:B:150:HIS:HE1	1.56	0.51
1:A:341:ALA:HB1	1:A:547:VAL:HG21	1.92	0.50
1:A:474:MET:HG2	1:A:506:VAL:HG21	1.93	0.50
1:A:74:LEU:HD21	1:A:96:VAL:HA	1.94	0.49
1:A:119:THR:H	1:A:150:HIS:CE1	2.30	0.49

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:418:LEU:HB3	1:B:433:LEU:HD13	1.95	0.49
1:A:118:LEU:HD13	1:A:147:TYR:CD1	2.48	0.49
1:B:237:ASP:HA	1:B:241:SER:HB2	1.95	0.48
1:B:269:CYS:HB2	1:B:391:ASN:ND2	2.27	0.48
1:A:41:ALA:O	1:A:45:ARG:HG3	2.13	0.48
1:B:239:GLY:HA2	1:B:498:ASN:OD1	2.14	0.48
1:A:304:THR:HG23	1:B:463:LYS:HE2	1.96	0.47
1:A:169:VAL:CG2	1:A:264:SER:HB3	2.44	0.47
1:A:193:SER:HB2	1:A:401:LEU:HD12	1.96	0.47
1:B:69:LEU:HD12	1:B:70:ASP:N	2.31	0.46
1:A:271:TYR:HB3	1:B:449:HIS:CG	2.50	0.46
1:B:119:THR:OG1	1:B:150:HIS:HE1	1.99	0.46
1:A:43:ARG:NH2	1:A:160:GLU:OE1	2.48	0.46
1:B:456:GLN:HG3	5:B:2018:HOH:O	2.15	0.46
1:A:212:LYS:NZ	1:A:538:ALA:O	2.49	0.45
1:A:55:MET:O	1:A:59:VAL:HG23	2.17	0.44
1:B:483:THR:N	1:B:484:PRO:CD	2.81	0.44
1:A:285:ARG:HD3	5:A:2004:HOH:O	2.17	0.43
1:B:41:ALA:O	1:B:45:ARG:CG	2.66	0.43
1:B:193:SER:C	2:B:1574:PIX:H12C	2.39	0.43
1:A:262:SER:HB2	1:A:302:LEU:HD11	2.01	0.43
1:A:437:ARG:O	1:A:439:ARG:NH1	2.51	0.43
1:A:472:THR:HG21	1:A:550:GLN:NE2	2.32	0.43
1:A:483:THR:N	1:A:484:PRO:CD	2.81	0.43
1:B:169:VAL:CG2	1:B:264:SER:HB3	2.49	0.43
1:B:197:SER:HB3	1:B:203:GLN:HB2	2.01	0.43
1:B:86:LEU:HD11	1:B:96:VAL:HG11	2.01	0.43
1:A:177:GLN:HG2	1:A:296:ALA:HB1	2.01	0.43
1:B:416:ARG:NH1	1:B:420:LEU:HD21	2.34	0.43
1:A:426:PHE:CB	1:A:429:LEU:HD22	2.48	0.42
1:B:74:LEU:C	1:B:74:LEU:HD12	2.40	0.42
1:A:237:ASP:HA	1:A:241:SER:HB2	2.01	0.41
1:B:177:GLN:HG2	1:B:296:ALA:HB1	2.01	0.41
1:A:428:ARG:HG3	1:A:527:PHE:CE2	2.55	0.41
1:A:51:SER:O	1:A:55:MET:HG3	2.19	0.41
1:B:60:GLN:O	1:B:64:LEU:HD13	2.21	0.41
1:B:269:CYS:HB2	1:B:391:ASN:CG	2.41	0.41
1:B:85:LYS:HD3	1:B:90:GLU:OE1	2.21	0.41
1:A:244:PHE:HZ	1:A:491:VAL:HA	1.86	0.41
1:B:211:SER:O	1:B:481:LEU:HB2	2.21	0.40
1:B:236:THR:HB	1:B:278:LEU:HD11	2.02	0.40

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:74:LEU:HD21	1:B:96:VAL:HA	2.02	0.40
1:B:131:GLN:NE2	1:B:131:GLN:HA	2.36	0.40
1:B:488:THR:HB	1:B:531:TRP:CE2	2.56	0.40
1:A:194:TYR:CD1	1:A:484:PRO:HB3	2.57	0.40
1:A:262:SER:CB	1:A:302:LEU:HD11	2.52	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	539/543 (99%)	518 (96%)	21 (4%)	0	100	100
1	B	541/543 (100%)	519 (96%)	22 (4%)	0	100	100
All	All	1080/1086 (99%)	1037 (96%)	43 (4%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	459/461 (100%)	401 (87%)	58 (13%)	4	14
1	B	461/461 (100%)	409 (89%)	52 (11%)	6	18
All	All	920/922 (100%)	810 (88%)	110 (12%)	5	15

All (110) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	ARG
1	A	36	LYS
1	A	46	GLN
1	A	53	GLU
1	A	60	GLN
1	A	61	ARG
1	A	65	GLN
1	A	69	LEU
1	A	74	LEU
1	A	77	LEU
1	A	80	LEU
1	A	82	LEU
1	A	84	GLN
1	A	86	LEU
1	A	96	VAL
1	A	101	LEU
1	A	115	THR
1	A	120	ASP
1	A	125	LEU
1	A	127	GLN
1	A	130	ARG
1	A	131	GLN
1	A	133	LEU
1	A	156	LEU
1	A	158	LEU
1	A	192	PHE
1	A	200	LEU
1	A	234	LEU
1	A	252	CYS
1	A	269	CYS
1	A	270	VAL
1	A	273	GLN
1	A	280	LEU
1	A	297	LEU
1	A	320	ARG
1	A	323	ARG
1	A	359	THR
1	A	374	VAL
1	A	397	VAL
1	A	401	LEU
1	A	416	ARG
1	A	418	LEU

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	420	LEU
1	A	422	LEU
1	A	423	LYS
1	A	429	LEU
1	A	439	ARG
1	A	444	LEU
1	A	466	ASN
1	A	469	VAL
1	A	507	VAL
1	A	521	GLU
1	A	524	LYS
1	A	530	ILE
1	A	537	LYS
1	A	543	VAL
1	A	549	VAL
1	A	563	ARG
1	B	32	THR
1	B	36	LYS
1	B	48	GLN
1	B	52	LEU
1	B	55	MET
1	B	65	GLN
1	B	72	GLU
1	B	74	LEU
1	B	84	GLN
1	B	86	LEU
1	B	96	VAL
1	B	101	LEU
1	B	120	ASP
1	B	125	LEU
1	B	127	GLN
1	B	130	ARG
1	B	154	LEU
1	B	156	LEU
1	B	158	LEU
1	B	162	MET
1	B	184	HIS
1	B	192	PHE
1	B	203	GLN
1	B	273	GLN
1	B	286	ASP
1	B	287	VAL

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	B	292	LEU
1	B	297	LEU
1	B	316	GLU
1	B	320	ARG
1	B	332	THR
1	B	360	LEU
1	B	374	VAL
1	B	401	LEU
1	B	416	ARG
1	B	418	LEU
1	B	422	LEU
1	B	425	LEU
1	B	439	ARG
1	B	444	LEU
1	B	469	VAL
1	B	472	THR
1	B	491	VAL
1	B	506	VAL
1	B	507	VAL
1	B	510	THR
1	B	521	GLU
1	B	530	ILE
1	B	549	VAL
1	B	559	GLU
1	B	560	LEU
1	B	563	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	150	HIS
1	A	203	GLN
1	A	351	GLN
1	B	46	GLN
1	B	48	GLN
1	B	66	ASN
1	B	131	GLN
1	B	150	HIS
1	B	203	GLN
1	B	351	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PIX	B	1574	1	28,28,29	0.66	1 (3%)	39,39,41	1.74	7 (17%)
2	PIX	A	1574	1	28,28,29	0.72	1 (3%)	39,39,41	1.45	6 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PIX	B	1574	1	-	1/16/26/28	0/3/3/3
2	PIX	A	1574	1	-	0/16/26/28	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1574	PIX	C19-N24	2.49	1.36	1.32

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1574	PIX	C19-N24	2.44	1.36	1.32

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1574	PIX	C23-N24-C19	5.95	121.86	116.63
2	A	1574	PIX	C23-N24-C19	4.33	120.44	116.63
2	B	1574	PIX	C24-C23-N24	-3.69	119.81	123.34
2	A	1574	PIX	C4-N3-C2	3.63	120.00	114.08
2	B	1574	PIX	C20-C19-N24	-3.23	120.05	124.87
2	B	1574	PIX	C4-N3-C2	3.19	119.27	114.08
2	B	1574	PIX	C1-C2-N3	-3.08	105.66	110.27
2	A	1574	PIX	C24-C23-N24	-2.72	120.75	123.34
2	B	1574	PIX	O18-C19-C20	2.58	120.53	115.35
2	A	1574	PIX	C20-C19-N24	-2.57	121.04	124.87
2	A	1574	PIX	O18-C19-C20	2.49	120.34	115.35
2	A	1574	PIX	C2-N3-C8	-2.37	119.66	122.66
2	B	1574	PIX	C4-N3-C8	-2.12	119.97	122.66

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1574	PIX	N24-C19-O18-C16


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1574	PIX	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.



## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	541/543 (99%)	-0.04	7 (1%) 77 72	30, 34, 37, 42	0
1	B	543/543 (100%)	-0.13	7 (1%) 77 72	30, 34, 37, 43	0
All	All	1084/1086 (99%)	-0.09	14 (1%) 77 72	30, 34, 37, 43	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	33	GLY	6.2
1	B	31	TRP	4.7
1	B	126	SER	3.5
1	A	69	LEU	2.7
1	B	37	ALA	2.7
1	A	522	LEU	2.7
1	B	35	GLN	2.6
1	A	131	GLN	2.3
1	B	32	THR	2.2
1	A	524	LYS	2.2
1	B	68	ASP	2.2
1	B	33	GLY	2.2
1	A	240	GLY	2.1
1	A	241	SER	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NA	B	1576	1/1	0.73	0.18	29,29,29,29	0
4	NA	A	1576	1/1	0.90	0.17	27,27,27,27	0
4	NA	B	1575	1/1	0.92	0.16	31,31,31,31	0
3	CL	A	1575	1/1	0.93	0.09	23,23,23,23	0
2	PIX	A	1574	26/27	0.96	0.18	29,32,35,36	0
2	PIX	B	1574	26/27	0.97	0.14	25,27,29,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.