



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 12, 2024 – 10:24 pm BST

PDB ID : 2X2U  
Title : First two Cadherin-like domains from Human RET  
Authors : Kjaer, S.; Hanrahan, S.; Purkiss-Trew, A.G.; Totty, N.; McDonald, N.Q.  
Deposited on : 2010-01-15  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : **FAILED**  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	A	246	
2	B	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	UNX	A	1287	-	-	X	-
3	UNX	A	1290	-	-	X	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	UNX	A	1291	-	-	X	-
5	BU1	A	1294	-	-	X	-

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 2237 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTO-ONCOGENE TYROSINE-PROTEIN KINASE RECEPTOR RET.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	236	1928	1248	329	344	7	0	2	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	87	ARG	CYS	engineered mutation	UNP P07949
A	98	GLN	ASN	engineered mutation	UNP P07949
A	199	GLN	ASN	engineered mutation	UNP P07949
A	216	SER	CYS	engineered mutation	UNP P07949
A	271	GLU	-	expression tag	UNP P07949
A	272	PHE	-	expression tag	UNP P07949
A	273	GLU	-	expression tag	UNP P07949
A	274	ASN	-	expression tag	UNP P07949

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	2	25	14	2	9	0	0	0

- Molecule 3 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	11	Total	X	0	0
			11	11		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	1
			10	8	2		

- Molecule 5 is 1,4-BUTANEDIOL (three-letter code: BU1) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 4 2	0	0
5	A	1	Total C O 6 4 2	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	216	Total O 216 216	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.

- Molecule 1: PROTO-ONCOGENE TYROSINE-PROTEIN KINASE RECEPTOR RET



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.07Å 86.07Å 110.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.86 – 2.00	Depositor
% Data completeness (in resolution range)	99.7 (33.86-2.00)	Depositor
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.49 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.181 , 0.228	Depositor
Wilson B-factor (Å <sup>2</sup> )	30.0	Xtrriage
Anisotropy	0.024	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2237	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.79% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, UNX, BU1, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.53	7/1997 (0.4%)	1.25	11/2719 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	251	GLU	CB-CG	-7.96	1.37	1.52
1	A	174	PHE	CE1-CZ	6.67	1.50	1.37
1	A	187	GLN	CB-CG	-6.31	1.35	1.52
1	A	241	ALA	CA-CB	5.54	1.64	1.52
1	A	205	ARG	CZ-NH2	5.42	1.40	1.33
1	A	205	ARG	CG-CD	5.33	1.65	1.51
1	A	139	TRP	CB-CG	-5.15	1.41	1.50

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	205	ARG	NE-CZ-NH2	11.12	125.86	120.30
1	A	144	ARG	NE-CZ-NH2	9.48	125.04	120.30
1	A	144	ARG	NE-CZ-NH1	-8.55	116.03	120.30
1	A	205	ARG	NE-CZ-NH1	-7.58	116.51	120.30
1	A	230	ASP	CB-CG-OD1	6.77	124.39	118.30
1	A	189	ARG	NE-CZ-NH2	6.32	123.46	120.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	230	ASP	CB-CG-OD2	-6.06	112.85	118.30
1	A	189	ARG	NE-CZ-NH1	-5.75	117.42	120.30
1	A	267	ASP	CB-CG-OD1	5.66	123.39	118.30
1	A	114	ARG	NE-CZ-NH2	5.28	122.94	120.30
1	A	77	ARG	NE-CZ-NH1	-5.07	117.76	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	206	LEU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1928	0	1822	26	0
2	B	25	0	21	0	0
3	A	11	0	0	6	0
4	A	45	0	0	3	0
5	A	12	0	20	10	0
6	A	216	0	0	6	0
All	All	2237	0	1863	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87[B]:ARG:NH2	6:A:2060:HOH:O	1.94	0.99
3:A:1283:UNX:UNK	3:A:1284:UNX:UNK	1.00	0.91
3:A:1285:UNX:UNK	3:A:1287:UNX:UNK	1.24	0.82
1:A:54:HIS:NE2	4:A:1282[A]:SO4:O3	2.29	0.65
1:A:164:GLU:HB3	5:A:1293:BU1:H31	1.77	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1287:UNX:UNK	3:A:1290:UNX:UNK	1.45	0.61
1:A:37:TRP:HD1	5:A:1294:BU1:H11	1.68	0.58
1:A:37:TRP:CD1	5:A:1294:BU1:H11	2.38	0.58
1:A:168:PRO:HA	5:A:1294:BU1:H32	1.87	0.57
3:A:1291:UNX:UNK	3:A:1292:UNX:UNK	1.49	0.57
1:A:42:VAL:O	1:A:43:ASP:HB2	2.05	0.55
1:A:168:PRO:HB3	5:A:1294:BU1:H32	1.89	0.54
1:A:168:PRO:CA	5:A:1294:BU1:H32	2.36	0.54
1:A:79:ARG:NH2	6:A:2053:HOH:O	2.27	0.54
1:A:155:PRO:HG2	1:A:160:LEU:HD21	1.92	0.51
1:A:102:ASP:HB2	6:A:2015:HOH:O	2.10	0.51
1:A:265:GLU:HG3	6:A:2125:HOH:O	2.10	0.50
1:A:209:GLY:O	1:A:212:LEU:HB2	2.12	0.49
1:A:168:PRO:CB	5:A:1294:BU1:H32	2.43	0.48
1:A:169:GLU:H	5:A:1294:BU1:H21	1.78	0.47
1:A:39:LYS:NZ	5:A:1293:BU1:H21	2.29	0.47
3:A:1287:UNX:UNK	3:A:1288:UNX:UNK	1.59	0.46
1:A:144:ARG:HH11	1:A:144:ARG:HD2	1.58	0.46
1:A:52:TYR:CD1	4:A:1282[B]:SO4:O1	2.72	0.42
1:A:139:TRP:CG	1:A:140:PRO:HA	2.54	0.42
3:A:1290:UNX:UNK	3:A:1291:UNX:UNK	1.64	0.42
1:A:42:VAL:HG11	1:A:103:HIS:CE1	2.55	0.41
1:A:217:ALA:O	1:A:218:PRO:C	2.57	0.41
1:A:169:GLU:H	5:A:1294:BU1:C2	2.33	0.41
4:A:1277:SO4:O4	6:A:2203:HOH:O	2.19	0.41
1:A:210:GLU:H	1:A:210:GLU:CD	2.25	0.41
1:A:29:LEU:N	1:A:58:ASP:OD2	2.54	0.40
1:A:196:LEU:HD11	6:A:2019:HOH:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	232/246 (94%)	226 (97%)	6 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/221 (92%)	194 (95%)	10 (5%)	21	18

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	GLU
1	A	67	ARG
1	A	89	GLN
1	A	127	LEU
1	A	171	ARG
1	A	180	ARG
1	A	205	ARG
1	A	230	ASP
1	A	238	GLU
1	A	265	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	246	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	1,2	14,14,15	1.34	3 (21%)	17,19,21	3.04	8 (47%)
2	NAG	B	2	2	11,11,15	0.85	1 (9%)	12,15,21	2.13	2 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/2/19/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	NAG	C1-C2	2.87	1.56	1.52
2	B	1	NAG	C2-N2	2.34	1.50	1.46
2	B	2	NAG	C1-C2	2.14	1.54	1.52
2	B	1	NAG	C4-C3	2.05	1.57	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	C1-O5-C5	8.54	123.77	112.19
2	B	2	NAG	C1-O5-C5	6.10	120.45	112.19

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	C2-N2-C7	4.68	129.56	122.90
2	B	1	NAG	O5-C1-C2	4.43	118.28	111.29
2	B	1	NAG	C3-C4-C5	3.33	116.18	110.24
2	B	2	NAG	O5-C5-C6	3.11	112.08	107.20
2	B	1	NAG	O5-C5-C4	2.37	116.60	110.83
2	B	1	NAG	O4-C4-C3	2.28	115.63	110.35
2	B	1	NAG	O7-C7-C8	-2.24	117.90	122.06
2	B	1	NAG	C1-C2-N2	2.10	114.08	110.49

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	2	NAG	O5-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6
2	B	1	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 11 are unknown - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	SO4	A	1275	-	4,4,4	1.09	1 (25%)	6,6,6	1.60	1 (16%)
4	SO4	A	1276	-	4,4,4	0.39	0	6,6,6	1.51	1 (16%)
4	SO4	A	1279	-	4,4,4	0.29	0	6,6,6	0.43	0
4	SO4	A	1281	-	4,4,4	0.18	0	6,6,6	1.18	0
4	SO4	A	1278	-	4,4,4	0.46	0	6,6,6	0.62	0
5	BU1	A	1294	-	5,5,5	0.83	0	4,4,4	1.44	1 (25%)
4	SO4	A	1277	-	4,4,4	0.31	0	6,6,6	0.40	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	SO4	A	1280	-	4,4,4	0.27	0	6,6,6	0.40	0
4	SO4	A	1282[A]	-	4,4,4	0.45	0	6,6,6	0.50	0
5	BU1	A	1293	-	5,5,5	1.47	1 (20%)	4,4,4	2.34	1 (25%)
4	SO4	A	1282[B]	-	4,4,4	0.28	0	6,6,6	1.68	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BU1	A	1293	-	-	2/3/3/3	-
5	BU1	A	1294	-	-	2/3/3/3	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1293	BU1	O6-C4	2.29	1.53	1.42
4	A	1275	SO4	O1-S	2.12	1.57	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1293	BU1	C3-C2-C1	-4.55	84.21	114.02
5	A	1294	BU1	C3-C2-C1	-2.40	98.31	114.02
4	A	1275	SO4	O3-S-O1	2.28	121.19	109.31
4	A	1276	SO4	O4-S-O3	-2.27	99.38	109.06
4	A	1282[B]	SO4	O3-S-O2	-2.11	98.31	109.31

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1293	BU1	O5-C1-C2-C3
5	A	1293	BU1	C2-C3-C4-O6
5	A	1294	BU1	C2-C3-C4-O6
5	A	1294	BU1	O5-C1-C2-C3

There are no ring outliers.

5 monomers are involved in 13 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1294	BU1	8	0
4	A	1277	SO4	1	0
4	A	1282[A]	SO4	1	0
5	A	1293	BU1	2	0
4	A	1282[B]	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.


### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS failed to run properly - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



### 6.4 Ligands [i](#)

EDS failed to run properly - this section is therefore empty.

## 6.5 Other polymers

EDS failed to run properly - this section is therefore empty.