



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 6, 2024 – 10:47 am BST

PDB ID : 2X93
Title : Crystal structure of AnCE-trandolaprilat complex
Authors : Akif, M.; Georgiadis, D.; Mahajan, A.; Dive, V.; Sturrock, E.D.; Isaac, R.E.; Acharya, K.R.
Deposited on : 2010-03-14
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1356 (1.98-1.98)
Clashscore	180529	1437 (1.98-1.98)
Ramachandran outliers	177936	1426 (1.98-1.98)
Sidechain outliers	177891	1426 (1.98-1.98)
RSRZ outliers	164620	1356 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	599	 6% 95% 5%
2	B	6	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EPE	A	1240	-	-	X	-

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 5564 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ANGIOTENSIN CONVERTING ENZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	598	4883	3124	806	933	20	0	2	0

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	6	72	40	2	30	0	0	0

- Molecule 3 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	15	8	2	4	1	0	0
3	A	1	15	8	2	4	1	0	0

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
4	A	1	1	1	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
5	A	1	14	8	1	5	0	0

- Molecule 6 is TRANDOLAPRILAT (three-letter code: X93) (formula: $C_{22}H_{30}N_2O_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	A	1	29	22	2	5	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	521	Total 521	O 521	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ANGIOTENSIN CONVERTING ENZYME



- Molecule 2: beta-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics i

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	172.86Å 172.86Å 103.25Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.76 – 1.98 28.76 – 1.98	Depositor EDS
% Data completeness (in resolution range)	97.6 (28.76-1.98) 94.4 (28.76-1.98)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 1.98Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.190 , 0.207 0.190 , 0.206	Depositor DCC
R_{free} test set	3746 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	24.8	Xtriage
Anisotropy	0.012	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 41.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.008 for $-2/3^*h-1/3^*k-4/3^*l,-1/3^*h-2/3^*k+4/3^*l,-1/3^*h+1/3^*k+1/3^*l$ 0.008 for $-h,1/3^*h-1/3^*k-4/3^*l,-1/3^*h-2/3^*k+1/3^*l$ 0.006 for $-1/3^*h+1/3^*k+4/3^*l,-k,2/3^*h+1/3^*k+1/3^*l$ 0.006 for $-h,2/3^*h+1/3^*k+4/3^*l,1/3^*h+2/3^*k-1/3^*l$ 0.011 for $-1/3^*h-2/3^*k+4/3^*l,-2/3^*h-1/3^*k-4/3^*l,1/3^*h-1/3^*k-1/3^*l$ 0.008 for $1/3^*h+2/3^*k-4/3^*l,-k,-2/3^*h-1/3^*k-1/3^*l$ 0.013 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5564	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, X93, EPE, ZN, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.40	2/5015 (0.0%)	0.47	0/6796

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	346	THR	CB-OG1	5.46	1.54	1.43
1	A	607	THR	CB-OG1	5.32	1.53	1.43

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4883	0	4677	15	0
2	B	72	0	61	0	0
3	A	30	0	34	8	0
4	A	1	0	0	0	0
5	A	28	0	26	0	0
6	A	29	0	28	1	0
7	A	521	0	0	3	0
All	All	5564	0	4826	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1240:EPE:H81	7:A:2504:HOH:O	1.78	0.81
1:A:506:ARG:HH12	3:A:1240:EPE:C2	2.05	0.69
1:A:506:ARG:HH22	3:A:1240:EPE:H22	1.58	0.68
3:A:1240:EPE:H72	7:A:2430:HOH:O	1.95	0.66
1:A:276:PRO:HB3	1:A:592:TRP:CH2	2.36	0.61
1:A:178:ARG:NH1	7:A:2164:HOH:O	2.39	0.54
1:A:506:ARG:HH12	3:A:1240:EPE:H21	1.74	0.51
1:A:306:PHE:CD2	1:A:401:LEU:HD13	2.47	0.49
1:A:502:VAL:HG22	3:A:1240:EPE:H71	1.93	0.49
1:A:296:PRO:HG2	1:A:326:GLU:HG2	1.94	0.49
3:A:1240:EPE:H31	6:A:1623:X93:HAH	1.95	0.48
1:A:133:CYS:HA	1:A:141:CYS:HA	1.98	0.44
1:A:395:GLU:HB2	1:A:510:SER:HB2	2.00	0.43
1:A:280:LYS:HB3	1:A:281:PRO:HD2	2.02	0.42
1:A:506:ARG:NH2	3:A:1240:EPE:H22	2.31	0.41
1:A:61:LYS:O	1:A:65:ILE:HG12	2.20	0.41
1:A:146:ASP:HA	1:A:147:PRO:HA	1.82	0.41
1:A:266:GLN:NE2	1:A:437:ASP:OD1	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	598/599 (100%)	591 (99%)	7 (1%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	521/521 (100%)	515 (99%)	6 (1%)	67 65

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	245	ASP
1	A	374	TYR
1	A	378	TYR
1	A	394	HIS
1	A	446	PHE
1	A	540	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	266	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	2,1	14,14,15	0.52	0	17,19,21	0.62	0
2	NAG	B	2	2	14,14,15	0.47	0	17,19,21	1.36	2 (11%)
2	BMA	B	3	2	11,11,12	0.64	0	15,15,17	0.93	0
2	MAN	B	4	2	11,11,12	0.62	0	15,15,17	0.70	0
2	BMA	B	5	2	11,11,12	0.58	0	15,15,17	1.33	2 (13%)
2	MAN	B	6	2	11,11,12	0.59	0	15,15,17	0.92	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	3/6/23/26	0/1/1/1
2	BMA	B	3	2	-	1/2/19/22	0/1/1/1
2	MAN	B	4	2	-	2/2/19/22	0/1/1/1
2	BMA	B	5	2	-	2/2/19/22	0/1/1/1
2	MAN	B	6	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	5	BMA	C1-O5-C5	3.83	117.39	112.19
2	B	2	NAG	C2-N2-C7	3.16	127.40	122.90
2	B	2	NAG	C1-O5-C5	2.89	116.10	112.19
2	B	6	MAN	C1-O5-C5	2.45	115.51	112.19
2	B	5	BMA	C1-C2-C3	2.42	112.64	109.67

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	4	MAN	C4-C5-C6-O6

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	5	BMA	O5-C5-C6-O6
2	B	4	MAN	O5-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6
2	B	5	BMA	C4-C5-C6-O6
2	B	2	NAG	C3-C2-N2-C7
2	B	2	NAG	O5-C5-C6-O6
2	B	3	BMA	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EPE	A	1241	-	15,15,15	0.83	1 (6%)	18,20,20	1.94	6 (33%)
5	NAG	A	1622	1	14,14,15	0.58	0	17,19,21	0.69	0
6	X93	A	1623	4	31,31,31	0.81	0	40,43,43	1.42	4 (10%)
5	NAG	A	1621	1	14,14,15	0.55	0	17,19,21	0.66	0
3	EPE	A	1240	-	15,15,15	0.84	1 (6%)	18,20,20	1.93	6 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EPE	A	1241	-	-	2/9/19/19	0/1/1/1
5	NAG	A	1622	1	-	2/6/23/26	0/1/1/1
6	X93	A	1623	4	-	4/25/48/48	0/3/3/3
5	NAG	A	1621	1	-	2/6/23/26	0/1/1/1
3	EPE	A	1240	-	-	8/9/19/19	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1240	EPE	C10-S	2.85	1.81	1.77
3	A	1241	EPE	C10-S	2.81	1.81	1.77

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1241	EPE	C5-N4-C3	4.94	119.95	108.83
3	A	1240	EPE	C5-N4-C3	4.94	119.94	108.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	1623	X93	CAP-CBB-CAZ	-4.80	106.80	113.04
3	A	1241	EPE	C7-N4-C3	3.41	119.95	111.23
3	A	1240	EPE	C7-N4-C3	3.40	119.93	111.23
3	A	1240	EPE	C7-N4-C5	3.39	119.90	111.23
3	A	1241	EPE	C7-N4-C5	3.38	119.87	111.23
6	A	1623	X93	O-C-NBC	-3.12	116.94	121.68
6	A	1623	X93	CBA-NBC-CBB	-2.61	107.98	112.22
3	A	1240	EPE	O3S-S-C10	2.29	109.48	105.77
3	A	1241	EPE	O3S-S-C10	2.27	109.44	105.77
3	A	1241	EPE	O1S-S-C10	2.13	109.48	106.92
3	A	1241	EPE	O2S-S-C10	2.13	109.48	106.92
3	A	1240	EPE	O2S-S-C10	2.11	109.45	106.92
3	A	1240	EPE	O1S-S-C10	2.10	109.45	106.92
6	A	1623	X93	CAO-CAZ-CBB	-2.07	108.92	113.73

There are no chirality outliers.

All (18) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1240	EPE	C8-C7-N4-C5
3	A	1240	EPE	S-C10-C9-N1
3	A	1241	EPE	C8-C7-N4-C5
5	A	1621	NAG	O5-C5-C6-O6
5	A	1622	NAG	O5-C5-C6-O6
3	A	1240	EPE	N4-C7-C8-O8
3	A	1240	EPE	C9-C10-S-O3S
3	A	1241	EPE	N4-C7-C8-O8
5	A	1622	NAG	C4-C5-C6-O6
5	A	1621	NAG	C4-C5-C6-O6
3	A	1240	EPE	C9-C10-S-O1S
3	A	1240	EPE	C9-C10-S-O2S
6	A	1623	X93	CAN-CAQ-CAY-CAT
3	A	1240	EPE	C10-C9-N1-C2
3	A	1240	EPE	C10-C9-N1-C6
6	A	1623	X93	CAQ-CAN-CAW-CAK
6	A	1623	X93	CAQ-CAN-CAW-CAJ
6	A	1623	X93	CAN-CAQ-CAY-N

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1623	X93	1	0
3	A	1240	EPE	8	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	598/599 (99%)	0.28	34 (5%) 30 42	14, 24, 35, 43	9 (1%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	139	THR	5.9
1	A	331	GLY	5.5
1	A	17	ALA	4.8
1	A	614	SER	4.5
1	A	19	VAL	4.4
1	A	329	THR	3.5
1	A	141	CYS	3.3
1	A	461	ASP	3.3
1	A	18	LEU	3.2
1	A	437	ASP	3.2
1	A	318	ASP	3.0
1	A	330	ASP	2.9
1	A	57	GLU	2.8
1	A	140	LYS	2.8
1	A	137	ASP	2.7
1	A	30	GLU	2.7
1	A	61	LYS	2.7
1	A	78	SER	2.6
1	A	81	THR	2.5
1	A	467	CYS	2.4
1	A	404	SER	2.4
1	A	360	ASP	2.4
1	A	526	GLY	2.4
1	A	547	ALA	2.3
1	A	75	GLU	2.3
1	A	530	PRO	2.2
1	A	24	GLN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	179	SER	2.2
1	A	533	VAL	2.2
1	A	531	ASP	2.1
1	A	82	LYS	2.1
1	A	64	GLU	2.1
1	A	73	MET	2.1
1	A	136	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MAN	B	6	11/12	0.66	0.20	56,58,58,59	0
2	BMA	B	5	11/12	0.73	0.23	55,56,57,58	0
2	BMA	B	3	11/12	0.86	0.16	45,48,51,54	0
2	NAG	B	2	14/15	0.88	0.14	32,36,38,41	0
2	MAN	B	4	11/12	0.89	0.13	51,51,53,54	0
2	NAG	B	1	14/15	0.94	0.09	22,24,26,29	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	NAG	A	1622	14/15	0.73	0.32	44,45,45,45	14
3	EPE	A	1240	15/15	0.77	0.26	41,44,47,47	9
5	NAG	A	1621	14/15	0.81	0.23	32,33,34,34	14
3	EPE	A	1241	15/15	0.84	0.18	46,47,49,49	0
6	X93	A	1623	29/29	0.94	0.08	19,20,25,25	0
4	ZN	A	1615	1/1	1.00	0.02	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.