



Full wwPDB X-ray Structure Validation Report

Jun 12, 2024 – 07:15 PM EDT

PDB ID : 2Y8L
Title : Structure of the regulatory fragment of mammalian aMPK in complex with two ADP
Authors : Xiao, B.; Sanders, M.J.; Underwood, E.; Heath, R.; Mayer, F.; Carmena, D.; Jing, C.; Walker, P.A.; Eccleston, J.F.; Haire, L.F.; Saiu, P.; Howell, S.A.; Aasland, R.; Martin, S.R.; Carling, D.; Gamblin, S.J.
Deposited on : 2011-02-07
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 1.20.1
EDS : 2.36.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	173	
2	B	87	
3	E	330	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4045 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 5'-AMP-ACTIVATED PROTEIN KINASE CATALYTIC SUB-UNIT ALPHA-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	104	855	539	152	157	7	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	378	MET	-	expression tag	UNP P54645
A	379	SER	-	expression tag	UNP P54645
A	380	HIS	-	expression tag	UNP P54645
A	381	HIS	-	expression tag	UNP P54645
A	382	HIS	-	expression tag	UNP P54645
A	383	HIS	-	expression tag	UNP P54645
A	384	HIS	-	expression tag	UNP P54645
A	385	HIS	-	expression tag	UNP P54645
A	386	SER	-	expression tag	UNP P54645
A	387	GLY	-	expression tag	UNP P54645
A	388	LEU	-	expression tag	UNP P54645
A	389	VAL	-	expression tag	UNP P54645
A	390	PRO	-	expression tag	UNP P54645
A	391	ARG	-	expression tag	UNP P54645
A	392	GLY	-	expression tag	UNP P54645
A	393	SER	-	cloning artifact	UNP P54645
A	394	MET	-	cloning artifact	UNP P54645
A	395	ALA	-	cloning artifact	UNP P54645
A	545	ASN	-	cloning artifact	UNP P54645
A	546	SER	-	cloning artifact	UNP P54645
A	547	CYS	-	cloning artifact	UNP P54645
A	548	THR	-	cloning artifact	UNP P54645
A	549	VAL	-	cloning artifact	UNP P54645
A	550	ASN	-	cloning artifact	UNP P54645

- Molecule 2 is a protein called 5'-AMP-ACTIVATED PROTEIN KINASE SUBUNIT BETA-

2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	69	570	374	97	96	3	0	0	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	186	MET	-	cloning artifact	UNP O43741

- Molecule 3 is a protein called 5'-AMP-ACTIVATED PROTEIN KINASE SUBUNIT GAMMA-1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	305	2442	1584	408	443	7	0	0	1

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	E	1	27	10	5	10	2	0	0
4	E	1	27	10	5	10	2	0	0

- Molecule 5 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: $C_{10}H_{14}N_5O_7P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	E	1	23	10	5	7	1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	23	Total	O	0	0
			23	23		
6	B	17	Total	O	0	0
			17	17		
6	E	61	Total	O	0	0
			61	61		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-AMP-ACTIVATED PROTEIN KINASE CATALYTIC SUBUNIT ALPHA-1



- Molecule 2: 5'-AMP-ACTIVATED PROTEIN KINASE SUBUNIT BETA-2



- Molecule 3: 5'-AMP-ACTIVATED PROTEIN KINASE SUBUNIT GAMMA-1



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.57Å 121.46Å 125.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	87.42 – 2.50 19.93 – 2.50	Depositor EDS
% Data completeness (in resolution range)	97.8 (87.42-2.50) 98.0 (19.93-2.50)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.73 (at 2.50Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.228 , 0.253 0.224 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	50.8	Xtrriage
Anisotropy	0.220	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.014 for -h,l,k	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4045	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AMP, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.49	1/873 (0.1%)	0.66	1/1178 (0.1%)
2	B	0.45	0/585	0.67	0/789
3	E	0.51	0/2494	0.65	0/3386
All	All	0.50	1/3952 (0.0%)	0.65	1/5353 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
3	E	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	547	CYS	CB-SG	-5.04	1.73	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	460	LEU	CA-CB-CG	6.86	131.08	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	209	PRO	Peptide
3	E	182	PHE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	855	0	845	24	0
2	B	570	0	595	9	0
3	E	2442	0	2506	25	0
4	E	54	0	24	0	0
5	E	23	0	12	0	0
6	A	23	0	0	0	0
6	B	17	0	0	0	0
6	E	61	0	0	0	0
All	All	4045	0	3982	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:208:LEU:O	2:B:210:PRO:HD3	1.44	1.14
1:A:416:ALA:HB2	1:A:546:SER:HB3	1.70	0.72
3:E:84:MET:HE3	3:E:153:PRO:HG3	1.77	0.65
2:B:208:LEU:O	2:B:210:PRO:CD	2.36	0.65
1:A:543:ILE:O	1:A:547:CYS:HB2	1.98	0.64
1:A:428:ASN:HB2	1:A:431:TYR:HB3	1.79	0.63
1:A:447:MET:HE3	1:A:542:LEU:HD12	1.80	0.62
2:B:200:GLU:HG3	2:B:203:LYS:HB2	1.83	0.61
3:E:108:LEU:O	3:E:111:HIS:HB2	2.03	0.59
3:E:282:GLU:HB2	3:E:287:ILE:HD11	1.85	0.59
1:A:417:ILE:HG23	1:A:422:TYR:HB2	1.85	0.57
1:A:532:THR:H	3:E:161:ASN:HD21	1.53	0.57
3:E:55:GLN:HB2	3:E:58:LYS:HD2	1.88	0.56
1:A:431:TYR:OH	1:A:433:ARG:NH1	2.38	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:447:MET:CE	1:A:542:LEU:HD12	2.37	0.54
1:A:393:SER:N	3:E:65:THR:HG22	2.23	0.54
1:A:409:ILE:O	1:A:413:VAL:HG13	2.08	0.53
3:E:57:LYS:HD2	3:E:109:GLU:HG3	1.89	0.53
3:E:193:GLU:OE1	3:E:280:LEU:HD22	2.08	0.53
3:E:56:VAL:HG13	3:E:60:PHE:CE2	2.44	0.53
3:E:109:GLU:OE1	3:E:109:GLU:HA	2.09	0.53
3:E:60:PHE:CE1	3:E:90:PHE:HB2	2.44	0.52
3:E:84:MET:CE	3:E:153:PRO:HG3	2.40	0.51
1:A:526:PRO:HG3	3:E:128:LEU:HD23	1.93	0.50
2:B:235:HIS:O	2:B:237:MET:HG2	2.10	0.50
1:A:396:TRP:CZ3	2:B:254:SER:HB2	2.47	0.50
1:A:532:THR:H	3:E:161:ASN:ND2	2.10	0.49
1:A:396:TRP:HB2	2:B:216:ILE:HB	1.93	0.49
1:A:405:ARG:HB3	1:A:407:ASN:OD1	2.13	0.48
3:E:47:LEU:HD21	3:E:162:THR:HB	1.97	0.47
3:E:246:ILE:H	3:E:246:ILE:HG13	1.61	0.47
3:E:283:THR:O	3:E:287:ILE:HD12	2.15	0.46
1:A:422:TYR:CE1	1:A:436:ARG:HG3	2.50	0.46
1:A:416:ALA:CB	1:A:546:SER:HB3	2.42	0.46
1:A:528:PRO:HA	3:E:80:SER:HB3	1.98	0.46
1:A:410:MET:HA	1:A:410:MET:CE	2.46	0.45
1:A:417:ILE:HG13	1:A:542:LEU:HD21	1.99	0.45
3:E:84:MET:HE3	3:E:128:LEU:HD11	1.99	0.44
3:E:54:LEU:HD13	3:E:59:ALA:HB2	2.00	0.43
1:A:543:ILE:O	1:A:547:CYS:CB	2.66	0.42
2:B:236:VAL:HG13	2:B:236:VAL:O	2.18	0.42
3:E:278:CYS:HB3	3:E:299:LEU:HD13	2.02	0.41
3:E:56:VAL:HG13	3:E:60:PHE:HE2	1.85	0.41
2:B:235:HIS:O	2:B:237:MET:N	2.54	0.41
1:A:533:ILE:HG21	3:E:74:TRP:CD2	2.55	0.41
3:E:61:PHE:O	3:E:65:THR:OG1	2.30	0.41
1:A:407:ASN:OD1	1:A:407:ASN:N	2.53	0.40
1:A:393:SER:HB2	2:B:244:LEU:HD21	2.04	0.40
3:E:108:LEU:HA	3:E:111:HIS:ND1	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	100/173 (58%)	98 (98%)	2 (2%)	0	100	100
2	B	65/87 (75%)	57 (88%)	7 (11%)	1 (2%)	10	18
3	E	303/330 (92%)	287 (95%)	14 (5%)	2 (1%)	22	39
All	All	468/590 (79%)	442 (94%)	23 (5%)	3 (1%)	25	43

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	236	VAL
3	E	122	GLN
3	E	182	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	97/157 (62%)	92 (95%)	5 (5%)	23	44
2	B	65/81 (80%)	62 (95%)	3 (5%)	27	50
3	E	277/299 (93%)	248 (90%)	29 (10%)	7	13
All	All	439/537 (82%)	402 (92%)	37 (8%)	11	21

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	413	VAL
1	A	420	LEU
1	A	433	ARG
1	A	460	LEU
1	A	547	CYS
2	B	201	ARG
2	B	237	MET
2	B	254	SER
3	E	24	ASN
3	E	26	SER
3	E	29	THR
3	E	40	LEU
3	E	54	LEU
3	E	94	LEU
3	E	103	VAL
3	E	109	GLU
3	E	110	GLU
3	E	123	ASP
3	E	124	SER
3	E	129	VAL
3	E	196	GLN
3	E	224	VAL
3	E	232	GLU
3	E	246	ILE
3	E	247	ASN
3	E	261	VAL
3	E	267	HIS
3	E	270	HIS
3	E	275	VAL
3	E	284	LEU
3	E	291	LEU
3	E	296	VAL
3	E	298	ARG
3	E	302	VAL
3	E	315	SER
3	E	318	LEU
3	E	322	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	397	HIS
3	E	24	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	E	122	GLN
3	E	147	ASN
3	E	161	ASN
3	E	168	HIS
3	E	222	HIS
3	E	266	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ADP	E	1327	-	24,29,29	0.96	1 (4%)	29,45,45	1.55	5 (17%)
5	AMP	E	1329	-	21,25,25	0.81	0	23,38,38	1.49	5 (21%)
4	ADP	E	1328	-	24,29,29	0.85	0	29,45,45	1.35	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	E	1327	-	-	4/12/32/32	0/3/3/3
5	AMP	E	1329	-	-	0/6/26/26	0/3/3/3
4	ADP	E	1328	-	-	7/12/32/32	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	E	1327	ADP	O4'-C1'	2.31	1.43	1.40

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	1328	ADP	N3-C2-N1	-4.44	122.64	128.67
4	E	1327	ADP	N3-C2-N1	-4.44	122.65	128.67
5	E	1329	AMP	N3-C2-N1	-4.36	122.75	128.67
4	E	1327	ADP	O3A-PA-O1A	-2.88	102.04	110.70
4	E	1327	ADP	O3B-PB-O2B	2.58	117.47	107.80
5	E	1329	AMP	C4'-O4'-C1'	2.51	112.23	109.92
4	E	1327	ADP	O2A-PA-O3A	2.37	113.69	107.27
5	E	1329	AMP	C4-C5-N7	-2.34	106.86	109.34
4	E	1327	ADP	C4-C5-N7	-2.31	106.90	109.34
5	E	1329	AMP	O5'-P-O1P	-2.22	100.43	106.44
4	E	1328	ADP	C4'-O4'-C1'	2.05	111.80	109.92
5	E	1329	AMP	O3P-P-O2P	2.03	115.41	107.80

There are no chirality outliers.


All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	E	1328	ADP	PA-O3A-PB-O3B
4	E	1328	ADP	C5'-O5'-PA-O1A
4	E	1328	ADP	C5'-O5'-PA-O2A
4	E	1328	ADP	C5'-O5'-PA-O3A
4	E	1328	ADP	O4'-C4'-C5'-O5'
4	E	1328	ADP	C3'-C4'-C5'-O5'
4	E	1327	ADP	PB-O3A-PA-O1A
4	E	1327	ADP	C5'-O5'-PA-O1A
4	E	1327	ADP	C5'-O5'-PA-O3A
4	E	1328	ADP	PA-O3A-PB-O1B
4	E	1327	ADP	PB-O3A-PA-O2A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	104/173 (60%)	0.37	8 (7%) 13 13	30, 54, 91, 111	0
2	B	69/87 (79%)	1.17	18 (26%) 0 0	32, 80, 102, 119	0
3	E	305/330 (92%)	0.39	32 (10%) 6 6	28, 50, 111, 199	0
All	All	478/590 (81%)	0.50	58 (12%) 4 4	28, 53, 105, 199	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	E	271	TYR	10.8
3	E	182	PHE	7.9
3	E	270	HIS	7.8
3	E	23	SER	6.9
3	E	125	PHE	6.8
3	E	253	THR	6.3
1	A	393	SER	6.1
1	A	394	MET	6.1
3	E	24	ASN	5.9
3	E	123	ASP	5.9
2	B	190	GLY	5.8
3	E	326	GLY	5.8
2	B	200	GLU	5.7
3	E	269	SER	5.6
2	B	235	HIS	5.3
3	E	25	SER	4.9
3	E	324	THR	4.7
2	B	209	PRO	4.7
2	B	217	LEU	4.6
1	A	525	ALA	4.6
2	B	210	PRO	4.4
3	E	245	VAL	4.3
2	B	201	ARG	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	E	250	ALA	4.1
3	E	252	LYS	4.1
2	B	218	ASN	4.0
3	E	267	HIS	4.0
2	B	234	ASN	3.8
1	A	468	ASP	3.8
3	E	124	SER	3.8
3	E	272	PHE	3.6
3	E	325	GLY	3.6
1	A	524	VAL	3.3
2	B	233	PRO	3.3
1	A	526	PRO	3.2
3	E	254	TYR	3.2
2	B	272	ILE	3.2
1	A	469	GLU	3.0
3	E	122	GLN	3.0
3	E	249	ALA	2.9
2	B	248	ASP	2.8
2	B	211	HIS	2.8
3	E	248	LEU	2.8
2	B	198	SER	2.7
3	E	120	TYR	2.7
3	E	327	GLU	2.7
2	B	199	GLU	2.6
2	B	197	ARG	2.6
2	B	203	LYS	2.5
1	A	528	PRO	2.4
2	B	236	VAL	2.3
3	E	273	GLU	2.3
3	E	251	GLU	2.3
3	E	118	GLU	2.2
3	E	321	LEU	2.2
3	E	126	LYS	2.2
3	E	246	ILE	2.0
3	E	97	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	ADP	E	1328	27/27	0.92	0.12	52,62,71,73	0
5	AMP	E	1329	23/23	0.95	0.12	37,44,47,49	0
4	ADP	E	1327	27/27	0.98	0.11	39,43,47,51	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around AMP E 1329:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.