



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 3, 2024 – 06:08 AM EST

PDB ID : 2ZTT  
Title : Crystal Structure of RNA polymerase PB1-PB2 subunits from Influenza A Virus  
Authors : Sugiyama, K.; Obayashi, E.; Park, S.-Y.  
Deposited on : 2008-10-08  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	79	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 58%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 30%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: grey;"></div> </div> <p style="font-size: small; margin-top: 5px;"> <span style="display: inline-block; width: 2%; height: 10px; background-color: red; margin-right: 5px;"></span> %            58% 30% 8%         </p>
1	C	79	<div style="display: flex; align-items: center;"> <div style="width: 47%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 37%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: grey;"></div> </div> <p style="font-size: small; margin-top: 5px;">47% 37% 6% 10%</p>
2	B	40	<div style="display: flex; align-items: center;"> <div style="width: 60%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 25%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: grey;"></div> </div> <p style="font-size: small; margin-top: 5px;">60% 25% 8% 8%</p>
2	D	40	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 52%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 32%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: grey;"></div> </div> <p style="font-size: small; margin-top: 5px;">2% 52% 32% 8% 8%</p>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1830 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-directed RNA polymerase catalytic subunit.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	73	606	376	109	114	3	4	0	0	0
1	C	71	585	364	102	112	3	4	0	0	0

- Molecule 2 is a protein called Polymerase basic protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
2	B	37	303	187	57	56	3	0	0	0
2	D	37	303	187	57	56	3	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLY	-	expression tag	UNP P03428
B	-1	GLY	-	expression tag	UNP P03428
B	0	SER	-	expression tag	UNP P03428
D	-2	GLY	-	expression tag	UNP P03428
D	-1	GLY	-	expression tag	UNP P03428
D	0	SER	-	expression tag	UNP P03428

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	15	Total	O	0	0
			15	15		
3	B	4	Total	O	0	0
			4	4		
3	C	8	Total	O	0	0
			8	8		

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
3	D	6	Total O 6 6	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RNA-directed RNA polymerase catalytic subunit

Chain A: 



- Molecule 1: RNA-directed RNA polymerase catalytic subunit


Chain C: 



GLN  
LYS


- Molecule 2: Polymerase basic protein 2

Chain B: 



- Molecule 2: Polymerase basic protein 2

Chain D: 



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	44.27Å 61.48Å 45.47Å 90.00° 103.35° 90.00°	Depositor
Resolution (Å)	20.00 – 2.10 20.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-2.10) 92.8 (20.00-2.10)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.65 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.232 , 0.272 0.231 , 0.273	Depositor DCC
$R_{free}$ test set	634 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.4	Xtrriage
Anisotropy	0.554	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 48.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.040 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	1830	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.95	0/610	0.97	0/802
1	C	1.05	0/589	0.92	0/777
2	B	1.03	0/301	1.07	0/394
2	D	1.02	0/301	1.07	0/394
All	All	1.01	0/1801	0.99	0/2367

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	606	0	606	22	0
1	C	585	0	578	42	0
2	B	303	0	329	11	0
2	D	303	0	329	17	0
3	A	15	0	0	0	0
3	B	4	0	0	0	0
3	C	8	0	0	0	0
3	D	6	0	0	1	0
All	All	1830	0	1842	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:688:MSE:HE1	1:C:691:ARG:HH21	1.17	1.06
1:C:715:VAL:HA	1:C:718:MSE:HE3	1.36	1.04
1:C:742:GLU:OE1	2:D:1:MSE:HE1	1.73	0.86
1:C:740:PHE:CE1	1:C:744:MSE:HE3	2.10	0.86
1:A:734:ARG:HG3	1:A:734:ARG:HH11	1.40	0.86
1:C:705:TYR:CD1	1:C:705:TYR:N	2.45	0.84
1:C:740:PHE:HE1	1:C:744:MSE:HE3	1.43	0.83
1:C:714:MSE:C	1:C:718:MSE:HE2	1.98	0.83
1:C:726:ALA:HB3	1:C:744:MSE:HE2	1.63	0.81
1:C:726:ALA:HB3	1:C:744:MSE:CE	2.09	0.81
1:C:705:TYR:H	1:C:705:TYR:HD1	1.30	0.75
1:C:714:MSE:O	1:C:718:MSE:HE2	1.88	0.73
1:A:695:LEU:HG	1:A:718:MSE:HE1	1.71	0.72
1:C:726:ALA:CB	1:C:744:MSE:HE2	2.20	0.71
2:D:28:MSE:HE2	2:D:32:LYS:HB2	1.72	0.71
1:A:699:PHE:CE2	1:A:718:MSE:HE2	2.25	0.71
1:C:699:PHE:CZ	2:D:3:ARG:HG3	2.26	0.71
1:A:695:LEU:HG	1:A:718:MSE:CE	2.21	0.69
1:C:741:THR:O	1:C:745:LYS:HG2	1.93	0.69
1:C:688:MSE:HE1	1:C:691:ARG:NH2	2.02	0.68
1:A:685:ASP:HA	1:A:687:GLN:HG3	1.77	0.67
1:C:709:VAL:HG13	1:C:712:SER:HB2	1.75	0.66
2:D:32:LYS:O	2:D:32:LYS:HG2	1.95	0.65
1:C:745:LYS:HB3	1:C:745:LYS:NZ	2.12	0.65
2:D:30:ILE:HG13	2:D:30:ILE:O	1.95	0.65
1:A:745:LYS:HD3	2:D:14:SER:H	1.61	0.64
2:D:28:MSE:CE	2:D:32:LYS:HB2	2.27	0.64
2:B:17:ARG:O	2:B:21:THR:HG23	1.98	0.64
1:A:750:ILE:HD13	2:B:7:LEU:HD13	1.78	0.64
1:A:734:ARG:HG3	1:A:734:ARG:NH1	2.07	0.63
1:C:715:VAL:HA	1:C:718:MSE:CE	2.22	0.63
1:C:688:MSE:SE	2:D:15:ARG:HG2	2.49	0.62
1:A:705:TYR:CD2	1:A:705:TYR:N	2.67	0.62
1:C:726:ALA:HB3	1:C:744:MSE:HE1	1.82	0.62
1:C:715:VAL:CA	1:C:718:MSE:HE3	2.22	0.61
1:C:740:PHE:CE1	1:C:744:MSE:CE	2.85	0.59
1:C:715:VAL:HG23	2:D:7:LEU:HD21	1.84	0.59
2:B:12:SER:HB3	1:C:746:ILE:HD12	1.87	0.57
1:C:737:LYS:O	1:C:741:THR:HG23	2.05	0.57
1:C:752:GLU:O	1:C:754:ARG:HD3	2.05	0.56

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:9:ASN:OD1	2:D:8:ARG:NH2	2.35	0.56
2:B:2:GLU:O	2:B:6:GLU:HG3	2.06	0.56
2:B:14:SER:HB3	1:C:745:LYS:HZ1	1.72	0.55
1:C:701:PRO:O	1:C:704:SER:HB2	2.06	0.55
1:C:745:LYS:HB3	1:C:745:LYS:HZ2	1.73	0.54
1:A:685:ASP:C	1:A:687:GLN:H	2.10	0.54
1:C:688:MSE:HA	1:C:688:MSE:HE2	1.90	0.53
2:B:33:LYS:O	2:B:33:LYS:HG2	2.09	0.53
2:B:33:LYS:O	2:B:33:LYS:CG	2.57	0.52
1:A:729:ASP:OD1	1:A:734:ARG:HD2	2.10	0.51
1:A:691:ARG:NE	2:B:10:LEU:HD11	2.26	0.51
1:C:714:MSE:C	1:C:718:MSE:CE	2.76	0.49
1:C:745:LYS:NZ	1:C:745:LYS:CB	2.77	0.47
1:C:709:VAL:HG13	1:C:712:SER:CB	2.42	0.47
1:C:719:VAL:HG13	1:C:747:CYS:HB3	1.95	0.47
1:A:699:PHE:HE2	1:A:718:MSE:HE2	1.75	0.47
1:C:739:GLU:HG3	2:D:1:MSE:HG2	1.97	0.46
1:A:689:TYR:CE2	1:A:693:CYS:SG	3.08	0.46
1:A:723:ARG:O	1:A:727:ARG:HD3	2.15	0.46
1:A:740:PHE:O	1:A:744:MSE:HG2	2.16	0.45
1:A:695:LEU:CG	1:A:718:MSE:HE1	2.44	0.45
2:D:5:LYS:CE	3:D:38:HOH:O	2.65	0.45
1:C:750:ILE:HD13	2:D:7:LEU:HD22	1.98	0.44
1:A:751:GLU:OE2	1:A:754:ARG:HD3	2.17	0.44
2:B:16:THR:O	2:B:20:LEU:HB2	2.19	0.43
1:C:729:ASP:O	1:C:734:ARG:HB2	2.19	0.43
1:A:713:SER:OG	1:A:716:GLU:HG2	2.19	0.42
1:C:685:ASP:HB3	2:D:15:ARG:HH21	1.85	0.42
1:A:696:PHE:HA	1:A:718:MSE:HE3	2.02	0.42
1:C:742:GLU:CD	2:D:1:MSE:HE1	2.36	0.42
2:D:22:LYS:HB3	2:D:22:LYS:NZ	2.34	0.41
2:D:2:GLU:O	2:D:6:GLU:HG3	2.21	0.41
1:C:716:GLU:O	1:C:717:ALA:C	2.59	0.41
1:A:745:LYS:HZ3	1:A:745:LYS:HG3	1.64	0.41
2:B:14:SER:H	1:C:745:LYS:HZ2	1.68	0.40
1:A:734:ARG:HH11	1:A:734:ARG:CG	2.19	0.40
1:C:695:LEU:HD12	1:C:698:LYS:HD3	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	71/79 (90%)	68 (96%)	3 (4%)	0	100	100
1	C	69/79 (87%)	64 (93%)	5 (7%)	0	100	100
2	B	35/40 (88%)	34 (97%)	1 (3%)	0	100	100
2	D	35/40 (88%)	35 (100%)	0	0	100	100
All	All	210/238 (88%)	201 (96%)	9 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	68/69 (99%)	60 (88%)	8 (12%)	4	2
1	C	66/69 (96%)	59 (89%)	7 (11%)	5	3
2	B	35/33 (106%)	31 (89%)	4 (11%)	4	2
2	D	35/33 (106%)	29 (83%)	6 (17%)	1	1
All	All	204/204 (100%)	179 (88%)	25 (12%)	4	2

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	686	GLU
1	A	707	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	709	VAL
1	A	716	GLU
1	A	727	ARG
1	A	734	ARG
1	A	736	LYS
1	A	757	LYS
2	B	7	LEU
2	B	10	LEU
2	B	15	ARG
2	B	20	LEU
1	C	685	ASP
1	C	702	SER
1	C	704	SER
1	C	705	TYR
1	C	707	ARG
1	C	745	LYS
1	C	754	ARG
2	D	1	MSE
2	D	10	LEU
2	D	20	LEU
2	D	22	LYS
2	D	30	ILE
2	D	35	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	687	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	69/79 (87%)	0.20	1 (1%) 73 74	34, 47, 71, 81	0
1	C	67/79 (84%)	0.19	0 100 100	34, 46, 76, 80	0
2	B	34/40 (85%)	0.04	0 100 100	31, 44, 54, 58	0
2	D	34/40 (85%)	0.08	1 (2%) 54 55	34, 46, 61, 68	0
All	All	204/238 (85%)	0.15	2 (0%) 79 80	31, 45, 72, 81	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	705	TYR	2.4
2	D	35	THR	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.