



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 13, 2024 – 07:26 AM EDT

PDB ID : 3AVM
Title : Crystal structures of novel allosteric peptide inhibitors of HIV integrase in the LEDGF binding site
Authors : Peat, T.S.; Deadman, J.J.; Newman, J.; Rhodes, D.I.
Deposited on : 2011-03-05
Resolution : 1.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1090 (1.88-1.88)
Clashscore	180529	1144 (1.88-1.88)
Ramachandran outliers	177936	1135 (1.88-1.88)
Sidechain outliers	177891	1135 (1.88-1.88)
RSRZ outliers	164620	1090 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	183	 68% 12% 19%
1	B	183	 63% 17% 19%
2	D	8	 88% 12%
2	F	8	 88% 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	B	3	-	-	X	-
5	ACY	A	214	-	-	X	-
5	ACY	A	215	-	-	X	-
5	ACY	A	3	-	-	X	-
5	ACY	B	214	-	-	X	-

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 2807 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	149	1220	783	207	226	4	0	11	0
1	B	149	1250	802	212	231	5	0	14	0

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	30	MET	-	expression tag	UNP P12497
A	31	GLY	-	expression tag	UNP P12497
A	32	SER	-	expression tag	UNP P12497
A	33	SER	-	expression tag	UNP P12497
A	34	HIS	-	expression tag	UNP P12497
A	35	HIS	-	expression tag	UNP P12497
A	36	HIS	-	expression tag	UNP P12497
A	37	HIS	-	expression tag	UNP P12497
A	38	HIS	-	expression tag	UNP P12497
A	39	HIS	-	expression tag	UNP P12497
A	40	SER	-	expression tag	UNP P12497
A	41	SER	-	expression tag	UNP P12497
A	42	GLY	-	expression tag	UNP P12497
A	43	LEU	-	expression tag	UNP P12497
A	44	VAL	-	expression tag	UNP P12497
A	45	PRO	-	expression tag	UNP P12497
A	46	ARG	-	expression tag	UNP P12497
A	47	GLY	-	expression tag	UNP P12497
A	48	SER	-	expression tag	UNP P12497
A	49	HIS	-	expression tag	UNP P12497
A	56	SER	CYS	engineered mutation	UNP P12497
A	139	ASP	PHE	engineered mutation	UNP P12497
A	185	HIS	PHE	engineered mutation	UNP P12497
B	30	MET	-	expression tag	UNP P12497
B	31	GLY	-	expression tag	UNP P12497

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	32	SER	-	expression tag	UNP P12497
B	33	SER	-	expression tag	UNP P12497
B	34	HIS	-	expression tag	UNP P12497
B	35	HIS	-	expression tag	UNP P12497
B	36	HIS	-	expression tag	UNP P12497
B	37	HIS	-	expression tag	UNP P12497
B	38	HIS	-	expression tag	UNP P12497
B	39	HIS	-	expression tag	UNP P12497
B	40	SER	-	expression tag	UNP P12497
B	41	SER	-	expression tag	UNP P12497
B	42	GLY	-	expression tag	UNP P12497
B	43	LEU	-	expression tag	UNP P12497
B	44	VAL	-	expression tag	UNP P12497
B	45	PRO	-	expression tag	UNP P12497
B	46	ARG	-	expression tag	UNP P12497
B	47	GLY	-	expression tag	UNP P12497
B	48	SER	-	expression tag	UNP P12497
B	49	HIS	-	expression tag	UNP P12497
B	56	SER	CYS	engineered mutation	UNP P12497
B	139	ASP	PHE	engineered mutation	UNP P12497
B	185	HIS	PHE	engineered mutation	UNP P12497

- Molecule 2 is a protein called LEDGF peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	8	Total	C	N	O	0	0	0
			66	39	13	14			
2	F	8	Total	C	N	O	0	0	0
			66	39	13	14			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cl 1 1	0	0
4	B	1	Total Cl 1 1	0	0

- Molecule 5 is ACETIC ACID (three-letter code: ACY) (formula: C₂H₄O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	70	Total O 70 70	0	0
6	B	75	Total O 75 75	0	0
6	D	1	Total O 1 1	0	0
6	F	1	Total O 1 1	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Integrase

Chain A: 




- Molecule 1: Integrase

Chain B: 




- Molecule 2: LEDGF peptide

Chain D: 



- Molecule 2: LEDGF peptide

Chain F: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	70.77Å 70.77Å 67.28Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	61.29 – 1.88 61.29 – 1.88	Depositor EDS
% Data completeness (in resolution range)	100.0 (61.29-1.88) 100.0 (61.29-1.88)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 1.88Å)	Xtrriage
Refinement program	REFMAC 5.6.0062	Depositor
R, R_{free}	0.158 , 0.197 0.157 , 0.195	Depositor DCC
R_{free} test set	1547 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	24.8	Xtrriage
Anisotropy	0.047	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 36.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.025 for -h,-k,l 0.480 for h,-h-k,-l 0.025 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2807	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, SO4, ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.29	2/1271 (0.2%)	1.02	3/1724 (0.2%)
1	B	1.30	8/1298 (0.6%)	1.01	4/1758 (0.2%)
2	D	0.79	0/65	0.91	0/85
2	F	0.90	0/65	0.99	0/85
All	All	1.27	10/2699 (0.4%)	1.01	7/3652 (0.2%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	132	TRP	CE3-CZ3	6.23	1.49	1.38
1	B	119	SER	CB-OG	6.22	1.50	1.42
1	B	132	TRP	CE3-CZ3	6.03	1.48	1.38
1	B	107	ARG	CG-CD	5.89	1.66	1.51
1	B	100	PHE	CD1-CE1	5.67	1.50	1.39
1	B	176	VAL	CB-CG2	5.50	1.64	1.52
1	A	107	ARG	CG-CD	5.45	1.65	1.51
1	B	130	CYS	CB-SG	5.27	1.91	1.82
1	B	79	VAL	CB-CG2	5.09	1.63	1.52
1	B	176	VAL	CB-CG1	5.06	1.63	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	116	ASP	CB-CG-OD1	7.23	124.81	118.30
1	B	116	ASP	CB-CG-OD2	-7.12	111.89	118.30
1	A	116	ASP	CB-CG-OD1	6.24	123.91	118.30
1	A	116	ASP	CB-CG-OD2	-5.86	113.03	118.30
1	A	102	LEU	CB-CG-CD2	-5.70	101.31	111.00
1	B	130	CYS	CA-CB-SG	-5.55	104.01	114.00
1	B	173	LYS	CD-CE-NZ	-5.33	99.43	111.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1220	0	1252	31	0
1	B	1250	0	1280	45	0
2	D	66	0	67	0	0
2	F	66	0	67	1	0
3	A	20	0	0	1	0
3	B	20	0	0	3	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	12	0	9	15	0
5	B	4	0	3	2	0
6	A	70	0	0	5	0
6	B	75	0	0	6	0
6	D	1	0	0	0	0
6	F	1	0	0	0	0
All	All	2807	0	2678	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:141[B]:ILE:HD12	1:B:142:PRO:HD2	1.18	1.15
1:A:141[B]:ILE:HD12	1:A:142:PRO:HD2	1.29	1.06
1:B:139:ASP:OD1	1:B:141[B]:ILE:HG22	1.59	1.01
1:B:141[B]:ILE:HD12	1:B:142:PRO:CD	1.89	1.00
5:A:3:ACY:OXT	5:A:215:ACY:H1	1.61	0.99
1:B:141[B]:ILE:HG23	1:B:143:TYR:CE2	1.99	0.97
5:A:3:ACY:O	1:B:87[B]:GLU:OE1	1.84	0.96
1:A:103[B]:LYS:HZ1	1:B:103[B]:LYS:HZ1	1.14	0.95
1:A:141[B]:ILE:HD12	1:A:142:PRO:CD	1.96	0.95

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87[B]:GLU:OE1	5:A:215:ACY:O	1.83	0.95
1:A:103[B]:LYS:HZ1	1:B:103[B]:LYS:NZ	1.67	0.92
1:A:87[B]:GLU:OE2	1:B:103[B]:LYS:HE3	1.71	0.91
5:A:3:ACY:H1	5:A:215:ACY:OXT	1.73	0.87
1:A:103[B]:LYS:NZ	1:B:103[B]:LYS:NZ	2.24	0.86
1:A:103[B]:LYS:NZ	1:B:103[B]:LYS:HZ1	1.74	0.85
1:A:103[A]:LYS:NZ	5:A:3:ACY:H2	1.92	0.84
5:A:215:ACY:H2	1:B:103[A]:LYS:HZ2	1.42	0.84
1:A:103[A]:LYS:HZ3	5:A:3:ACY:H2	1.42	0.84
1:B:141[B]:ILE:CD1	1:B:142:PRO:HD2	2.08	0.83
3:B:5:SO4:O1	6:B:266:HOH:O	1.97	0.83
5:A:215:ACY:H2	1:B:103[A]:LYS:NZ	1.97	0.79
1:B:141[B]:ILE:CG2	1:B:143:TYR:CE2	2.65	0.79
1:B:145:PRO:HA	1:B:148:GLN:HE22	1.46	0.79
1:A:141[B]:ILE:HG23	1:A:143:TYR:CE1	2.19	0.77
5:B:214:ACY:H1	6:B:256:HOH:O	1.86	0.74
1:A:139:ASP:OD1	1:A:141[B]:ILE:HG22	1.88	0.73
1:B:142:PRO:HB2	1:B:148:GLN:HE21	1.58	0.69
1:A:209:GLN:NE2	1:B:202[B]:ASP:OD1	2.23	0.68
1:B:143:TYR:OH	6:B:263:HOH:O	2.08	0.68
1:A:202[B]:ASP:OD1	1:B:209:GLN:NE2	2.26	0.68
1:A:143:TYR:OH	6:A:260:HOH:O	2.11	0.68
1:B:171:HIS:HD2	3:B:3:SO4:O1	1.75	0.68
2:F:2:ARG:O	2:F:2:ARG:HG3	1.94	0.67
1:A:125[B]:THR:CG2	6:A:270:HOH:O	2.42	0.66
1:B:84:ILE:HG21	1:B:154[B]:MET:CE	2.27	0.65
1:B:145:PRO:HA	1:B:148:GLN:NE2	2.12	0.64
1:A:87[B]:GLU:OE2	1:B:103[B]:LYS:CE	2.45	0.63
1:B:139:ASP:OD1	1:B:141[B]:ILE:CG2	2.42	0.62
1:A:141[B]:ILE:CD1	1:A:142:PRO:HD2	2.18	0.61
1:B:84:ILE:HG21	1:B:154[B]:MET:HE1	1.83	0.60
5:B:214:ACY:CH3	6:B:256:HOH:O	2.47	0.60
1:A:141[B]:ILE:HD12	1:A:142:PRO:N	2.17	0.59
1:B:84:ILE:HD13	1:B:154[B]:MET:CE	2.33	0.59
5:A:3:ACY:C	5:A:215:ACY:H1	2.34	0.57
1:B:141[B]:ILE:HD12	1:B:142:PRO:N	2.21	0.55
1:B:152:GLU:CD	1:B:156[B]:LYS:NZ	2.59	0.55
1:A:103[B]:LYS:HE3	1:B:87[B]:GLU:OE2	2.06	0.55
1:A:171:HIS:HD2	3:A:4:SO4:O3	1.88	0.55
1:B:142:PRO:HB2	1:B:148:GLN:NE2	2.21	0.54
5:A:3:ACY:H3	6:B:269:HOH:O	2.07	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:152:GLU:CD	1:B:156[B]:LYS:HZ1	2.13	0.52
1:A:103[A]:LYS:HZ1	5:A:3:ACY:H2	1.76	0.51
1:B:141[B]:ILE:HG23	1:B:143:TYR:HE2	1.62	0.51
1:B:181:PHE:O	1:B:185:HIS:HD2	1.94	0.50
1:A:71:LYS:NZ	6:A:248:HOH:O	2.45	0.49
1:A:162:ILE:O	1:A:166:ARG:HB2	2.13	0.48
1:A:127:LYS:NZ	1:A:137:GLN:HE22	2.12	0.47
1:B:142:PRO:CB	1:B:148:GLN:HE21	2.27	0.47
1:A:197:GLY:HA3	5:A:214:ACY:H3	1.97	0.46
1:A:209:GLN:HE22	1:B:202[B]:ASP:CG	2.16	0.46
1:B:139:ASP:CG	1:B:141[B]:ILE:HG22	2.34	0.46
5:A:214:ACY:H2	6:A:8:HOH:O	2.15	0.46
1:B:84:ILE:HG21	1:B:154[B]:MET:HE2	1.96	0.45
1:A:139:ASP:OD1	1:A:141[B]:ILE:CG2	2.61	0.45
1:B:171:HIS:CD2	3:B:3:SO4:O1	2.65	0.44
1:B:209:GLN:HA	1:B:209:GLN:OE1	2.17	0.44
1:B:136[A]:LYS:NZ	6:B:228:HOH:O	2.50	0.44
1:A:62[A]:GLN:HG2	1:A:151:ILE:HD11	1.99	0.43
1:A:202[B]:ASP:CG	1:B:209:GLN:HE22	2.21	0.43
1:B:151[B]:ILE:O	1:B:154[B]:MET:HB3	2.19	0.43
5:A:215:ACY:H3	6:A:249:HOH:O	2.19	0.42
1:A:139:ASP:CG	1:A:141[B]:ILE:HG22	2.40	0.42
5:A:3:ACY:H1	5:A:215:ACY:C	2.46	0.42
1:B:105:ALA:HB2	1:B:135:ILE:HD11	2.01	0.42
1:B:92[B]:GLU:OE2	1:B:120:ASN:ND2	2.53	0.41
1:B:141[B]:ILE:HG21	1:B:143:TYR:CZ	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	156/183 (85%)	154 (99%)	2 (1%)	0	100	100
1	B	159/183 (87%)	158 (99%)	1 (1%)	0	100	100
2	D	6/8 (75%)	5 (83%)	1 (17%)	0	100	100
2	F	6/8 (75%)	5 (83%)	1 (17%)	0	100	100
All	All	327/382 (86%)	322 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/150 (89%)	130 (97%)	4 (3%)	36	19
1	B	137/150 (91%)	135 (98%)	2 (2%)	60	49
2	D	8/8 (100%)	7 (88%)	1 (12%)	3	0
2	F	8/8 (100%)	8 (100%)	0	100	100
All	All	287/316 (91%)	280 (98%)	7 (2%)	40	28

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	SER
1	A	137	GLN
1	A	188	LYS
1	A	209	GLN
1	B	147	SER
1	B	148	GLN
2	D	1	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	137	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	155	ASN
1	A	171	HIS
1	B	148	GLN
1	B	171	HIS
1	B	185	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 2 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ACY	B	214	-	3,3,3	0.87	0	3,3,3	1.47	0
3	SO4	B	5	-	4,4,4	0.32	0	6,6,6	1.72	1 (16%)
3	SO4	A	2	-	4,4,4	0.13	0	6,6,6	0.98	0
3	SO4	B	3	-	4,4,4	0.29	0	6,6,6	0.76	0
3	SO4	B	8	-	4,4,4	1.01	0	6,6,6	0.48	0
3	SO4	B	1	-	4,4,4	0.22	0	6,6,6	0.96	0
5	ACY	A	215	-	3,3,3	1.15	0	3,3,3	0.25	0
5	ACY	A	214	-	3,3,3	0.82	0	3,3,3	1.50	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	7	-	4,4,4	1.11	1 (25%)	6,6,6	0.59	0
3	SO4	A	6	-	4,4,4	0.26	0	6,6,6	2.05	2 (33%)
5	ACY	A	3	-	3,3,3	1.12	0	3,3,3	0.16	0
3	SO4	A	4	-	4,4,4	0.25	0	6,6,6	0.63	0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	7	SO4	O2-S	2.09	1.57	1.44

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	6	SO4	O4-S-O2	-3.85	89.45	109.56
3	B	5	SO4	O2-S-O1	-2.88	88.74	109.06
3	A	6	SO4	O4-S-O3	2.11	120.20	108.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	214	ACY	2	0
3	B	5	SO4	1	0
3	B	3	SO4	2	0
5	A	215	ACY	8	0
5	A	214	ACY	2	0
5	A	3	ACY	9	0
3	A	4	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	149/183 (81%)	-1.23	0 100 100	11, 22, 42, 61	30 (20%)
1	B	149/183 (81%)	-1.23	0 100 100	10, 21, 42, 61	31 (20%)
2	D	8/8 (100%)	-0.61	0 100 100	33, 40, 66, 69	1 (12%)
2	F	8/8 (100%)	-0.69	0 100 100	33, 39, 68, 73	1 (12%)
All	All	314/382 (82%)	-1.20	0 100 100	10, 22, 47, 73	63 (20%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	8	5/5	0.96	0.09	34,38,46,48	5
5	ACY	A	3	4/4	0.97	0.08	28,29,31,33	0
4	CL	A	213	1/1	0.98	0.05	53,53,53,53	0
4	CL	B	213	1/1	0.98	0.05	52,52,52,52	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	ACY	A	214	4/4	0.98	0.05	36,40,49,52	0
3	SO4	A	7	5/5	0.98	0.06	31,31,44,44	5
5	ACY	A	215	4/4	0.98	0.06	27,30,32,33	0
5	ACY	B	214	4/4	0.98	0.07	30,33,42,44	4
3	SO4	A	6	5/5	0.99	0.04	26,30,36,42	5
3	SO4	A	2	5/5	0.99	0.05	33,38,43,53	5
3	SO4	B	1	5/5	0.99	0.04	32,39,41,53	5
3	SO4	B	3	5/5	0.99	0.04	32,39,44,46	5
3	SO4	B	5	5/5	0.99	0.04	27,31,35,39	5
3	SO4	A	4	5/5	0.99	0.04	27,37,40,47	5

6.5 Other polymers [i](#)

There are no such residues in this entry.