



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2023 – 09:41 PM EDT

PDB ID : 3E6Y  
Title : Structure of 14-3-3 in complex with the differentiation-inducing agent  
Cotylenin A  
Authors : Ottmann, C.; Weyand, M.; Wittinghofer, A.; Oecking, C.  
Deposited on : 2008-08-17  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	 69% 18% 10%
1	B	260	 69% 17% 11%
2	C	5	 40% 20% 40%
2	D	5	 60% 20% 20%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
3	CW1	A	261	X	-	-	-
3	CW1	B	261	X	-	-	-



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			46	34	12		
3	B	1	Total	C	O	0	0
			46	34	12		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	55	Total	O	0	0
			55	55		
5	B	42	Total	O	0	0
			42	42		
5	C	3	Total	O	0	0
			3	3		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 14-3-3-like protein C



- Molecule 1: 14-3-3-like protein C



- Molecule 2: H<sup>+</sup>-ATPase phosphopeptide QSYpTV



- Molecule 2: H<sup>+</sup>-ATPase phosphopeptide QSYpTV



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	41.80Å 141.90Å 51.40Å 90.00° 114.10° 90.00°	Depositor
Resolution (Å)	14.95 – 2.50 14.95 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (14.95-2.50) 98.4 (14.95-2.50)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.62 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.5.0035	Depositor
R, $R_{free}$	0.195 , 0.285 0.209 , 0.295	Depositor DCC
$R_{free}$ test set	930 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.4	Xtrriage
Anisotropy	0.166	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 9.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	0.177 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3926	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CW1, CL, TPO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.86	0/1868	0.97	5/2524 (0.2%)
1	B	0.83	0/1828	0.82	3/2471 (0.1%)
2	C	1.09	0/34	0.73	0/42
2	D	0.87	0/34	0.69	0/42
All	All	0.85	0/3764	0.90	8/5079 (0.2%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	63	ARG	NE-CZ-NH1	12.30	126.45	120.30
1	A	63	ARG	NE-CZ-NH2	-9.47	115.56	120.30
1	A	110	LEU	CA-CB-CG	6.78	130.89	115.30
1	A	38	LEU	CA-CB-CG	6.10	129.34	115.30
1	B	110	LEU	CA-CB-CG	6.02	129.15	115.30
1	B	89	ARG	NE-CZ-NH1	5.40	123.00	120.30
1	B	92	ARG	NE-CZ-NH1	5.36	122.98	120.30
1	A	232	ASP	CB-CG-OD2	-5.05	113.75	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1840	0	1787	26	0
1	B	1801	0	1720	30	0
2	C	46	0	38	4	0
2	D	46	0	38	1	0
3	A	46	0	52	3	0
3	B	46	0	52	5	0
4	A	1	0	0	1	0
5	A	55	0	0	2	0
5	B	42	0	0	3	0
5	C	3	0	0	1	0
All	All	3926	0	3687	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:233:ASN:HD21	2:C:3:TYR:H	1.27	0.79
1:B:128:LEU:HB3	1:B:161:ALA:HB2	1.70	0.74
1:B:173:HIS:HD2	1:B:175:ILE:H	1.33	0.74
1:A:96:GLU:HG2	1:A:141:PHE:CD2	2.30	0.67
1:B:73:GLU:HB3	1:B:88:ILE:HG21	1.77	0.66
1:B:162:GLN:O	1:B:166:THR:HG22	1.96	0.65
1:A:181:LEU:CD1	2:C:4:TPO:HA	2.26	0.65
1:A:69:ILE:HD13	1:B:21:ALA:HB2	1.80	0.63
1:B:52:SER:OG	5:B:275:HOH:O	2.15	0.61
1:B:156:THR:HG23	1:B:157:ALA:N	2.16	0.61
1:B:76:GLU:HG3	1:B:81:ASN:HB2	1.82	0.60
1:A:53:VAL:HG21	3:A:261:CW1:HAA	1.83	0.60
1:B:187:TYR:HA	1:B:191:LEU:HB2	1.84	0.60
1:A:76:GLU:OE1	1:A:81:ASN:ND2	2.28	0.60
1:B:173:HIS:CD2	1:B:175:ILE:H	2.19	0.58
1:B:118:ALA:HB1	1:B:123:SER:HB2	1.86	0.57
1:A:35:SER:O	1:A:38:LEU:HG	2.05	0.57
1:A:199:ASN:HB3	5:A:264:HOH:O	2.07	0.55
1:B:156:THR:CG2	1:B:157:ALA:N	2.71	0.54
3:B:261:CW1:HBP	3:B:261:CW1:CBF	2.38	0.53
1:B:156:THR:CG2	1:B:157:ALA:H	2.22	0.53
1:A:173:HIS:HE1	1:A:175:ILE:HD12	1.74	0.53
1:A:20:GLN:HG2	1:B:72:ILE:HD11	1.91	0.53
1:A:209:ILE:HG22	5:A:276:HOH:O	2.08	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:110:LEU:HD21	1:B:130:MET:HB3	1.92	0.52
1:B:12:VAL:HA	1:B:31:MET:CE	2.41	0.51
1:A:69:ILE:CD1	1:B:21:ALA:HB2	2.40	0.50
1:B:164:ILE:O	1:B:168:GLU:HB2	2.12	0.50
1:B:205:PHE:O	1:B:209:ILE:HG12	2.12	0.49
1:B:153:SER:O	1:B:156:THR:HG22	2.13	0.49
3:B:261:CW1:CAA	5:B:264:HOH:O	2.61	0.48
3:B:261:CW1:HAAB	5:B:264:HOH:O	2.12	0.48
1:A:132:GLY:HA3	1:A:158:TYR:CE1	2.48	0.48
1:A:63:ARG:HG3	4:A:262:CL:CL	2.50	0.48
1:B:73:GLU:HB3	1:B:88:ILE:CG2	2.43	0.48
1:A:173:HIS:CE1	1:A:175:ILE:HD12	2.49	0.48
3:B:261:CW1:HAAA	3:B:261:CW1:HBK	1.63	0.47
1:B:25:GLU:O	1:B:29:GLU:HG3	2.13	0.47
1:A:236:LEU:HD12	1:A:236:LEU:O	2.15	0.46
1:B:120:SER:OG	1:B:122:ASP:HB2	2.15	0.46
1:A:228:GLN:HA	1:A:231:ARG:HD2	1.97	0.46
1:B:142:LYS:O	1:B:147:ARG:HG3	2.15	0.46
1:A:5:PRO:O	1:A:9:GLU:OE1	2.34	0.45
1:A:31:MET:O	1:A:35:SER:HB2	2.16	0.45
2:C:1:GLN:HG2	5:C:27:HOH:O	2.15	0.45
1:B:164:ILE:O	1:B:168:GLU:CB	2.64	0.45
1:A:115:ILE:HB	1:A:116:PRO:HD3	1.98	0.45
3:A:261:CW1:CBF	3:A:261:CW1:HBP	2.47	0.45
3:B:261:CW1:CBF	3:B:261:CW1:CBP	2.95	0.44
1:B:172:THR:O	1:B:177:LEU:HD12	2.18	0.44
1:A:203:GLN:O	1:A:207:GLU:HG3	2.18	0.43
1:A:63:ARG:NH2	2:C:4:TPO:O1P	2.52	0.42
1:B:81:ASN:O	1:B:85:VAL:HG23	2.19	0.42
1:A:21:ALA:HB2	1:B:69:ILE:CD1	2.49	0.42
1:B:63:ARG:HH22	2:D:4:TPO:P	2.43	0.41
3:A:261:CW1:CBF	3:A:261:CW1:CBP	2.98	0.41
1:B:187:TYR:CD1	1:B:191:LEU:HD12	2.55	0.41
1:A:187:TYR:CD1	1:A:191:LEU:HD12	2.55	0.41
1:A:155:LEU:HD13	1:A:186:PHE:HE2	1.86	0.41
1:A:220:TYR:O	1:A:224:THR:HG23	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	232/260 (89%)	223 (96%)	8 (3%)	1 (0%)	34	54
1	B	228/260 (88%)	210 (92%)	17 (8%)	1 (0%)	34	54
2	C	2/5 (40%)	2 (100%)	0	0	100	100
2	D	2/5 (40%)	2 (100%)	0	0	100	100
All	All	464/530 (88%)	437 (94%)	25 (5%)	2 (0%)	34	54

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	6	THR
1	B	39	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/221 (86%)	169 (88%)	22 (12%)	5	11
1	B	183/221 (83%)	166 (91%)	17 (9%)	9	17
2	C	4/4 (100%)	3 (75%)	1 (25%)	0	1
2	D	4/4 (100%)	3 (75%)	1 (25%)	0	1
All	All	382/450 (85%)	341 (89%)	41 (11%)	6	13

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	GLU
1	A	20	GLN
1	A	32	GLU
1	A	35	SER
1	A	36	ASN
1	A	37	SER
1	A	43	LEU
1	A	56	LYS
1	A	110	LEU
1	A	123	SER
1	A	149	GLU
1	A	156	THR
1	A	166	THR
1	A	177	LEU
1	A	181	LEU
1	A	183	PHE
1	A	196	ARG
1	A	209	ILE
1	A	211	GLU
1	A	215	LEU
1	A	218	GLU
1	A	225	LEU
1	B	32	GLU
1	B	52	SER
1	B	70	SER
1	B	76	GLU
1	B	78	SER
1	B	82	GLU
1	B	93	SER
1	B	110	LEU
1	B	122	ASP
1	B	123	SER
1	B	149	GLU
1	B	166	THR
1	B	168	GLU
1	B	183	PHE
1	B	219	SER
1	B	225	LEU
1	B	232	ASP
2	C	1	GLN
2	D	1	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	ASN
1	A	199	ASN
1	A	233	ASN
1	B	49	ASN
1	B	74	GLN
1	B	173	HIS
1	B	233	ASN
2	D	1	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	TPO	D	4	2	8,10,11	0.63	0	10,14,16	1.62	2 (20%)
2	TPO	C	4	2	8,10,11	0.90	0	10,14,16	1.23	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TPO	D	4	2	-	3/9/11/13	-
2	TPO	C	4	2	-	2/9/11/13	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	4	TPO	CG2-CB-CA	-2.85	107.54	113.16
2	D	4	TPO	OG1-P-O1P	-2.61	99.32	109.39
2	C	4	TPO	O-C-CA	-2.23	118.93	124.78

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	4	TPO	CB-OG1-P-O1P
2	C	4	TPO	CB-OG1-P-O2P
2	D	4	TPO	CB-OG1-P-O2P
2	C	4	TPO	O-C-CA-CB
2	D	4	TPO	O-C-CA-CB

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	4	TPO	1	0
2	C	4	TPO	2	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	CW1	B	261	-	48,51,51	2.70	13 (27%)	44,81,81	2.40	14 (31%)
3	CW1	A	261	-	48,51,51	2.67	11 (22%)	44,81,81	3.46	23 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CW1	B	261	-	1/1/17/19	7/21/119/119	1/7/6/6
3	CW1	A	261	-	1/1/17/19	7/21/119/119	1/7/6/6

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	261	CW1	CAM-CBE	11.33	1.42	1.33
3	A	261	CW1	CAM-CBE	10.39	1.41	1.33
3	A	261	CW1	CBR-CBM	8.79	1.67	1.54
3	B	261	CW1	CBR-CBM	6.42	1.63	1.54
3	B	261	CW1	CBD-CBF	5.77	1.41	1.33
3	A	261	CW1	CBD-CBF	5.76	1.41	1.33
3	B	261	CW1	OBC-CBR	5.51	1.55	1.45
3	A	261	CW1	OBB-CBM	-4.33	1.34	1.41
3	B	261	CW1	CBG-CBD	3.75	1.55	1.51
3	A	261	CW1	CBG-CBD	3.72	1.55	1.51
3	A	261	CW1	CBT-CBN	-3.47	1.49	1.54
3	A	261	CW1	OBC-CBR	3.42	1.52	1.45
3	B	261	CW1	CBT-CBN	-3.37	1.49	1.54
3	B	261	CW1	CAU-CBS	-3.24	1.48	1.52
3	A	261	CW1	CAS-CBS	-3.05	1.49	1.53
3	B	261	CW1	CBQ-CBF	2.80	1.55	1.50
3	B	261	CW1	OAY-CBK	-2.78	1.40	1.44
3	B	261	CW1	CBP-CBH	2.77	1.58	1.56
3	A	261	CW1	OAZ-CBM	2.67	1.46	1.41
3	B	261	CW1	OAY-CBL	2.57	1.48	1.41
3	A	261	CW1	CBT-CBK	2.49	1.58	1.54
3	A	261	CW1	CBP-CBE	2.35	1.56	1.52
3	B	261	CW1	CBP-CBE	2.31	1.56	1.52
3	B	261	CW1	OBB-CBT	-2.26	1.39	1.43

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	261	CW1	OAZ-CBM-OBB	-12.09	90.97	105.85
3	A	261	CW1	CBG-CBD-CBF	-6.74	120.72	129.59
3	B	261	CW1	OAZ-CBM-OBB	-6.58	97.75	105.85
3	A	261	CW1	CBT-CBN-CBI	-6.53	106.20	114.33

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	261	CW1	OAW-CAU-CBS	-5.91	103.97	109.88
3	B	261	CW1	CAS-CBS-CBE	5.33	108.87	103.65
3	A	261	CW1	CBP-CBH-CBJ	-5.08	107.04	112.39
3	A	261	CW1	CAO-CBD-CBG	5.06	127.22	120.06
3	B	261	CW1	CBT-CBN-CBI	-4.85	108.29	114.33
3	B	261	CW1	OBC-CBR-CAG	4.80	117.86	108.61
3	A	261	CW1	OAZ-CBN-CBT	-4.79	96.94	103.70
3	B	261	CW1	CBS-CBE-CBP	-4.69	105.33	108.97
3	A	261	CW1	OBB-CBM-CBR	-4.57	100.87	104.57
3	B	261	CW1	CAB-OAW-CAU	-4.56	102.08	112.16
3	A	261	CW1	OBC-CBR-CAG	3.85	116.04	108.61
3	A	261	CW1	CAG-CBR-CAQ	3.69	117.00	111.67
3	A	261	CW1	CBS-CBE-CBP	-3.69	106.11	108.97
3	A	261	CW1	CAR-CAS-CBS	3.64	107.98	104.03
3	A	261	CW1	CAD-CBG-CBD	3.63	116.87	111.35
3	A	261	CW1	OBA-CBL-OAY	-3.47	100.97	110.67
3	A	261	CW1	CAS-CBS-CBE	3.35	106.93	103.65
3	B	261	CW1	CAE-CBH-CBP	3.32	117.56	114.85
3	B	261	CW1	CAG-CBR-CAQ	3.16	116.22	111.67
3	B	261	CW1	OAZ-CBN-CBT	-3.02	99.43	103.70
3	A	261	CW1	CAR-CBP-CBE	2.95	107.05	103.02
3	A	261	CW1	CAN-CAQ-CBR	-2.88	108.03	116.27
3	A	261	CW1	CAE-CBH-CBP	2.78	117.12	114.85
3	B	261	CW1	CBP-CBH-CBJ	-2.78	109.47	112.39
3	B	261	CW1	CAN-CAQ-CBR	-2.77	108.33	116.27
3	A	261	CW1	CAC-CBG-CBD	-2.76	107.15	111.35
3	B	261	CW1	OAW-CAU-CBS	2.59	112.47	109.88
3	A	261	CW1	CAA-OAV-CAT	-2.42	102.27	112.42
3	B	261	CW1	CBG-CBD-CBF	-2.32	126.54	129.59
3	A	261	CW1	CAE-CBH-CBJ	2.31	115.29	111.35
3	B	261	CW1	CAR-CBP-CBE	2.06	105.83	103.02
3	A	261	CW1	CAR-CBP-CBH	-2.05	106.10	110.81
3	A	261	CW1	CAU-CBS-CBE	2.02	116.60	112.21

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	261	CW1	CBR
3	B	261	CW1	CBR

All (14) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	261	CW1	OAX-CAN-CAQ-CBR
3	A	261	CW1	CAN-CAQ-CBR-CAG
3	A	261	CW1	OAV-CAT-CBK-OAY
3	A	261	CW1	OAW-CAU-CBS-CBE
3	A	261	CW1	OAW-CAU-CBS-OAK
3	A	261	CW1	OAW-CAU-CBS-CAS
3	B	261	CW1	CAN-CAQ-CBR-CAG
3	B	261	CW1	OAV-CAT-CBK-OAY
3	B	261	CW1	OAW-CAU-CBS-CBE
3	B	261	CW1	OAW-CAU-CBS-OAK
3	B	261	CW1	OAW-CAU-CBS-CAS
3	A	261	CW1	OAV-CAT-CBK-CBT
3	B	261	CW1	OAH-CAL-OAX-CAN
3	B	261	CW1	CBK-CAT-OAV-CAA


All (2) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	261	CW1	CBI-CBK-CBL-CBN-CBT-OAY
3	B	261	CW1	CBI-CBK-CBL-CBN-CBT-OAY

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	261	CW1	5	0
3	A	261	CW1	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	234/260 (90%)	-0.47	0 <b>100</b> <b>100</b>	13, 21, 33, 39	0
1	B	232/260 (89%)	-0.28	3 (1%) 77 79	12, 21, 36, 46	0
2	C	4/5 (80%)	0.24	0 <b>100</b> <b>100</b>	15, 16, 17, 19	0
2	D	4/5 (80%)	0.46	0 <b>100</b> <b>100</b>	15, 19, 20, 24	0
All	All	474/530 (89%)	-0.36	3 (0%) 89 90	12, 21, 34, 46	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	220	TYR	2.4
1	B	240	ASP	2.3
1	B	145	ALA	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	TPO	C	4	11/12	0.98	0.12	8,11,14,16	0
2	TPO	D	4	11/12	0.98	0.12	14,16,17,17	0

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	CL	A	262	1/1	0.83	0.25	43,43,43,43	0
3	CW1	B	261	46/46	0.92	0.14	15,20,34,42	0
3	CW1	A	261	46/46	0.95	0.11	10,17,26,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.