



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 26, 2023 – 10:39 PM EDT

PDB ID : 3GP8
Title : Crystal structure of the binary complex of RecD2 with DNA
Authors : Saikrishnan, K.; Cook, N.; Wigley, D.B.
Deposited on : 2009-03-23
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	574	 7% 76% 18%
2	X	14	 7% 21% 21% 14% 43%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4435 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Exodeoxyribonuclease V, subunit RecD, putative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	551	4088	2565	763	750	10	0	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	150	MET	-	expression tag	UNP Q9RT63
A	716	LEU	-	expression tag	UNP Q9RT63
A	717	GLU	-	expression tag	UNP Q9RT63
A	718	HIS	-	expression tag	UNP Q9RT63
A	719	HIS	-	expression tag	UNP Q9RT63
A	720	HIS	-	expression tag	UNP Q9RT63
A	721	HIS	-	expression tag	UNP Q9RT63
A	722	HIS	-	expression tag	UNP Q9RT63
A	723	HIS	-	expression tag	UNP Q9RT63

- Molecule 2 is a DNA chain called 5'-D(*TP*TP*TP*TP*TP*T*TP*TP*TP*TP*TP*TP*TP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	X	8	157	80	16	54	7	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	186	Total	O	0	0
			186	186		
3	X	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Exodeoxyribonuclease V, subunit RecD, putative



- Molecule 2: 5'-D(*TP*TP*TP*TP*TP*T*TP*TP*TP*TP*TP*TP*TP*TP*T)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.14Å 90.32Å 142.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.18 – 2.50 38.88 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.0 (45.18-2.50) 99.0 (38.88-2.50)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	0.03	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.22 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.228 , 0.268 0.230 , 0.268	Depositor DCC
R_{free} test set	1256 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	43.2	Xtrriage
Anisotropy	0.499	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 53.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4435	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/4157	0.60	0/5646
2	X	0.95	0/172	1.74	4/264 (1.5%)
All	All	0.45	0/4329	0.69	4/5910 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	X	2	DT	C5-C4-O4	-6.01	120.69	124.90
2	X	2	DT	N3-C4-O4	5.52	123.21	119.90
2	X	5	DT	C4-C5-C7	5.52	122.31	119.00
2	X	7	DT	O4'-C1'-N1	-5.33	104.27	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4088	0	4121	89	0
2	X	157	0	98	5	0
3	A	186	0	0	1	0
3	X	4	0	0	0	0
All	All	4435	0	4219	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 11.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:435:ASP:HA	1:A:461:VAL:HG13	1.46	0.95
1:A:591:VAL:CG1	1:A:637:LEU:HD23	1.97	0.94
1:A:591:VAL:HG11	1:A:637:LEU:HD23	1.50	0.91
1:A:656:VAL:CG1	1:A:680:ALA:HB2	2.02	0.88
1:A:591:VAL:HG11	1:A:637:LEU:CD2	2.07	0.84
1:A:357:VAL:HB	1:A:485:THR:HG22	1.60	0.83
1:A:665:MET:HE1	1:A:668:LEU:HD12	1.58	0.83
1:A:560:MET:CE	1:A:644:THR:HG23	2.08	0.82
1:A:560:MET:HE3	1:A:644:THR:HG23	1.63	0.78
1:A:560:MET:HE2	1:A:644:THR:N	1.98	0.77
1:A:480:ALA:HA	1:A:485:THR:HG21	1.67	0.76
1:A:592:VAL:HG11	1:A:643:LEU:HD21	1.69	0.73
1:A:180:PHE:CZ	1:A:194:THR:CG2	2.79	0.66
1:A:483:ALA:O	1:A:485:THR:HG23	1.95	0.66
1:A:560:MET:CE	1:A:644:THR:H	2.10	0.65
1:A:197:GLU:OE2	1:A:275:ARG:HD2	1.98	0.64
1:A:665:MET:CE	1:A:668:LEU:HD12	2.28	0.62
1:A:560:MET:CE	1:A:644:THR:N	2.63	0.61
1:A:592:VAL:HG12	1:A:607:LEU:HD23	1.83	0.61
1:A:160:LEU:C	1:A:160:LEU:HD23	2.22	0.60
1:A:434:VAL:HG21	1:A:448:LEU:HD21	1.84	0.60
1:A:237:GLN:HB2	2:X:2:DT:O4	2.02	0.60
1:A:578:VAL:HG23	1:A:613:ALA:HB1	1.85	0.59
1:A:357:VAL:HB	1:A:485:THR:CG2	2.32	0.59
1:A:560:MET:HE2	1:A:644:THR:HG23	1.82	0.58
1:A:592:VAL:HG13	1:A:640:GLY:HA3	1.86	0.58
1:A:564:ASN:HD21	1:A:642:ALA:H	1.52	0.58
1:A:198:VAL:HG21	1:A:201:ILE:HD12	1.86	0.57
1:A:313:ARG:HG3	1:A:715:HIS:HB3	1.86	0.56
1:A:672:LEU:C	1:A:672:LEU:HD13	2.27	0.54
1:A:390:THR:HG22	2:X:5:DT:H1'	1.90	0.54
1:A:188:LEU:HD22	1:A:195:LEU:HG	1.90	0.53
1:A:191:ASP:O	1:A:194:THR:HB	2.08	0.53
1:A:610:VAL:HA	1:A:620:VAL:HG22	1.91	0.53
1:A:384:VAL:HG12	1:A:386:LEU:HD12	1.91	0.53
1:A:538:VAL:HG13	1:A:547:VAL:CG1	2.38	0.53
1:A:312:ILE:HD13	1:A:479:LEU:HD21	1.90	0.52
1:A:398:GLY:HA2	1:A:401:THR:HG22	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:520:LEU:HD23	1:A:520:LEU:O	2.10	0.52
1:A:656:VAL:HG13	1:A:680:ALA:HB2	1.88	0.52
1:A:412:LEU:HD11	1:A:448:LEU:HA	1.93	0.51
1:A:560:MET:HE3	1:A:647:ARG:HH11	1.76	0.51
1:A:591:VAL:HG13	1:A:638:GLN:O	2.12	0.50
2:X:3:DT:H72	2:X:4:DT:O4	2.11	0.50
1:A:563:LEU:O	1:A:567:LEU:HD22	2.12	0.49
1:A:591:VAL:CG1	1:A:592:VAL:N	2.75	0.49
1:A:198:VAL:CG2	1:A:201:ILE:HD12	2.42	0.49
1:A:384:VAL:HG12	1:A:386:LEU:CD1	2.42	0.49
1:A:180:PHE:HZ	1:A:194:THR:CG2	2.23	0.49
1:A:386:LEU:HB3	1:A:397:LEU:HD13	1.93	0.49
1:A:665:MET:HE1	1:A:695:ILE:HG21	1.94	0.49
1:A:592:VAL:HG13	1:A:640:GLY:CA	2.43	0.48
1:A:656:VAL:HG11	1:A:680:ALA:HB2	1.92	0.48
1:A:578:VAL:CG2	1:A:613:ALA:HB1	2.44	0.48
1:A:603:PHE:CE2	1:A:606:THR:HG23	2.49	0.48
1:A:376:LEU:HD22	1:A:380:LEU:HD11	1.96	0.48
1:A:259:THR:H	1:A:262:GLN:NE2	2.12	0.48
1:A:610:VAL:HG22	1:A:620:VAL:CG2	2.45	0.47
1:A:647:ARG:HG2	3:A:763:HOH:O	2.15	0.46
1:A:665:MET:CE	1:A:695:ILE:HG21	2.45	0.46
1:A:470:VAL:HB	2:X:5:DT:H73	1.98	0.45
1:A:259:THR:H	1:A:262:GLN:HE21	1.65	0.45
1:A:246:SER:CB	1:A:293:GLU:OE1	2.65	0.45
1:A:610:VAL:HG22	1:A:620:VAL:HG22	1.97	0.45
1:A:613:ALA:HB2	1:A:618:LEU:HD23	1.97	0.45
1:A:620:VAL:HG12	1:A:621:ASP:N	2.32	0.45
1:A:591:VAL:HG12	1:A:637:LEU:HD23	1.89	0.45
1:A:614:GLU:O	1:A:615:GLY:C	2.55	0.45
1:A:401:THR:HG23	1:A:403:ARG:HB2	1.99	0.44
1:A:551:THR:O	1:A:644:THR:HA	2.18	0.44
1:A:180:PHE:CZ	1:A:194:THR:HG22	2.53	0.44
1:A:246:SER:HB3	1:A:293:GLU:OE1	2.18	0.44
1:A:470:VAL:HG11	2:X:4:DT:H1'	1.99	0.43
1:A:609:MET:O	1:A:620:VAL:HG13	2.18	0.43
1:A:486:ILE:N	1:A:486:ILE:HD12	2.33	0.43
1:A:517:ASP:OD1	1:A:518:LYS:N	2.51	0.43
1:A:535:ALA:HB1	1:A:570:LEU:HD12	2.00	0.43
1:A:245:ARG:HH21	1:A:268:GLU:HG3	1.84	0.42
1:A:591:VAL:HG11	1:A:637:LEU:HD22	1.97	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:591:VAL:HG12	1:A:592:VAL:N	2.35	0.42
1:A:627:VAL:HG22	1:A:628:GLU:H	1.84	0.42
1:A:398:GLY:CA	1:A:401:THR:HG22	2.49	0.42
1:A:331:VAL:O	1:A:331:VAL:HG23	2.19	0.42
1:A:376:LEU:O	1:A:379:SER:OG	2.29	0.41
1:A:395:ARG:CD	1:A:395:ARG:C	2.89	0.41
1:A:180:PHE:CZ	1:A:194:THR:HG23	2.55	0.41
1:A:401:THR:HG23	1:A:403:ARG:N	2.36	0.41
1:A:660:LEU:O	1:A:688:GLY:HA3	2.21	0.41
1:A:193:PHE:CZ	1:A:222:ARG:HG3	2.55	0.40
1:A:386:LEU:HD22	1:A:401:THR:HG21	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	541/574 (94%)	519 (96%)	21 (4%)	1 (0%)	47 68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	615	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	403/435 (93%)	366 (91%)	37 (9%)	9 18

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	158	ARG
1	A	159	LEU
1	A	185	LEU
1	A	192	LEU
1	A	198	VAL
1	A	204	LEU
1	A	213	ARG
1	A	223	LEU
1	A	236	THR
1	A	237	GLN
1	A	247	ARG
1	A	302	LEU
1	A	308	LEU
1	A	313	ARG
1	A	322	ASP
1	A	341	GLU
1	A	355	LEU
1	A	376	LEU
1	A	395	ARG
1	A	396	ARG
1	A	397	LEU
1	A	444	LEU
1	A	459	LEU
1	A	477	LEU
1	A	519	ARG
1	A	523	THR
1	A	567	LEU
1	A	570	LEU
1	A	579	ARG
1	A	612	LYS
1	A	627	VAL
1	A	637	LEU
1	A	638	GLN
1	A	656	VAL
1	A	662	GLU
1	A	701	ARG
1	A	709	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	237	GLN
1	A	262	GLN
1	A	343	GLN
1	A	466	GLN
1	A	502	GLN
1	A	564	ASN
1	A	593	GLN
1	A	638	GLN
1	A	649	GLN
1	A	664	HIS
1	A	671	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	551/574 (95%)	0.66	40 (7%) 15 15	38, 46, 52, 59	1 (0%)
2	X	8/14 (57%)	0.09	1 (12%) 3 3	40, 59, 72, 86	0
All	All	559/588 (95%)	0.65	41 (7%) 15 15	38, 46, 53, 86	1 (0%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	632	ALA	8.1
1	A	613	ALA	7.2
1	A	579	ARG	6.2
1	A	635	PHE	6.2
1	A	618	LEU	5.6
1	A	615	GLY	5.2
1	A	585	ALA	5.1
1	A	614	GLU	5.1
1	A	578	VAL	4.8
1	A	631	GLY	4.5
1	A	577	GLY	4.5
1	A	289	ALA	4.3
1	A	576	GLY	4.2
1	A	338	GLY	4.0
1	A	583	GLY	3.8
1	A	617	ARG	3.4
1	A	581	ALA	3.4
1	A	634	LEU	3.4
1	A	625	ASN	3.2
1	A	290	ALA	3.1
1	A	627	VAL	2.9
2	X	8	DT	2.7
1	A	630	THR	2.7
1	A	629	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	520	LEU	2.6
1	A	619	THR	2.5
1	A	607	LEU	2.5
1	A	288	ALA	2.5
1	A	153	GLN	2.4
1	A	616	ALA	2.4
1	A	715	HIS	2.3
1	A	541	LEU	2.3
1	A	331	VAL	2.2
1	A	417	GLN	2.2
1	A	284	PHE	2.1
1	A	467	LEU	2.1
1	A	628	GLU	2.1
1	A	423	HIS	2.1
1	A	519	ARG	2.1
1	A	291	THR	2.0
1	A	603	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.