



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 14, 2024 – 04:19 PM EST

PDB ID : 3HSY
Title : High resolution structure of a dimeric GluR2 N-terminal domain (NTD)
Authors : Rossmann, M.; Sukumaran, M.; Penn, A.C.; Veprintsev, D.B.; Greger, I.H.
Deposited on : 2009-06-11
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.21
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2888 (1.76-1.76)
Clashscore	180529	3097 (1.76-1.76)
Ramachandran outliers	177936	3072 (1.76-1.76)
Sidechain outliers	177891	3072 (1.76-1.76)
RSRZ outliers	164620	2887 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	376	<div style="display: flex; align-items: center;"> <div style="width: 9%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 82%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: grey;"></div> </div>
1	B	376	<div style="display: flex; align-items: center;"> <div style="width: 8%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 87%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: yellow;"></div> </div>
2	C	3	<div style="display: flex; align-items: center;"> <div style="width: 33%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 67%; height: 10px; background-color: yellow;"></div> </div>

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 6601 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	354	2851	1816	491	535	9	0	7	0
1	B	376	3027	1930	514	573	10	0	7	0

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	3	39	22	2	15	0	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
3	A	1	14	8	1	5	0	0
3	B	1	14	8	1	5	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
			Total	O			S
4	A	1	5	4	1	0	0
4	B	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	282	Total	O	0	0
			282	282		
5	B	354	Total	O	0	0
			354	354		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamate receptor 2



- Molecule 1: Glutamate receptor 2



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	82.45Å 93.69Å 101.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.53 – 1.75 42.53 – 1.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (42.53-1.75) 100.0 (42.53-1.75)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.51 (at 1.75Å)	Xtrriage
Refinement program	PHENIX 1.6.1_357	Depositor
R, R_{free}	0.178 , 0.205 0.182 , 0.208	Depositor DCC
R_{free} test set	1589 reflections (1.99%)	wwPDB-VP
Wilson B-factor (Å ²)	21.0	Xtrriage
Anisotropy	0.028	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6601	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, SO4, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/2923	0.50	0/3956
1	B	0.35	0/3103	0.51	0/4200
All	All	0.34	0/6026	0.50	0/8156

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2851	0	2808	39	0
1	B	3027	0	2971	45	0
2	C	39	0	34	0	0
3	A	14	0	13	5	0
3	B	14	0	13	0	0
4	A	5	0	0	0	0
4	B	15	0	0	0	0
5	A	282	0	0	10	0
5	B	354	0	0	6	0
All	All	6601	0	5839	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 7.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:235:ASN:ND2	3:A:401:NAG:O5	1.87	1.08
1:A:235:ASN:ND2	3:A:401:NAG:C1	2.17	1.08
1:B:229:ILE:HD12	1:B:230:GLN:N	1.91	0.86
1:A:358:LYS:HG3	1:A:363:ARG:HD3	1.60	0.82
1:A:54[B]:ASN:OD1	1:B:310:LEU:HD22	1.82	0.80
1:B:329:GLN:HG3	5:B:573:HOH:O	1.84	0.78
1:A:235:ASN:HD22	3:A:401:NAG:C1	1.97	0.75
1:A:23:PHE:CZ	1:A:27:MET:HE2	2.21	0.75
1:B:227:LEU:HD12	1:B:230:GLN:NE2	2.03	0.73
1:B:100:THR:HG23	5:B:540:HOH:O	1.89	0.72
1:B:242:VAL:HG11	1:B:272[B]:ILE:HD11	1.72	0.71
1:A:224:GLY:HA3	5:A:509:HOH:O	1.91	0.69
1:B:344:ASN:HB3	5:B:540:HOH:O	1.95	0.66
1:B:101[A]:HIS:CE1	1:B:346:LYS:HE3	2.31	0.66
1:B:229:ILE:HD12	1:B:230:GLN:HG3	1.77	0.65
1:A:65:TYR:O	5:A:501:HOH:O	2.13	0.64
1:A:8:ILE:HD11	1:A:27:MET:HE1	1.79	0.63
1:B:148:ALA:HB3	5:B:511:HOH:O	1.97	0.62
1:A:23:PHE:CZ	1:A:27:MET:CE	2.83	0.62
1:B:101[A]:HIS:N	1:B:101[A]:HIS:ND1	2.47	0.62
1:A:147:ALA:HA	1:A:152:TRP:HB2	1.82	0.60
1:A:44:LEU:N	1:A:44:LEU:HD23	2.16	0.60
1:B:44:LEU:H	1:B:44:LEU:HD23	1.65	0.60
1:B:229:ILE:CD1	1:B:230:GLN:HG3	2.30	0.60
1:A:80[A]:THR:HG21	5:A:638:HOH:O	2.03	0.59
1:A:346:LYS:HE3	5:A:746:HOH:O	2.02	0.58
1:A:370:GLU:HB2	5:A:523:HOH:O	2.02	0.58
1:B:14:ARG:NH2	1:B:45:GLU:HG3	2.18	0.58
1:B:62:ARG:HG2	1:B:302:ARG:O	2.04	0.58
1:A:105:ILE:HD12	1:A:105:ILE:N	2.17	0.58
1:B:101[A]:HIS:HE1	1:B:346:LYS:HE3	1.67	0.57
1:B:300:ILE:N	1:B:300:ILE:HD12	2.19	0.57
1:A:100:THR:HG22	1:A:102:PRO:HD3	1.87	0.56
1:A:213:HIS:ND1	3:A:401:NAG:H4	2.20	0.56
1:B:83[B]:CYS:SG	1:B:90:PHE:HB2	2.46	0.55
1:A:235:ASN:CG	3:A:401:NAG:C1	2.75	0.55
1:A:184:ARG:HB2	1:A:212:TYR:HA	1.89	0.55
1:B:13:PRO:HG3	1:B:46:VAL:HG11	1.89	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:370:GLU:O	1:A:373:LYS:NZ	2.23	0.54
1:B:229:ILE:HD12	1:B:230:GLN:H	1.70	0.54
1:B:27:MET:HE1	1:B:37:LEU:O	2.08	0.53
1:A:70:PHE:CE1	1:A:93:PRO:HG2	2.44	0.53
1:B:272[B]:ILE:HD13	1:B:277:ALA:HB2	1.91	0.52
1:B:13:PRO:HD3	1:B:70:PHE:CD1	2.44	0.52
1:B:273:LYS:HD3	5:B:780:HOH:O	2.08	0.52
1:A:8:ILE:HD11	1:A:27:MET:CE	2.40	0.51
1:A:302:ARG:HD2	5:A:690:HOH:O	2.11	0.50
1:A:44:LEU:HD23	1:A:44:LEU:H	1.76	0.50
1:B:70:PHE:CE2	1:B:93:PRO:HG2	2.47	0.49
1:B:6:ILE:HG23	1:B:65:TYR:CD2	2.47	0.49
1:B:203:ILE:HD12	1:B:209:VAL:HG21	1.95	0.49
1:A:194:LYS:HD2	5:A:611:HOH:O	2.13	0.48
1:A:240[B]:GLN:NE2	5:A:503:HOH:O	2.46	0.48
1:B:27:MET:HE1	1:B:39:PRO:HD3	1.96	0.47
1:B:44:LEU:HD23	1:B:44:LEU:N	2.29	0.47
1:B:11:LEU:CD2	1:B:42:ASP:HB2	2.44	0.47
1:A:83:CYS:SG	1:A:90:PHE:HB2	2.55	0.46
1:B:97:THR:OG1	1:B:101[A]:HIS:CD2	2.69	0.46
1:B:176:GLN:HG3	5:B:591:HOH:O	2.16	0.46
1:A:76:VAL:O	1:A:80[A]:THR:HG23	2.16	0.45
1:A:287:THR:O	1:A:291:ARG:HG3	2.17	0.44
1:B:210:LYS:HG2	1:B:232:GLY:O	2.17	0.44
1:B:252:PHE:HE2	1:B:272[B]:ILE:HG21	1.81	0.44
1:B:101[A]:HIS:CE1	1:B:346:LYS:CE	3.00	0.43
1:A:203:ILE:HD11	1:A:230:GLN:O	2.18	0.43
1:A:371:VAL:HG13	1:A:372:ASP:OD1	2.19	0.43
1:A:250:SER:O	1:A:254:GLU:HG3	2.18	0.43
1:B:151:LYS:HD2	1:B:151:LYS:N	2.33	0.43
1:A:65:TYR:HD2	1:A:323:ILE:HD11	1.84	0.43
1:B:200:ASP:O	1:B:204:THR:HG23	2.19	0.42
1:B:44:LEU:N	1:B:44:LEU:CD2	2.82	0.42
1:B:300:ILE:HD12	1:B:300:ILE:H	1.83	0.42
1:A:265:PRO:HG2	5:A:673:HOH:O	2.18	0.42
1:A:44:LEU:N	1:A:44:LEU:CD2	2.82	0.41
1:B:4:ASN:HD22	1:B:4:ASN:N	2.18	0.41
1:B:12:PHE:HA	1:B:13:PRO:HD3	1.88	0.41
1:A:59:GLN:HB3	1:A:64:VAL:HG21	2.03	0.41
1:B:97:THR:HG1	1:B:101[A]:HIS:CD2	2.38	0.41
1:A:135[B]:ARG:HE	1:A:135[B]:ARG:HB2	1.46	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:203:ILE:CD1	1:B:209:VAL:HB	2.50	0.41
1:A:359:THR:HG23	5:A:715:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	357/376 (95%)	346 (97%)	10 (3%)	1 (0%)	37	22
1	B	381/376 (101%)	373 (98%)	8 (2%)	0	100	100
All	All	738/752 (98%)	719 (97%)	18 (2%)	1 (0%)	48	32

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	224	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	309/325 (95%)	307 (99%)	2 (1%)	84	78
1	B	328/325 (101%)	322 (98%)	6 (2%)	54	37
All	All	637/650 (98%)	629 (99%)	8 (1%)	70	51

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	184	ARG
1	A	372	ASP
1	B	4	ASN
1	B	43	ASN
1	B	101[A]	HIS
1	B	101[B]	HIS
1	B	106	GLN
1	B	110	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	235	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	2,1	14,14,15	1.01	1 (7%)	17,19,21	2.67	7 (41%)
2	NAG	C	2	2	14,14,15	0.83	0	17,19,21	0.95	0
2	BMA	C	3	2	11,11,12	0.84	0	15,15,17	1.39	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	1/6/23/26	0/1/1/1
2	NAG	C	2	2	-	1/6/23/26	0/1/1/1
2	BMA	C	3	2	-	2/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	NAG	C1-C2	2.03	1.55	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	NAG	C1-O5-C5	8.01	122.92	112.19
2	C	1	NAG	C3-C4-C5	3.81	117.14	110.23
2	C	1	NAG	O5-C5-C6	2.80	113.11	107.66
2	C	1	NAG	O4-C4-C5	-2.75	102.54	109.32
2	C	3	BMA	C1-C2-C3	2.56	113.38	109.64
2	C	3	BMA	C1-O5-C5	2.52	115.56	112.19
2	C	1	NAG	C8-C7-N2	2.45	120.17	116.12
2	C	1	NAG	O5-C5-C4	2.29	116.41	110.83
2	C	3	BMA	O4-C4-C5	2.16	114.65	109.32
2	C	1	NAG	O6-C6-C5	-2.13	104.09	111.33

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	3	BMA	O5-C5-C6-O6
2	C	3	BMA	C4-C5-C6-O6
2	C	1	NAG	C4-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	B	404	-	4,4,4	0.25	0	6,6,6	0.11	0
3	NAG	A	401	-	14,14,15	1.83	3 (21%)	17,19,21	2.78	4 (23%)
3	NAG	B	401	1	14,14,15	0.60	0	17,19,21	1.70	4 (23%)
4	SO4	B	403	-	4,4,4	0.28	0	6,6,6	0.25	0
4	SO4	B	402	-	4,4,4	0.24	0	6,6,6	0.19	0
4	SO4	A	402	-	4,4,4	0.24	0	6,6,6	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	401	-	-	3/6/23/26	0/1/1/1
3	NAG	B	401	1	-	0/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	401	NAG	C1-C2	4.45	1.58	1.52
3	A	401	NAG	O5-C1	-3.95	1.37	1.43
3	A	401	NAG	C4-C5	2.02	1.57	1.53

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	NAG	C4-C3-C2	-6.52	101.47	111.02
3	A	401	NAG	C1-O5-C5	-6.39	103.62	112.19
3	A	401	NAG	O4-C4-C5	5.27	122.29	109.32
3	A	401	NAG	O5-C5-C6	2.90	113.32	107.66
3	B	401	NAG	O3-C3-C2	-2.50	104.22	109.40
3	B	401	NAG	O4-C4-C5	2.47	115.41	109.32
3	B	401	NAG	C1-C2-N2	-2.46	106.56	110.43
3	B	401	NAG	C3-C4-C5	2.42	114.62	110.23

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	401	NAG	O5-C5-C6-O6
3	A	401	NAG	C1-C2-N2-C7
3	A	401	NAG	C3-C2-N2-C7

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	NAG	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	354/376 (94%)	0.34	35 (9%)	14 16	9, 23, 59, 147	7 (1%)
1	B	376/376 (100%)	0.10	30 (7%)	20 22	7, 19, 63, 110	7 (1%)
All	All	730/752 (97%)	0.21	65 (8%)	17 19	7, 21, 61, 147	14 (1%)

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	162	ILE	6.6
1	A	371	VAL	6.2
1	B	300	ILE	5.3
1	B	229	ILE	5.2
1	A	306	ALA	4.8
1	A	100	THR	4.7
1	A	65	TYR	4.6
1	A	300	ILE	4.3
1	B	164	ASN	4.2
1	A	317	TRP	3.9
1	B	298	ILE	3.8
1	A	32	THR	3.7
1	A	377	THR	3.7
1	A	315	VAL	3.6
1	B	296	GLN	3.6
1	A	304	GLY	3.6
1	B	301	SER	3.5
1	A	302	ARG	3.4
1	B	297	ARG	3.4
1	A	316	PRO	3.3
1	A	308	ASP	3.3
1	B	165	ASP	3.3
1	A	311	ALA	3.3
1	A	360	ASN	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	302	ARG	3.2
1	A	224	GLY	3.1
1	B	101[A]	HIS	3.0
1	A	307	GLY	3.0
1	A	313	PRO	3.0
1	B	293	LEU	2.9
1	B	166	LYS	2.9
1	A	312	ASN	2.9
1	B	168	ASP	2.9
1	B	292	ASN	2.9
1	B	225	ASP	2.8
1	A	161	ASN	2.8
1	A	305	ASN	2.7
1	A	151	LYS	2.7
1	A	303	ARG	2.7
1	B	230	GLN	2.7
1	B	15	GLY	2.6
1	B	44	LEU	2.6
1	B	14	ARG	2.6
1	B	294	ARG	2.5
1	A	208	HIS	2.5
1	B	228	LYS	2.5
1	B	98	ASP	2.5
1	A	163	ASN	2.5
1	B	4	ASN	2.5
1	A	192	ARG	2.4
1	B	100	THR	2.4
1	A	203	ILE	2.3
1	B	176	GLN	2.3
1	B	169	GLU	2.3
1	A	193	ASP	2.3
1	A	98	ASP	2.3
1	A	370	GLU	2.3
1	A	314	ALA	2.2
1	B	359	THR	2.2
1	B	148	ALA	2.1
1	B	42	ASP	2.1
1	A	200	ASP	2.1
1	B	208	HIS	2.1
1	A	204	THR	2.0
1	A	4	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BMA	C	3	11/12	0.69	0.12	52,55,61,64	0
2	NAG	C	2	14/15	0.83	0.11	38,44,48,52	0
2	NAG	C	1	14/15	0.86	0.10	26,30,39,43	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	A	401	14/15	0.58	0.22	46,51,56,59	0
3	NAG	B	401	14/15	0.70	0.16	38,48,51,51	0
4	SO4	B	404	5/5	0.74	0.18	75,75,82,82	0
4	SO4	A	402	5/5	0.75	0.14	43,63,75,77	0
4	SO4	B	402	5/5	0.94	0.08	34,41,45,52	0
4	SO4	B	403	5/5	0.96	0.13	23,24,31,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.