



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 27, 2023 – 07:26 AM EDT

PDB ID : 3I8V  
Title : Crystal structure of human PDE4a with 4-(3-butoxy-4-methoxyphenyl)methyl-2-imidazolidone  
Authors : Cheng, R.K.Y.; Crawley, L.; Barker, J.; Wood, M.; Felicetti, B.; Whittaker, M.  
Deposited on : 2009-07-10  
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	354	 2% 83% 11% • 5%
1	B	354	 2% 87% 6% • 6%

## 2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 5833 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP-specific 3',5'-cyclic phosphodiesterase 4A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	335	2708	1712	460	515	21	0	0	0
1	B	334	2713	1718	457	517	21	0	2	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	269	MET	-	expression tag	UNP P27815
A	270	GLY	-	expression tag	UNP P27815
A	271	SER	-	expression tag	UNP P27815
A	272	SER	-	expression tag	UNP P27815
A	273	HIS	-	expression tag	UNP P27815
A	274	HIS	-	expression tag	UNP P27815
A	275	HIS	-	expression tag	UNP P27815
A	276	HIS	-	expression tag	UNP P27815
A	277	HIS	-	expression tag	UNP P27815
A	278	HIS	-	expression tag	UNP P27815
A	279	SER	-	expression tag	UNP P27815
A	280	SER	-	expression tag	UNP P27815
A	281	GLY	-	expression tag	UNP P27815
A	282	LEU	-	expression tag	UNP P27815
A	283	VAL	-	expression tag	UNP P27815
A	284	PRO	-	expression tag	UNP P27815
A	285	ARG	-	expression tag	UNP P27815
A	286	GLY	-	expression tag	UNP P27815
A	287	SER	-	expression tag	UNP P27815
A	288	HIS	-	expression tag	UNP P27815
A	289	MET	-	expression tag	UNP P27815
B	269	MET	-	expression tag	UNP P27815
B	270	GLY	-	expression tag	UNP P27815
B	271	SER	-	expression tag	UNP P27815
B	272	SER	-	expression tag	UNP P27815

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	273	HIS	-	expression tag	UNP P27815
B	274	HIS	-	expression tag	UNP P27815
B	275	HIS	-	expression tag	UNP P27815
B	276	HIS	-	expression tag	UNP P27815
B	277	HIS	-	expression tag	UNP P27815
B	278	HIS	-	expression tag	UNP P27815
B	279	SER	-	expression tag	UNP P27815
B	280	SER	-	expression tag	UNP P27815
B	281	GLY	-	expression tag	UNP P27815
B	282	LEU	-	expression tag	UNP P27815
B	283	VAL	-	expression tag	UNP P27815
B	284	PRO	-	expression tag	UNP P27815
B	285	ARG	-	expression tag	UNP P27815
B	286	GLY	-	expression tag	UNP P27815
B	287	SER	-	expression tag	UNP P27815
B	288	HIS	-	expression tag	UNP P27815
B	289	MET	-	expression tag	UNP P27815

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	B	1	Total Mg 1 1	0	0

- Molecule 4 is (4R)-4-(3-butoxy-4-methoxybenzyl)imidazolidin-2-one (three-letter code: OMO) (formula: C<sub>15</sub>H<sub>22</sub>N<sub>2</sub>O<sub>3</sub>).



*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Co	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	172	Total	O	0	0
			172	172		
7	B	177	Total	O	0	0
			177	177		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4A



- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.41Å 105.41Å 165.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.33 – 2.25 45.33 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.9 (45.33-2.25) 100.0 (45.33-2.25)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.56 (at 2.24Å)	Xtrriage
Refinement program	REFMAC 5.5.0088	Depositor
R, $R_{free}$	0.187 , 0.222 0.185 , 0.220	Depositor DCC
$R_{free}$ test set	2262 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.3	Xtrriage
Anisotropy	0.053	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 35.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5833	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CO, OMO, ZN, MG, GOL, CSS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.54	0/2756	0.61	1/3736 (0.0%)
1	B	0.55	0/2767	0.63	1/3751 (0.0%)
All	All	0.54	0/5523	0.62	2/7487 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	472	LEU	CA-CB-CG	7.55	132.67	115.30
1	A	472	LEU	CA-CB-CG	7.51	132.57	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2708	0	2648	19	0
1	B	2713	0	2656	14	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	20	0	22	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	20	0	22	0	0
5	A	6	0	8	2	0
5	B	12	0	16	0	0
6	A	1	0	0	0	0
7	A	172	0	0	3	0
7	B	177	0	0	1	0
All	All	5833	0	5372	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:415:ASP:OD1	5:A:3:GOL:H2	1.76	0.84
1:A:366:HIS:HE1	5:A:3:GOL:H32	1.71	0.56
1:B:334:MET:HB3	1:B:358:MET:SD	2.47	0.55
1:B:396:VAL:HG11	1:B:512:LEU:HD12	1.87	0.55
1:A:539:GLU:HB2	7:A:83:HOH:O	2.09	0.51
1:B:498:MET:HE2	1:B:599:LEU:HD13	1.93	0.50
1:B:535:THR:HB	1:B:607:ILE:HG23	1.93	0.50
1:A:519:ILE:O	1:A:523:ARG:HG3	2.11	0.49
1:A:334:MET:HB3	1:A:358:MET:SD	2.52	0.49
1:A:457:ASN:ND2	7:A:16:HOH:O	2.38	0.48
1:B:514:ASN:HD22	1:B:514:ASN:N	2.12	0.47
1:A:535:THR:HB	1:A:607:ILE:HG23	1.97	0.46
1:A:556:GLY:HA3	1:A:570:CYS:O	2.15	0.46
1:B:619:TYR:HA	1:B:622:ILE:HD12	1.98	0.46
1:B:297:LYS:HG3	1:B:298:THR:N	2.31	0.46
1:A:585:ILE:HA	1:A:589:VAL:HB	1.99	0.45
7:A:652:HOH:O	1:B:429:SER:HB2	2.16	0.45
1:B:390:THR:HG22	1:B:393:LEU:HD12	2.00	0.43
1:B:514:ASN:N	1:B:514:ASN:ND2	2.67	0.43
1:A:296:VAL:HG21	1:A:325:ALA:HB1	2.01	0.42
1:A:505:THR:CG2	1:A:511:LEU:HD12	2.49	0.42
1:A:505:THR:HG22	1:A:511:LEU:HD12	2.00	0.42
1:A:512:LEU:HD22	1:A:517:ASP:HB3	2.00	0.42
1:A:363:ASP:OD1	1:B:558:ARG:HD3	2.19	0.42
1:A:317:ASN:ND2	1:A:540:LEU:HD21	2.34	0.41
1:A:328:ARG:NH2	1:B:561:GLU:OE1	2.50	0.41
1:A:533:ASN:HB2	1:A:534:PRO:HD3	2.01	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:482:ALA:HB1	1:A:491:LEU:HD21	2.02	0.41
1:B:502:LYS:NZ	7:B:669:HOH:O	2.52	0.41
1:A:333:ILE:HG23	1:A:381:LEU:HD13	2.03	0.41
1:B:556:GLY:HA3	1:B:570:CYS:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	332/354 (94%)	328 (99%)	3 (1%)	1 (0%)	41	46
1	B	333/354 (94%)	328 (98%)	5 (2%)	0	100	100
All	All	665/708 (94%)	656 (99%)	8 (1%)	1 (0%)	47	55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	506	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/316 (95%)	289 (96%)	11 (4%)	34	40

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	301/316 (95%)	295 (98%)	6 (2%)	55	64
All	All	601/632 (95%)	584 (97%)	17 (3%)	43	52

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	289	MET
1	A	351	VAL
1	A	381	LEU
1	A	390	THR
1	A	469	ARG
1	A	472	LEU
1	A	473	ARG
1	A	489	MET
1	A	504	VAL
1	A	565	GLU
1	A	620	SER
1	B	390	THR
1	B	451	LYS
1	B	472	LEU
1	B	514	ASN
1	B	598	ASP
1	B	606	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	290	ASN
1	A	457	ASN
1	A	470	GLN
1	A	524	ASN
1	B	366	HIS
1	B	457	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CSS	A	320	1	4,6,7	0.79	0	1,6,8	0.16	0
1	CSS	B	320	1	4,6,7	0.90	0	1,6,8	1.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSS	A	320	1	-	0/1/5/7	-
1	CSS	B	320	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 5 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	0MO	A	1	-	20,21,21	1.21	3 (15%)	25,27,27	3.21	7 (28%)
4	0MO	B	2	-	20,21,21	1.07	1 (5%)	25,27,27	2.45	7 (28%)
5	GOL	B	624	-	5,5,5	0.26	0	5,5,5	0.92	0
5	GOL	B	625	-	5,5,5	0.32	0	5,5,5	0.32	0
5	GOL	A	3	-	5,5,5	0.34	0	5,5,5	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	0MO	A	1	-	-	3/11/20/20	0/2/2/2
4	0MO	B	2	-	-	2/11/20/20	0/2/2/2
5	GOL	B	624	-	-	2/4/4/4	-
5	GOL	B	625	-	-	1/4/4/4	-
5	GOL	A	3	-	-	3/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1	0MO	C6-C5	3.93	1.49	1.40
4	B	2	0MO	C6-C5	3.83	1.48	1.40
4	A	1	0MO	C12-N2	-2.23	1.43	1.46
4	A	1	0MO	C11-C12	-2.13	1.49	1.53

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1	0MO	C11-C12-N2	-13.98	97.97	113.10
4	B	2	0MO	C11-C12-N2	-9.07	103.29	113.10
4	B	2	0MO	C11-C10-C15	-3.43	114.55	120.44
4	B	2	0MO	C7-O2-C6	3.35	122.58	117.53

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	2	0MO	O1-C5-C6	3.34	122.63	115.73
4	A	1	0MO	O1-C5-C6	3.27	122.48	115.73
4	A	1	0MO	C7-O2-C6	3.01	122.08	117.53
4	B	2	0MO	C13-C12-C11	-2.96	109.04	113.72
4	A	1	0MO	O2-C6-C5	2.88	119.42	115.41
4	A	1	0MO	C11-C10-C15	-2.68	115.83	120.44
4	A	1	0MO	O3-C14-N2	-2.61	122.19	125.94
4	A	1	0MO	C9-C10-C15	2.19	121.60	118.54
4	B	2	0MO	O1-C5-C15	-2.07	118.83	123.58
4	B	2	0MO	O2-C6-C5	2.05	118.26	115.41

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1	0MO	C10-C11-C12-N2
4	A	1	0MO	C10-C11-C12-C13
4	B	2	0MO	C10-C11-C12-N2
4	B	2	0MO	C10-C11-C12-C13
5	A	3	GOL	C1-C2-C3-O3
5	B	624	GOL	O1-C1-C2-O2
5	B	624	GOL	O1-C1-C2-C3
5	A	3	GOL	O1-C1-C2-C3
5	B	625	GOL	O1-C1-C2-C3
5	A	3	GOL	O2-C2-C3-O3
4	A	1	0MO	C1-C2-C3-C4


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	3	GOL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	334/354 (94%)	-0.27	7 (2%) 63 66	25, 35, 59, 75	0
1	B	333/354 (94%)	-0.28	7 (2%) 63 66	24, 36, 56, 70	0
All	All	667/708 (94%)	-0.28	14 (2%) 63 66	24, 36, 58, 75	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	507	SER	7.1
1	B	506	SER	4.8
1	A	506	SER	4.7
1	A	505	THR	4.1
1	A	513	ASP	3.6
1	A	508	GLY	3.6
1	B	508	GLY	3.6
1	B	507	SER	2.9
1	A	509	VAL	2.5
1	A	504	VAL	2.2
1	B	306	GLN	2.1
1	B	302	GLU	2.1
1	B	622	ILE	2.0
1	B	511	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	CSS	B	320	7/8	0.93	0.12	39,41,47,55	0
1	CSS	A	320	7/8	0.97	0.10	36,38,42,51	0

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	GOL	A	3	6/6	0.66	0.27	69,71,71,72	0
6	CO	A	624	1/1	0.80	0.10	110,110,110,110	0
5	GOL	B	625	6/6	0.81	0.20	77,78,79,79	0
5	GOL	B	624	6/6	0.81	0.20	50,52,52,52	0
4	OMO	A	1	20/20	0.88	0.18	37,43,54,55	0
4	OMO	B	2	20/20	0.93	0.14	38,42,55,56	0
3	MG	B	623	1/1	0.99	0.10	24,24,24,24	0
2	ZN	B	1	1/1	1.00	0.09	33,33,33,33	0
3	MG	A	623	1/1	1.00	0.13	26,26,26,26	0
2	ZN	A	2	1/1	1.00	0.09	31,31,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.