



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 2, 2024 – 05:33 PM EST

PDB ID : 3IBT
Title : Structure of 1H-3-hydroxy-4-oxoquinoline 2,4-dioxygenase (QDO)
Authors : Oakley, A.J.
Deposited on : 2009-07-16
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.21
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3775 (2.60-2.60)
Clashscore	180529	4181 (2.60-2.60)
Ramachandran outliers	177936	4129 (2.60-2.60)
Sidechain outliers	177891	4129 (2.60-2.60)
RSRZ outliers	164620	3775 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	264	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2170 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1H-3-hydroxy-4-oxoquinoline 2,4-dioxygenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	263	2138	1356	380	390	6	6	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	118	ILE	VAL	conflict	UNP O33472
A	200	ASP	GLU	conflict	UNP O33472
A	226	GLU	ASP	conflict	UNP O33472


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	32	Total	O	0	0
			32	32		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 1H-3-hydroxy-4-oxoquinoline 2,4-dioxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	90.14Å 90.14Å 168.74Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.60 20.00 – 2.60	Depositor EDS
% Data completeness (in resolution range)	96.3 (20.00-2.60) 96.0 (20.00-2.60)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.59 (at 2.59Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.237 , 0.279 0.241 , 0.298	Depositor DCC
R_{free} test set	617 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	63.2	Xtrriage
Anisotropy	0.061	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 78.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.83	EDS
Total number of atoms	2170	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.95	9/2202 (0.4%)	0.87	4/2992 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	56	ARG	CZ-NH1	9.14	1.45	1.33
1	A	76	ASP	CG-OD1	8.82	1.45	1.25
1	A	176	GLU	CD-OE1	7.18	1.33	1.25
1	A	176	GLU	CD-OE2	6.44	1.32	1.25
1	A	56	ARG	NE-CZ	6.16	1.41	1.33
1	A	76	ASP	CG-OD2	6.04	1.39	1.25
1	A	176	GLU	CG-CD	5.46	1.60	1.51
1	A	7	ASN	CG-OD1	5.10	1.35	1.24
1	A	7	ASN	CG-ND2	5.02	1.45	1.32

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	56	ARG	NE-CZ-NH2	-7.64	116.48	120.30
1	A	33	ASP	CB-CG-OD1	-6.42	112.52	118.30
1	A	76	ASP	CB-CG-OD1	-6.00	112.90	118.30
1	A	190	ARG	NE-CZ-NH1	5.01	122.81	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	109	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2138	0	2011	70	0
2	A	32	0	0	1	0
All	All	2170	0	2011	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:89:PHE:HE1	1:A:91:MSE:HE3	1.10	1.04
1:A:89:PHE:CE1	1:A:91:MSE:HE3	2.03	0.92
1:A:8:GLY:H	1:A:56:ARG:HH22	1.22	0.86
1:A:243:THR:CG2	1:A:245:PHE:H	1.97	0.78
1:A:10:LEU:HG	1:A:11:MSE:N	2.04	0.73
1:A:243:THR:HG22	1:A:245:PHE:H	1.53	0.73
1:A:214:GLN:HG2	1:A:215:PRO:HA	1.71	0.72
1:A:243:THR:HB	1:A:249:GLU:OE1	1.91	0.70
1:A:92:VAL:HG22	1:A:117:ILE:HB	1.75	0.69
1:A:26:LEU:HD13	1:A:37:PHE:CD2	2.29	0.68
1:A:8:GLY:N	1:A:56:ARG:HH22	1.92	0.67
1:A:73:LEU:O	1:A:76:ASP:HB2	1.95	0.67
1:A:121:TRP:H	1:A:210:HIS:HE2	1.43	0.65
1:A:113:LEU:HD12	1:A:113:LEU:O	1.97	0.63
1:A:1:MSE:N	1:A:1:MSE:HE3	2.15	0.61
1:A:87:ARG:O	1:A:112:ARG:NH1	2.33	0.61
1:A:3:SER:HB3	1:A:12:THR:HG22	1.81	0.60
1:A:10:LEU:HG	1:A:11:MSE:H	1.67	0.60
1:A:105:CYS:HA	1:A:113:LEU:HD11	1.83	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:69:ASP:HB3	1:A:71:GLN:H	1.68	0.58
1:A:146:ARG:HA	1:A:178:TRP:CZ2	2.40	0.56
1:A:182:CYS:O	1:A:186:GLU:HB2	2.06	0.56
1:A:78:LEU:HD22	1:A:108:LEU:HD11	1.89	0.54
1:A:208:ILE:HD12	1:A:233:TRP:CZ3	2.43	0.53
1:A:157:THR:HB	1:A:159:ASN:H	1.74	0.52
1:A:169:GLU:O	1:A:172:TRP:HB3	2.10	0.51
1:A:81:ILE:HD12	1:A:82:ASP:N	2.25	0.51
1:A:146:ARG:HG2	1:A:147:GLN:N	2.24	0.51
1:A:159:ASN:HD22	1:A:162:VAL:H	1.58	0.51
1:A:51:ILE:O	1:A:53:PRO:HD3	2.11	0.50
1:A:78:LEU:HD21	1:A:108:LEU:HD13	1.93	0.49
1:A:235:HIS:HB3	1:A:237:ARG:HH12	1.76	0.49
1:A:3:SER:HB3	1:A:12:THR:HA	1.94	0.49
1:A:46:ARG:HH21	1:A:259:ARG:HD3	1.78	0.49
1:A:119:ILE:CG2	1:A:246:PRO:HG2	2.43	0.49
1:A:11:MSE:HB3	1:A:54:ASP:O	2.12	0.49
1:A:8:GLY:H	1:A:56:ARG:NH2	2.03	0.49
1:A:78:LEU:CD2	1:A:108:LEU:CD1	2.90	0.49
1:A:3:SER:HB2	1:A:10:LEU:HD11	1.93	0.48
1:A:5:ASN:HB3	1:A:10:LEU:HD12	1.95	0.48
1:A:64:ASP:O	1:A:65:SER:HB2	2.13	0.48
1:A:46:ARG:NH2	1:A:259:ARG:HD3	2.29	0.48
1:A:4:LEU:HD21	1:A:83:ALA:HB3	1.96	0.48
1:A:225:LEU:HD13	2:A:265:HOH:O	2.13	0.48
1:A:55:TRP:O	1:A:58:HIS:HB2	2.13	0.48
1:A:75:GLN:HE22	1:A:107:GLN:HE22	1.63	0.47
1:A:61:LYS:O	1:A:62:GLN:C	2.53	0.46
1:A:78:LEU:HD22	1:A:108:LEU:CD1	2.45	0.46
1:A:243:THR:HG23	1:A:245:PHE:H	1.75	0.46
1:A:243:THR:HG22	1:A:245:PHE:N	2.28	0.45
1:A:128:GLY:O	1:A:132:GLN:HG3	2.16	0.45
1:A:118:ILE:HD12	1:A:121:TRP:HE3	1.81	0.45
1:A:192:TRP:CD2	1:A:198:ARG:HD3	2.52	0.45
1:A:26:LEU:HB2	1:A:52:CYS:SG	2.57	0.44
1:A:78:LEU:CD2	1:A:108:LEU:HD11	2.47	0.43
1:A:64:ASP:O	1:A:65:SER:CB	2.65	0.43
1:A:1:MSE:HE3	1:A:1:MSE:H3	1.82	0.43
1:A:62:GLN:HE22	1:A:174:HIS:H	1.66	0.43
1:A:67:ASP:HB2	1:A:180:ARG:NH2	2.34	0.43
1:A:82:ASP:OD1	1:A:82:ASP:C	2.58	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:ILE:HD11	1:A:262:LEU:CD1	2.50	0.42
1:A:10:LEU:CG	1:A:11:MSE:N	2.78	0.42
1:A:9:THR:HB	1:A:10:LEU:H	1.53	0.41
1:A:78:LEU:CD2	1:A:108:LEU:HD13	2.49	0.41
1:A:13:TYR:CD1	1:A:51:ILE:HG23	2.56	0.41
1:A:210:HIS:ND1	1:A:224:GLN:NE2	2.67	0.41
1:A:98:CYS:HB2	1:A:121:TRP:CE2	2.56	0.41
1:A:102:ILE:HD13	1:A:102:ILE:HA	1.92	0.40
1:A:200:ASP:O	1:A:205:LYS:NZ	2.54	0.40
1:A:55:TRP:O	1:A:56:ARG:C	2.60	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	261/264 (99%)	236 (90%)	23 (9%)	2 (1%)	16 34

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	65	SER
1	A	62	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	228/222 (103%)	201 (88%)	27 (12%)	4 8

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MSE
1	A	4	LEU
1	A	5	ASN
1	A	10	LEU
1	A	11	MSE
1	A	12	THR
1	A	51	ILE
1	A	64	ASP
1	A	67	ASP
1	A	77	LEU
1	A	80	PHE
1	A	82	ASP
1	A	88	ASP
1	A	94	THR
1	A	112	ARG
1	A	120	ASP
1	A	140	THR
1	A	146	ARG
1	A	155	GLU
1	A	156	THR
1	A	157	THR
1	A	159	ASN
1	A	225	LEU
1	A	237	ARG
1	A	242	ARG
1	A	243	THR
1	A	245	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	75	GLN
1	A	107	GLN
1	A	132	GLN
1	A	159	ASN
1	A	224	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Warning: The R factor obtained from EDS is 0.3778, which does not match the depositor's R factor of 0.23692. Please interpret the results in this section carefully.

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/264 (97%)	1.93	110 (42%) 1 1	65, 72, 83, 98	0

All (110) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	14	SER	7.0
1	A	6	VAL	5.5
1	A	79	ALA	5.5
1	A	5	ASN	4.9
1	A	10	LEU	4.7
1	A	172	TRP	4.7
1	A	232	SER	4.4
1	A	234	PHE	4.3
1	A	233	TRP	4.3
1	A	54	ASP	4.2
1	A	228	ALA	4.2
1	A	150	PHE	4.2
1	A	9	THR	4.1
1	A	63	THR	4.1
1	A	212	TYR	4.1
1	A	74	ALA	4.0
1	A	222	GLN	4.0
1	A	198	ARG	3.9
1	A	146	ARG	3.7
1	A	236	PRO	3.7
1	A	179	GLN	3.7
1	A	62	GLN	3.6
1	A	206	PRO	3.5
1	A	132	GLN	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	68	PHE	3.4
1	A	142	TYR	3.4
1	A	235	HIS	3.4
1	A	4	LEU	3.3
1	A	22	PRO	3.3
1	A	12	THR	3.2
1	A	58	HIS	3.2
1	A	225	LEU	3.2
1	A	117	ILE	3.1
1	A	218	GLN	3.1
1	A	263	GLN	3.1
1	A	64	ASP	3.1
1	A	231	HIS	3.1
1	A	195	PRO	3.1
1	A	205	LYS	3.1
1	A	8	GLY	3.0
1	A	2	GLN	3.0
1	A	262	LEU	3.0
1	A	50	VAL	3.0
1	A	252	VAL	3.0
1	A	59	ASP	2.9
1	A	214	GLN	2.9
1	A	159	ASN	2.9
1	A	173	PHE	2.9
1	A	176	GLU	2.9
1	A	210	HIS	2.9
1	A	111	ALA	2.9
1	A	3	SER	2.8
1	A	67	ASP	2.8
1	A	33	ASP	2.8
1	A	148	SER	2.7
1	A	227	PHE	2.7
1	A	192	TRP	2.7
1	A	83	ALA	2.7
1	A	224	GLN	2.7
1	A	28	SER	2.6
1	A	76	ASP	2.6
1	A	238	HIS	2.6
1	A	116	THR	2.6
1	A	34	HIS	2.6
1	A	93	SER	2.5
1	A	171	PRO	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	215	PRO	2.5
1	A	255	ALA	2.5
1	A	128	GLY	2.5
1	A	174	HIS	2.5
1	A	138	HIS	2.4
1	A	229	ALA	2.4
1	A	87	ARG	2.4
1	A	70	SER	2.4
1	A	52	CYS	2.4
1	A	219	ASP	2.4
1	A	220	TYR	2.4
1	A	175	GLY	2.4
1	A	72	THR	2.4
1	A	140	THR	2.4
1	A	19	PRO	2.4
1	A	108	LEU	2.3
1	A	20	HIS	2.3
1	A	92	VAL	2.3
1	A	217	SER	2.3
1	A	69	ASP	2.3
1	A	133	LEU	2.3
1	A	259	ARG	2.3
1	A	216	LEU	2.3
1	A	207	GLU	2.2
1	A	203	PRO	2.2
1	A	123	LEU	2.2
1	A	240	PRO	2.1
1	A	82	ASP	2.1
1	A	137	GLN	2.1
1	A	256	GLN	2.1
1	A	60	ALA	2.1
1	A	180	ARG	2.1
1	A	260	GLU	2.1
1	A	61	LYS	2.1
1	A	78	LEU	2.1
1	A	209	CYS	2.1
1	A	147	GLN	2.1
1	A	88	ASP	2.1
1	A	96	HIS	2.1
1	A	13	TYR	2.1
1	A	17	GLY	2.1
1	A	136	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	197	ASP	2.0
1	A	125	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.