



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2024 – 11:03 PM EST

PDB ID : 3PHG  
Title : Crystal structure of the Shikimate 5-Dehydrogenase (aroE) from *Helicobacter pylori*  
Authors : Cheng, W.C.; Lin, S.C.; Wang, W.C.  
Deposited on : 2010-11-04  
Resolution : 1.57 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5534 (1.60-1.56)
Clashscore	141614	5861 (1.60-1.56)
Ramachandran outliers	138981	5708 (1.60-1.56)
Sidechain outliers	138945	5703 (1.60-1.56)
RSRZ outliers	127900	5431 (1.60-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	 4% (poor fit), 78% (0-1 outliers), 15% (2-3 outliers), 3% (not modelled)
1	B	269	 6% (poor fit), 81% (0-1 outliers), 12% (2-3 outliers), 3% (not modelled)

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4374 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Shikimate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	258	2023	1318	333	362	10	0	0	0
1	B	255	1996	1307	323	356	10	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	264	HIS	-	expression tag	UNP P56119
A	265	HIS	-	expression tag	UNP P56119
A	266	HIS	-	expression tag	UNP P56119
A	267	HIS	-	expression tag	UNP P56119
A	268	HIS	-	expression tag	UNP P56119
A	269	HIS	-	expression tag	UNP P56119
B	264	HIS	-	expression tag	UNP P56119
B	265	HIS	-	expression tag	UNP P56119
B	266	HIS	-	expression tag	UNP P56119
B	267	HIS	-	expression tag	UNP P56119
B	268	HIS	-	expression tag	UNP P56119
B	269	HIS	-	expression tag	UNP P56119


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	176	Total 176	O 176	0	0
2	B	179	Total 179	O 179	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Shikimate dehydrogenase



- Molecule 1: Shikimate dehydrogenase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.33Å 46.18Å 118.70Å 90.00° 99.04° 90.00°	Depositor
Resolution (Å)	23.56 – 1.57 23.55 – 1.57	Depositor EDS
% Data completeness (in resolution range)	98.3 (23.56-1.57) 98.3 (23.55-1.57)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.76 (at 1.57Å)	Xtrriage
Refinement program	REFMAC 5.5.0088	Depositor
R, $R_{free}$	0.200 , 0.240 0.198 , 0.239	Depositor DCC
$R_{free}$ test set	3378 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.0	Xtrriage
Anisotropy	0.811	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 62.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.022 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4374	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.70	0/2067	0.79	1/2781 (0.0%)
1	B	0.71	0/2041	0.79	2/2747 (0.1%)
All	All	0.70	0/4108	0.79	3/5528 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	160	LEU	CA-CB-CG	5.92	128.91	115.30
1	B	38	HIS	O-C-N	5.20	131.01	122.70
1	A	44	LEU	CB-CG-CD2	5.06	119.60	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2023	0	2062	45	0
1	B	1996	0	2032	43	0
2	A	176	0	0	11	0
2	B	179	0	0	7	0
All	All	4374	0	4094	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3:LEU:HD21	1:B:38:HIS:CE1	1.85	1.09
1:B:3:LEU:HD21	1:B:38:HIS:ND1	1.74	1.02
1:B:38:HIS:HD2	1:B:40:HIS:NE2	1.58	1.01
1:B:3:LEU:CD2	1:B:38:HIS:CE1	2.45	0.99
1:A:254:LYS:HE2	1:A:254:LYS:HA	1.52	0.90
1:B:129:SER:HB3	1:B:208:LEU:HD21	1.59	0.84
1:A:14:LYS:HE3	2:B:318:HOH:O	1.79	0.83
1:A:14:LYS:HD3	1:B:185:HIS:CE1	2.15	0.81
1:A:50:ILE:HD11	1:A:68:PHE:HB3	1.64	0.80
1:B:129:SER:HB3	1:B:208:LEU:CD2	2.15	0.76
1:B:46:LEU:HD13	2:B:270:HOH:O	1.85	0.75
1:B:3:LEU:CD2	1:B:38:HIS:ND1	2.48	0.75
1:B:38:HIS:CD2	1:B:40:HIS:NE2	2.50	0.75
1:A:125:GLY:C	1:A:148:ASN:HD21	1.91	0.74
1:B:231:LYS:HE3	2:B:405:HOH:O	1.90	0.71
1:B:3:LEU:HD22	1:B:38:HIS:HE1	1.56	0.70
1:A:38:HIS:HD2	1:A:40:HIS:NE2	1.90	0.69
1:A:213:LEU:N	2:A:338:HOH:O	2.25	0.69
1:B:3:LEU:HD22	1:B:38:HIS:CE1	2.29	0.67
1:B:44:LEU:HD22	1:B:48:SER:CB	2.26	0.65
1:A:66:LEU:CD1	1:A:66:LEU:H	2.11	0.64
1:A:56:HIS:HD2	2:A:324:HOH:O	1.81	0.64
1:A:191:ASN:HD22	1:A:194:VAL:H	1.47	0.63
1:A:1:MET:HB2	2:A:389:HOH:O	1.99	0.62
1:A:15:HIS:HB3	1:B:14:LYS:NZ	2.14	0.62
1:A:73:PHE:CE1	1:A:79:ILE:HD13	2.35	0.62
1:B:44:LEU:HD22	1:B:48:SER:HB2	1.81	0.62
1:B:3:LEU:CD2	1:B:38:HIS:HE1	2.03	0.61
1:A:66:LEU:H	1:A:66:LEU:HD12	1.66	0.61
1:A:78:LYS:HE2	1:A:98:GLU:OE2	2.02	0.60
1:A:125:GLY:CA	1:A:148:ASN:HD21	2.15	0.60
1:B:129:SER:CB	1:B:208:LEU:CD2	2.80	0.59
1:A:80:LYS:HD3	1:A:100:VAL:HG11	1.85	0.58
1:A:139:LYS:HE3	2:A:402:HOH:O	2.02	0.58
1:B:82:ILE:HD11	1:B:139:LYS:NZ	2.19	0.57
1:A:254:LYS:HA	1:A:254:LYS:CE	2.30	0.56
1:B:46:LEU:HD12	1:B:46:LEU:N	2.21	0.55
1:B:44:LEU:HD21	1:B:53:GLU:HG3	1.87	0.54
1:B:231:LYS:HG3	1:B:263:PHE:CE2	2.42	0.54
1:A:190:LEU:HD12	1:A:195:LEU:HD13	1.91	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:GLU:HG3	1:A:140:GLN:HE21	1.72	0.53
1:A:186:ASN:ND2	2:A:277:HOH:O	2.42	0.52
1:B:48:SER:O	1:B:71:ARG:NH2	2.43	0.52
1:B:153:GLY:O	1:B:154:LEU:C	2.48	0.51
1:B:31:LYS:HG2	1:B:32:GLU:OE2	2.10	0.51
1:A:31:LYS:H	1:A:31:LYS:HD2	1.75	0.50
1:B:44:LEU:HD22	1:B:48:SER:HB3	1.94	0.50
1:A:15:HIS:HB3	1:B:14:LYS:HZ1	1.76	0.49
1:B:143:GLN:NE2	2:B:400:HOH:O	2.45	0.49
1:A:14:LYS:HG3	2:B:318:HOH:O	2.12	0.49
1:B:143:GLN:NE2	2:B:284:HOH:O	2.46	0.48
1:B:217:LEU:HD22	1:B:227:PHE:HB2	1.95	0.48
1:A:14:LYS:CD	1:B:185:HIS:CE1	2.94	0.48
1:A:191:ASN:HD22	1:A:194:VAL:HG23	1.79	0.48
1:A:136:GLU:HG3	1:A:140:GLN:NE2	2.27	0.47
1:A:257:GLU:HG2	2:A:377:HOH:O	2.14	0.47
1:B:47:GLU:H	1:B:47:GLU:CD	2.17	0.47
1:A:196:LYS:HG2	1:A:223:LEU:HD21	1.96	0.47
1:B:129:SER:CB	1:B:208:LEU:HD23	2.44	0.46
1:B:50:ILE:HD11	1:B:68:PHE:HB3	1.97	0.46
1:A:33:LEU:HD21	1:A:246:SER:HB2	1.97	0.46
1:A:254:LYS:HE3	2:A:377:HOH:O	2.15	0.46
1:A:31:LYS:HB3	1:A:31:LYS:HE3	1.75	0.46
1:B:46:LEU:N	1:B:46:LEU:CD1	2.78	0.45
1:B:143:GLN:HB3	2:B:400:HOH:O	2.17	0.44
1:A:221:LYS:HG3	1:A:227:PHE:CE2	2.53	0.44
1:A:16:SER:HB3	2:A:382:HOH:O	2.18	0.44
1:A:14:LYS:HE2	1:B:12:PRO:HG3	2.00	0.43
1:A:44:LEU:HD21	1:A:53:GLU:HG3	1.98	0.43
1:B:78:LYS:HB2	1:B:78:LYS:HE2	1.85	0.43
1:B:46:LEU:CD1	1:B:46:LEU:H	2.32	0.43
1:A:109:PHE:HE2	1:A:137:LEU:HD21	1.84	0.43
1:A:150:SER:OG	1:A:152:ARG:HG3	2.18	0.43
1:B:73:PHE:CE1	1:B:79:ILE:HD13	2.54	0.42
1:B:82:ILE:CD1	1:B:139:LYS:NZ	2.82	0.42
1:A:31:LYS:H	1:A:31:LYS:CD	2.32	0.42
1:A:114:LYS:HG2	2:A:434:HOH:O	2.19	0.41
1:A:66:LEU:HD12	1:A:66:LEU:N	2.31	0.41
1:A:153:GLY:CA	2:A:320:HOH:O	2.68	0.41
1:A:15:HIS:HB3	1:B:14:LYS:HZ3	1.86	0.41
1:B:11:ASN:ND2	1:B:46:LEU:HD12	2.36	0.40

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:47:GLU:HG2	2:A:429:HOH:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/269 (94%)	251 (100%)	1 (0%)	0	100	100
1	B	249/269 (93%)	247 (99%)	2 (1%)	0	100	100
All	All	501/538 (93%)	498 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/229 (96%)	210 (96%)	9 (4%)	30	8
1	B	215/229 (94%)	207 (96%)	8 (4%)	34	9
All	All	434/458 (95%)	417 (96%)	17 (4%)	32	9

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LYS
1	A	31	LYS
1	A	44	LEU
1	A	46	LEU
1	A	66	LEU
1	A	186	ASN
1	A	195	LEU
1	A	237	GLN
1	A	254	LYS
1	B	3	LEU
1	B	27	LEU
1	B	31	LYS
1	B	97	ASP
1	B	113	LEU
1	B	160	LEU
1	B	175	LEU
1	B	237	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	HIS
1	A	30	GLN
1	A	38	HIS
1	A	56	HIS
1	A	63	ASN
1	A	96	ASN
1	A	119	GLN
1	A	140	GLN
1	A	143	GLN
1	A	148	ASN
1	A	191	ASN
1	A	228	GLN
1	B	22	HIS
1	B	38	HIS
1	B	49	HIS
1	B	63	ASN
1	B	185	HIS
1	B	228	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	258/269 (95%)	0.27	11 (4%) 35 35	9, 16, 28, 38	0
1	B	255/269 (94%)	0.34	16 (6%) 20 20	10, 17, 28, 35	0
All	All	513/538 (95%)	0.31	27 (5%) 26 26	9, 16, 28, 38	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	7.5
1	B	96	ASN	4.8
1	A	152	ARG	4.7
1	B	153	GLY	4.7
1	B	1	MET	4.3
1	B	38	HIS	3.8
1	B	154	LEU	3.8
1	A	34	ARG	3.5
1	B	34	ARG	3.3
1	A	114	LYS	3.3
1	A	184	LEU	3.0
1	A	96	ASN	3.0
1	A	27	LEU	2.9
1	B	2	LYS	2.9
1	B	97	ASP	2.8
1	B	114	LYS	2.8
1	B	212	PHE	2.7
1	A	14	LYS	2.7
1	B	3	LEU	2.6
1	B	86	CYS	2.5
1	A	191	ASN	2.4
1	B	257	GLU	2.3
1	B	64	VAL	2.2
1	B	98	GLU	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	56	HIS	2.1
1	A	201	GLU	2.1
1	A	240	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.