



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 12, 2024 – 06:29 PM EST

PDB ID : 3T61
Title : Crystal Structure of a gluconokinase from Sinorhizobium meliloti 1021
Authors : Kumaran, D.; Chamala, S.; Evans, B.; Foti, R.; Gizzi, A.; Hillerich, B.; Kar, A.; LaFleur, J.; Seidel, R.; Villigas, G.; Zencheck, W.; Almo, S.C.; Swaminathan, S.; New York Structural Genomics Research Consortium (NYSGRG)
Deposited on : 2011-07-28
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	202	 65% 21% 13%
1	B	202	 65% 22% 13%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2860 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gluconokinase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	176	1353	848	247	252	2	4	0	0	0
1	B	175	1343	842	244	251	2	4	0	0	0

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MSE	-	expression tag	UNP Q92VK3
A	0	VAL	-	expression tag	UNP Q92VK3
A	179	ALA	-	expression tag	UNP Q92VK3
A	180	GLU	-	expression tag	UNP Q92VK3
A	181	ASN	-	expression tag	UNP Q92VK3
A	182	LEU	-	expression tag	UNP Q92VK3
A	183	TYR	-	expression tag	UNP Q92VK3
A	184	PHE	-	expression tag	UNP Q92VK3
A	185	GLN	-	expression tag	UNP Q92VK3
A	186	SER	-	expression tag	UNP Q92VK3
A	187	HIS	-	expression tag	UNP Q92VK3
A	188	HIS	-	expression tag	UNP Q92VK3
A	189	HIS	-	expression tag	UNP Q92VK3
A	190	HIS	-	expression tag	UNP Q92VK3
A	191	HIS	-	expression tag	UNP Q92VK3
A	192	HIS	-	expression tag	UNP Q92VK3
A	193	TRP	-	expression tag	UNP Q92VK3
A	194	SER	-	expression tag	UNP Q92VK3
A	195	HIS	-	expression tag	UNP Q92VK3
A	196	PRO	-	expression tag	UNP Q92VK3
A	197	GLN	-	expression tag	UNP Q92VK3
A	198	PHE	-	expression tag	UNP Q92VK3
A	199	GLU	-	expression tag	UNP Q92VK3
A	200	LYS	-	expression tag	UNP Q92VK3
B	-1	MSE	-	expression tag	UNP Q92VK3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	VAL	-	expression tag	UNP Q92VK3
B	179	ALA	-	expression tag	UNP Q92VK3
B	180	GLU	-	expression tag	UNP Q92VK3
B	181	ASN	-	expression tag	UNP Q92VK3
B	182	LEU	-	expression tag	UNP Q92VK3
B	183	TYR	-	expression tag	UNP Q92VK3
B	184	PHE	-	expression tag	UNP Q92VK3
B	185	GLN	-	expression tag	UNP Q92VK3
B	186	SER	-	expression tag	UNP Q92VK3
B	187	HIS	-	expression tag	UNP Q92VK3
B	188	HIS	-	expression tag	UNP Q92VK3
B	189	HIS	-	expression tag	UNP Q92VK3
B	190	HIS	-	expression tag	UNP Q92VK3
B	191	HIS	-	expression tag	UNP Q92VK3
B	192	HIS	-	expression tag	UNP Q92VK3
B	193	TRP	-	expression tag	UNP Q92VK3
B	194	SER	-	expression tag	UNP Q92VK3
B	195	HIS	-	expression tag	UNP Q92VK3
B	196	PRO	-	expression tag	UNP Q92VK3
B	197	GLN	-	expression tag	UNP Q92VK3
B	198	PHE	-	expression tag	UNP Q92VK3
B	199	GLU	-	expression tag	UNP Q92VK3
B	200	LYS	-	expression tag	UNP Q92VK3

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	78	Total	O	0	0
			78	78		
3	B	76	Total	O	0	0
			76	76		

4 Data and refinement statistics i

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	68.20Å 68.20Å 118.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.68 – 2.20 44.68 – 2.20	Depositor EDS
% Data completeness (in resolution range)	97.6 (44.68-2.20) 97.6 (44.68-2.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.63 (at 2.20Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.237 , 0.250 0.237 , 0.255	Depositor DCC
R_{free} test set	1862 reflections (6.93%)	wwPDB-VP
Wilson B-factor (Å ²)	32.1	Xtrriage
Anisotropy	0.107	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 23.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.488 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2860	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/1379	0.59	0/1861
1	B	0.36	0/1368	0.61	0/1846
All	All	0.36	0/2747	0.60	0/3707

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1353	0	1338	37	0
1	B	1343	0	1331	28	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	78	0	0	2	0
3	B	76	0	0	1	0
All	All	2860	0	2669	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:ARG:HH11	1:A:180:GLU:HA	1.20	1.01
1:A:13:ARG:NH1	1:A:180:GLU:HA	1.76	1.00
1:B:25:SER:H	1:B:142:GLN:HE21	1.23	0.86
1:A:25:SER:H	1:A:142:GLN:HE21	1.18	0.86
1:A:124:GLU:HG2	3:A:226:HOH:O	1.93	0.69
1:B:30:SER:O	1:B:34:GLU:HG3	1.93	0.68
1:B:135:PRO:HD2	1:B:138:LEU:HD12	1.76	0.66
1:B:56:ILE:HA	1:B:59:MSE:HE3	1.77	0.66
1:A:135:PRO:HD2	1:A:138:LEU:HD12	1.79	0.65
1:A:61:GLU:O	1:B:134:MSE:HG2	1.96	0.65
1:B:55:ASN:HB3	1:B:59:MSE:HE2	1.78	0.64
1:A:134:MSE:HG2	1:B:61:GLU:O	1.97	0.64
1:A:13:ARG:NH1	1:A:183:TYR:HB2	2.13	0.63
1:A:25:SER:H	1:A:142:GLN:NE2	1.94	0.62
1:A:136:SER:O	1:A:140:GLN:HG3	2.01	0.60
1:A:179:ALA:O	1:A:183:TYR:HD1	1.84	0.60
1:B:45:ILE:HB	1:B:89:VAL:HG22	1.85	0.59
1:A:140:GLN:O	1:A:144:GLU:HG3	2.04	0.58
1:A:180:GLU:HB3	3:A:221:HOH:O	2.04	0.56
1:A:119:GLU:OE1	1:A:140:GLN:HG2	2.07	0.55
1:B:96:ARG:HG3	1:B:96:ARG:HH11	1.70	0.55
1:B:50:LEU:HD12	1:B:77:ILE:HD13	1.88	0.54
1:B:51:HIS:CG	1:B:59:MSE:HE1	2.43	0.54
1:A:121:VAL:O	1:A:125:ARG:HG3	2.08	0.53
1:B:116:HIS:HE1	3:B:260:HOH:O	1.91	0.53
1:B:34:GLU:HG2	1:B:44:PHE:CD2	2.44	0.52
1:A:30:SER:O	1:A:34:GLU:HG3	2.10	0.51
1:A:50:LEU:HD12	1:A:77:ILE:HD13	1.92	0.51
1:A:183:TYR:O	1:A:187:HIS:HD2	1.94	0.50
1:B:173:ALA:O	1:B:177:ARG:HG3	2.10	0.50
1:A:15:PHE:CD2	1:A:19:ILE:HD11	2.47	0.49
1:A:45:ILE:HD12	1:A:87:VAL:HG21	1.94	0.49
1:A:59:MSE:HE2	1:B:60:SER:OG	2.12	0.48
1:B:15:PHE:HA	1:B:179:ALA:HB2	1.96	0.48
1:A:34:GLU:HG2	1:A:44:PHE:CD2	2.49	0.47
1:B:47:GLY:HA3	1:B:90:SER:O	2.14	0.47
1:B:56:ILE:CA	1:B:59:MSE:HE3	2.44	0.47
1:A:19:ILE:CD1	1:A:111:ALA:HB3	2.46	0.47
1:A:46:GLU:OE1	1:B:57:ARG:NH1	2.48	0.46
1:A:47:GLY:HA3	1:A:90:SER:O	2.15	0.46
1:A:13:ARG:HH12	1:A:183:TYR:HB2	1.80	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:PRO:HG3	1:B:49:ALA:HB2	1.98	0.44
1:A:19:ILE:HD13	1:A:111:ALA:HB3	2.00	0.43
1:B:181:ASN:O	1:B:185:GLN:HG3	2.17	0.43
1:A:45:ILE:HB	1:A:89:VAL:HG22	1.98	0.43
1:A:187:HIS:O	1:A:188:HIS:C	2.57	0.42
1:A:53:PRO:HB3	1:B:46:GLU:HG2	2.02	0.42
1:A:98:TYR:N	1:A:98:TYR:CD1	2.87	0.42
1:B:85:GLU:HA	1:B:86:PRO:HD3	1.91	0.42
1:B:127:HIS:HA	1:B:132:HIS:O	2.20	0.42
1:A:96:ARG:HD2	1:A:100:ASP:OD2	2.19	0.42
1:B:69:ASP:O	1:B:72:PRO:HD2	2.20	0.42
1:B:125:ARG:NH1	1:B:126:MSE:CE	2.83	0.42
1:A:55:ASN:O	1:A:59:MSE:HG3	2.20	0.41
1:A:127:HIS:HA	1:A:132:HIS:O	2.20	0.41
1:A:13:ARG:HB3	1:A:14:ARG:H	1.44	0.41
1:A:54:GLU:OE2	1:A:58:LYS:HE3	2.22	0.40
1:B:98:TYR:N	1:B:98:TYR:CD1	2.88	0.40
1:B:56:ILE:N	1:B:59:MSE:HE3	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	174/202 (86%)	168 (97%)	6 (3%)	0	100	100
1	B	173/202 (86%)	169 (98%)	4 (2%)	0	100	100
All	All	347/404 (86%)	337 (97%)	10 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	143/160 (89%)	142 (99%)	1 (1%)	81 90
1	B	142/160 (89%)	138 (97%)	4 (3%)	38 51
All	All	285/320 (89%)	280 (98%)	5 (2%)	54 69

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	SER
1	B	38	GLU
1	B	67	ASP
1	B	71	TRP
1	B	150	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	142	GLN
1	A	162	GLN
1	A	187	HIS
1	B	116	HIS
1	B	142	GLN
1	B	162	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	A	300	-	4,4,4	3.89	3 (75%)	6,6,6	1.05	0
2	PO4	B	300	-	4,4,4	4.00	3 (75%)	6,6,6	0.89	0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	300	PO4	P-O1	7.08	1.67	1.50
2	A	300	PO4	P-O1	6.78	1.66	1.50
2	A	300	PO4	P-O2	2.88	1.63	1.54
2	B	300	PO4	P-O2	2.55	1.62	1.54
2	B	300	PO4	P-O4	2.53	1.62	1.54
2	A	300	PO4	P-O4	2.37	1.61	1.54

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	172/202 (85%)	-1.40	0 100 100	24, 35, 58, 73	0
1	B	171/202 (84%)	-1.40	0 100 100	23, 35, 58, 71	0
All	All	343/404 (84%)	-1.40	0 100 100	23, 35, 58, 73	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	A	300	5/5	1.00	0.03	28,31,34,34	0
2	PO4	B	300	5/5	1.00	0.03	26,29,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.