

Full wwPDB X-ray Structure Validation Report (i)

Apr 29, 2025 – 01:28 PM EDT

PDB ID	:	$3\mathrm{UQZ} \ / \ \mathrm{pdb}_00003\mathrm{uqz}$
Title	:	X-ray structure of DNA processing protein A (DprA) from Streptococcus pneu-
		moniae
Authors	:	Quevillon-Cheruel, S.; Brooks, M.A.; Li de la Sierra-Gallay, I.
Deposited on	:	2011-11-21
Resolution	:	2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.43.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R _{free}	164625	3333 (2.70-2.70)
Clashscore	180529	3684 (2.70-2.70)
Ramachandran outliers	177936	3633 (2.70-2.70)
Sidechain outliers	177891	3633 (2.70-2.70)
RSRZ outliers	164620	3333 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	288	2% 8 5%	11%	••
1	В	288	80%	15%	••
1	С	288	9% 82%	12%	•••



 $\mathbf{2}$

Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 6732 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	1 A	283	Total	С	Ν	Ο	S	Se	0	0	0
1		203	2194	1401	370	412	6	5	0		
1	Р	280	Total	С	Ν	0	S	Se	0	0	0
1	ГБ		2164	1381	365	407	6	5	0		
1	1 C	000	Total	С	Ν	0	S	Se	0	0	0
	202	2184	1395	367	411	6	5	0	U	0	

• Molecule 1 is a protein called DNA processing protein DprA.

Chain	Residue	Modelled	Actual	Comment	Reference
А	283	HIS	-	expression tag	UNP Q97QF0
А	284	HIS	-	expression tag	UNP Q97QF0
А	285	HIS	-	expression tag	UNP Q97QF0
А	286	HIS	-	expression tag	UNP Q97QF0
А	287	HIS	-	expression tag	UNP Q97QF0
А	288	HIS	-	expression tag	UNP Q97QF0
В	283	HIS	-	expression tag	UNP Q97QF0
В	284	HIS	-	expression tag	UNP Q97QF0
В	285	HIS	-	expression tag	UNP Q97QF0
В	286	HIS	-	expression tag	UNP Q97QF0
В	287	HIS	-	expression tag	UNP Q97QF0
В	288	HIS	-	expression tag	UNP Q97QF0
С	283	HIS	-	expression tag	UNP Q97QF0
С	284	HIS	-	expression tag	UNP Q97QF0
С	285	HIS	-	expression tag	UNP Q97QF0
С	286	HIS	-	expression tag	UNP Q97QF0
С	287	HIS	-	expression tag	UNP Q97QF0
С	288	HIS	-	expression tag	UNP Q97QF0

There are 18 discrepancies between the modelled and reference sequences:

• Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
2	Δ	1	Total O S	0	1	
2	Л	1	10 8 2	0	1	
9	Λ	1	Total O S	O S O		
	Z A	1	5 4 1	0	0	
0	В	1	Total O S	0 0	0	
	D	1	5 4 1	0	0	
9	С	1	Total O S	0	0	
	U		5 4 1	0	0	

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	99	Total O 99 99	0	0
3	В	44	Total O 44 44	0	0
3	С	22	TotalO2222	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: DNA processing protein DprA









4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants	300.24Å 300.24Å 78.34Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
$\mathbf{Posolution} \left(\overset{\circ}{\mathbf{A}} \right)$	41.76 - 2.70	Depositor
Resolution (A)	41.76 - 2.70	EDS
% Data completeness	(Not available) $(41.76-2.70)$	Depositor
(in resolution range)	96.2(41.76-2.70)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.19 (at 2.69 \text{\AA})$	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
P. P.	0.219 , 0.245	Depositor
Π, Π_{free}	0.224 , 0.256	DCC
R_{free} test set	2420 reflections $(5.11%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	51.5	Xtriage
Anisotropy	0.166	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.34 , 47.3	EDS
L-test for $twinning^2$	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6732	wwPDB-VP
Average B, all atoms $(Å^2)$	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
	Unain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.76	0/2231	1.39	13/3003~(0.4%)	
1	В	0.83	1/2199~(0.0%)	1.38	13/2960~(0.4%)	
1	С	0.78	0/2220	1.39	11/2988~(0.4%)	
All	All	0.79	1/6650~(0.0%)	1.39	$37/8951 \ (0.4\%)$	

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	В	53	MSE	SE-CE	-8.25	1.70	1.95

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	226	MSE	N-CA-C	7.00	121.27	112.87
1	А	207	ALA	CA-C-N	6.43	128.90	120.28
1	А	207	ALA	C-N-CA	6.43	128.90	120.28
1	С	207	ALA	CA-C-N	6.43	128.90	120.28
1	С	207	ALA	C-N-CA	6.43	128.90	120.28
1	В	253	ASP	N-CA-C	6.35	119.19	111.82
1	А	253	ASP	N-CA-C	6.29	118.14	111.28
1	В	165	ILE	N-CA-C	6.28	118.79	108.99
1	В	164	VAL	CB-CA-C	6.17	118.74	110.42
1	В	207	ALA	CA-C-N	5.96	128.26	120.28
1	В	207	ALA	C-N-CA	5.96	128.26	120.28
1	В	82	PRO	CA-C-N	5.72	128.26	120.54
1	В	82	PRO	C-N-CA	5.72	128.26	120.54
1	С	164	VAL	CB-CA-C	5.72	118.81	110.98
1	С	244	VAL	N-CA-CB	5.49	118.78	111.64
1	А	42	ASP	CA-CB-CG	5.47	118.07	112.60
1	А	244	VAL	N-CA-CB	5.37	118.62	111.64

All (37) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	С	171	VAL	CB-CA-C	5.33	118.57	110.62
1	А	244	VAL	CB-CA-C	5.33	118.34	110.77
1	А	252	LEU	CA-C-N	5.31	127.40	120.28
1	А	252	LEU	C-N-CA	5.31	127.40	120.28
1	В	66	LYS	CA-C-N	5.30	127.33	120.44
1	В	66	LYS	C-N-CA	5.30	127.33	120.44
1	А	282	PHE	CA-C-N	5.28	131.20	121.70
1	А	282	PHE	C-N-CA	5.28	131.20	121.70
1	В	120	GLN	CA-C-N	5.27	125.95	119.99
1	В	120	GLN	C-N-CA	5.27	125.95	119.99
1	А	253	ASP	CA-CB-CG	5.25	117.85	112.60
1	А	66	LYS	CA-C-N	5.16	127.15	120.44
1	А	66	LYS	C-N-CA	5.16	127.15	120.44
1	С	253	ASP	CA-CB-CG	5.09	117.69	112.60
1	В	216	CYS	N-CA-C	5.08	118.01	110.14
1	С	244	VAL	CB-CA-C	5.07	117.97	110.77
1	С	66	LYS	CA-C-N	5.03	126.98	120.44
1	С	66	LYS	C-N-CA	5.03	126.98	120.44
1	С	253	ASP	N-CA-C	5.01	117.63	111.82
1	С	164	VAL	N-CA-CB	5.00	117.06	111.21

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2194	0	2205	15	0
1	В	2164	0	2183	17	0
1	С	2184	0	2198	15	0
2	А	15	0	0	0	0
2	В	5	0	0	0	0
2	С	5	0	0	0	0
3	А	99	0	0	0	0
3	В	44	0	0	1	0
3	С	22	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	6732	0	6586	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:C:226:MSE:HE3	1:C:258:GLY:HA2	1.67	0.76
1:A:164:VAL:O	1:A:208:ARG:HD3	2.00	0.61
1:C:142:LEU:H	1:C:151:HIS:HE1	1.50	0.60
1:A:142:LEU:H	1:A:151:HIS:HE1	1.50	0.60
1:B:137:VAL:HG22	1:B:160:LYS:HG3	1.85	0.59
1:B:142:LEU:H	1:B:151:HIS:HE1	1.50	0.58
1:C:137:VAL:HG22	1:C:160:LYS:HG3	1.85	0.57
1:A:151:HIS:HD2	1:A:161:THR:OG1	1.88	0.56
1:B:175:LYS:O	1:B:178:LYS:HG3	2.06	0.56
1:A:282:PHE:O	1:A:283:HIS:HB2	2.06	0.55
1:B:184:ILE:HG23	1:B:188:HIS:HB2	1.89	0.55
1:A:114:SER:HB2	1:A:223:GLU:O	2.08	0.53
1:B:63:HIS:O	1:B:63:HIS:ND1	2.43	0.52
1:A:164:VAL:HG12	1:A:208:ARG:HG2	1.91	0.52
1:A:226:MSE:HE1	1:A:261:HIS:HB2	1.91	0.52
1:C:217:ARG:HD3	1:C:281:GLU:HB3	1.93	0.51
1:C:164:VAL:HG22	1:C:208:ARG:HG2	1.94	0.50
1:B:165:ILE:HG12	3:B:422:HOH:O	2.12	0.49
1:C:114:SER:HB2	1:C:223:GLU:O	2.13	0.49
1:A:22:LEU:O	1:A:26:GLU:HG3	2.12	0.49
1:A:115:ARG:HH22	1:A:208:ARG:HH22	1.60	0.48
1:B:174:PRO:HG2	1:B:177:ASN:HD22	1.78	0.47
1:B:114:SER:HB2	1:B:223:GLU:O	2.15	0.46
1:B:115:ARG:HH22	1:B:208:ARG:HH22	1.64	0.46
1:B:143:ALA:HA	1:B:174:PRO:HD3	1.98	0.45
1:B:114:SER:N	1:B:146:ILE:HD11	2.33	0.44
1:C:115:ARG:HH22	1:C:208:ARG:HH22	1.65	0.44
1:A:36:LEU:HD21	1:C:51:VAL:HG21	1.99	0.44
1:A:115:ARG:HH22	1:A:208:ARG:NH2	2.16	0.43
1:A:222:ALA:HB2	1:A:247:ILE:HD12	2.00	0.42
1:A:283:HIS:CD2	1:C:176:ALA:HB2	2.54	0.42
1:C:255:LEU:H	1:C:255:LEU:HG	1.71	0.42
1:C:125:VAL:HG21	1:C:146:ILE:HG23	2.02	0.42



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:222:ALA:HB2	1:C:247:ILE:HD12	2.01	0.42
1:B:115:ARG:HA	1:B:144:LYS:O	2.20	0.42
1:B:151:HIS:HD2	1:B:161:THR:OG1	2.03	0.42
1:B:115:ARG:HH22	1:B:208:ARG:NH2	2.17	0.41
1:C:144:LYS:HG3	1:C:174:PRO:HG3	2.00	0.41
1:C:85:LEU:HD22	1:C:215:LEU:HD13	2.03	0.41
1:C:201:LEU:HD23	1:C:203:PHE:HE2	1.86	0.41
1:B:172:PHE:CB	1:B:178:LYS:HG2	2.51	0.41
1:A:85:LEU:HD22	1:A:215:LEU:HD13	2.04	0.40
1:B:222:ALA:HB2	1:B:247:ILE:HD12	2.03	0.40
1:A:125:VAL:HG21	1:A:146:ILE:HG23	2.02	0.40
1:B:202:LYS:HD3	1:B:202:LYS:HA	1.76	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	281/288 (98%)	273~(97%)	8(3%)	0	100 100
1	В	278/288~(96%)	267~(96%)	11 (4%)	0	100 100
1	С	280/288~(97%)	276 (99%)	3 (1%)	1 (0%)	30 55
All	All	839/864~(97%)	816 (97%)	22 (3%)	1 (0%)	48 73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	187	ASP



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percer	ntiles
1	А	235/235~(100%)	218~(93%)	17 (7%)	12	30
1	В	232/235~(99%)	211 (91%)	21 (9%)	7	19
1	С	234/235~(100%)	212 (91%)	22 (9%)	7	18
All	All	701/705~(99%)	641 (91%)	60 (9%)	8	21

All (60) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	16	LEU
1	А	51	VAL
1	А	79	ASP
1	А	93	VAL
1	А	135	GLU
1	А	142	LEU
1	А	156	GLN
1	А	208	ARG
1	А	210	ARG
1	А	215	LEU
1	А	221	VAL
1	А	230	SER
1	А	238	MSE
1	А	244	VAL
1	А	274	GLN
1	А	279	GLU
1	А	283	HIS
1	В	16	LEU
1	В	54	GLU
1	В	55	ARG
1	В	93	VAL
1	В	117	CYS
1	В	123	LYS
1	В	126	GLU
1	В	135	GLU
1	В	142	LEU



Mol	Chain	Res	Type
1	В	146	ILE
1	В	164	VAL
1	В	189	LEU
1	В	202	LYS
1	В	208	ARG
1	В	210	ARG
1	В	215	LEU
1	В	221	VAL
1	В	230	SER
1	В	238	MSE
1	В	268	LYS
1	В	274	GLN
1	С	16	LEU
1	С	54	GLU
1	С	79	ASP
1	С	93	VAL
1	С	119	LYS
1	С	135	GLU
1	С	142	LEU
1	С	164	VAL
1	С	171	VAL
1	С	186	ASN
1	С	202	LYS
1	С	208	ARG
1	С	210	ARG
1	С	215	LEU
1	С	221	VAL
1	С	226	MSE
1	С	230	SER
1	С	238	MSE
1	С	244	VAL
1	С	255	LEU
1	С	274	GLN
1	С	281	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	100	ASN
1	А	151	HIS
1	А	199	GLN
1	В	48	ASN



001000	iraca jien	e proces	o uo puge
\mathbf{Mol}	Chain	Res	Type
1	В	151	HIS
1	В	177	ASN
1	В	188	HIS
1	В	199	GLN
1	С	19	GLN
1	С	100	ASN
1	С	151	HIS
1	С	157	ASN
1	С	199	GLN

Continued from previous page...

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	301[A]	-	4,4,4	0.33	0	6,6,6	0.09	0
2	SO4	А	301[B]	-	4,4,4	0.35	0	6,6,6	0.18	0
2	SO4	А	302	-	4,4,4	0.31	0	6,6,6	0.13	0
2	SO4	С	300	-	4,4,4	0.33	0	6,6,6	0.11	0
2	SO4	В	300	-	4,4,4	0.30	0	6,6,6	0.19	0



There are no bond length outliers. There are no bond angle outliers. There are no chirality outliers. There are no torsion outliers. There are no ring outliers. No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	278/288~(96%)	-0.07	7 (2%) 58 57	26, 43, 65, 91	0
1	В	275/288 (95%)	1.23	75 (27%) 2 2	34, 59, 86, 105	0
1	С	277/288 (96%)	0.87	25 (9%) 17 15	46, 67, 89, 109	0
All	All	830/864~(96%)	0.68	107 (12%) 9 8	26, 57, 85, 109	0

All (107) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	119	LYS	5.8
1	В	174	PRO	5.6
1	С	282	PHE	5.4
1	В	177	ASN	5.2
1	В	155	LEU	4.9
1	В	158	GLY	4.9
1	В	159	GLY	4.9
1	В	183	TYR	4.4
1	В	184	ILE	4.4
1	В	185	GLY	4.3
1	В	106	PHE	4.2
1	В	175	LYS	4.1
1	В	98	LYS	4.0
1	В	173	TYR	3.9
1	С	119	LYS	3.9
1	В	154	ALA	3.9
1	С	236	ARG	3.9
1	В	280	PHE	3.8
1	В	180	LEU	3.8
1	В	202	LYS	3.8
1	В	156	GLN	3.7
1	В	199	GLN	3.7
1	В	146	ILE	3.7



Mol	Iol Chain R		Type	RSRZ	
1	В	151	HIS	3.7	
1	В	143	ALA	3.6	
1	А	283	HIS	3.6	
1	В	188	HIS	3.6	
1	В	133	GLU	3.5	
1	В	66	LYS	3.4	
1	В	117	CYS	3.4	
1	В	134	ASN	3.4	
1	В	172	PHE	3.4	
1	В	120	GLN	3.3	
1	С	227	ARG	3.3	
1	С	29	GLU	3.2	
1	В	176	ALA	3.2	
1	С	225	LYS	3.2	
1	А	48	ASN	3.2	
1	В	189	LEU	3.1	
1	В	217	ARG	3.1	
1	В	171	VAL	3.1	
1	В	190	VAL	3.0	
1	В	132	LEU	3.0	
1	С	57	PHE	3.0	
1	А	30	ASN	3.0	
1	С	83	TRP	2.9	
1	В	83	TRP	2.8	
1	В	187	ASP	2.8	
1	В	200	PRO	2.8	
1	С	31	VAL	2.8	
1	В	97	TYR	2.8	
1	В	147	ASP	2.8	
1	В	142	LEU	2.8	
1	В	196	PRO	2.7	
1	В	115	ARG	2.7	
1	C	89	TYR	2.7	
1	В	63	HIS	2.7	
1	С	2	LYS	2.6	
1	В	148	THR	2.6	
1	В	116	ALA	2.6	
1	С	107	PRO	2.5	
1	В	179	ARG	2.5	
1	С	123	LYS	2.5	
1	В	255	LEU	2.5	
1	С	115	ARG	2.5	



Mol	Chain	Res	Type	RSRZ
1	В	60	ASP	2.5
1	С	241	GLY	2.5
1	В	161	THR	2.4
1	С	34	GLU	2.4
1	В	104	LEU	2.4
1	С	35	LEU	2.4
1	В	122	ALA	2.4
1	С	116	ALA	2.3
1	В	101	LEU	2.3
1	В	181	GLN	2.3
1	В	121	GLY	2.3
1	В	259	CYS	2.3
1	С	158	GLY	2.3
1	А	252	LEU	2.3
1	А	281	GLU	2.2
1	В	198	GLU	2.2
1	С	33	GLN	2.2
1	В	170	ASP	2.2
1	В	150	ALA	2.2
1	В	113	GLY	2.2
1	В	197	GLY	2.2
1	В	178	LYS	2.2
1	В	225	LYS	2.2
1	В	30	ASN	2.2
1	А	239	GLU	2.1
1	С	147	ASP	2.1
1	В	203	PHE	2.1
1	В	103	LEU	2.1
1	C	48	ASN	2.1
1	В	127	LYS	2.1
1	С	127	LYS	2.1
1	В	253	ASP	2.1
1	В	12	LYS	2.1
1	В	144	LYS	2.1
1	В	160	LYS	2.1
1	В	167	THR	2.1
1	A	253	ASP	2.0
1	С	27	TYR	2.0
1	B	99	GLY	2.0
1	В	201	LEU	2.0
1	С	280	PHE	2.0
1	В	157	ASN	2.0

Continued from previous page...



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
2	SO4	В	300	5/5	0.78	0.17	116,120,120,122	0
2	SO4	А	301[B]	5/5	0.85	0.30	35,39,40,41	5
2	SO4	А	301[A]	5/5	0.85	0.30	64,68,69,69	5
2	SO4	С	300	5/5	0.85	0.14	98,102,103,103	0
2	SO4	А	302	5/5	0.97	0.07	$50,\!53,\!55,\!57$	0

6.5 Other polymers (i)

There are no such residues in this entry.

