



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 19, 2024 – 03:52 AM EDT

PDB ID : 3ZJE
Title : A20 OTU domain in reversibly oxidised (SOH) state
Authors : Kulathu, Y.; Garcia, F.J.; Mevissen, T.E.T.; Busch, M.; Arnaudo, N.; Carroll, K.S.; Barford, D.; Komander, D.
Deposited on : 2013-01-17
Resolution : 1.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4003 (1.86-1.82)
Clashscore	141614	4233 (1.86-1.82)
Ramachandran outliers	138981	4185 (1.86-1.82)
Sidechain outliers	138945	4186 (1.86-1.82)
RSRZ outliers	127900	3957 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	 14% 76% 11% 10%
1	B	366	 11% 74% 9% 5% 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EDO	A	1358	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5697 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called A20P50.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	329	2694	1732	468	477	17	28	8	0
1	B	324	2664	1718	460	470	16	54	3	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	114	SER	GLY	engineered mutation	UNP P21580
B	114	SER	GLY	engineered mutation	UNP P21580

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	4	2	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Cl	0	0
			2	2		
3	B	4	Total	Cl	0	0
			4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	169	Total	O	0	0
			169	169		
4	B	156	Total	O	0	0
			156	156		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: A20P50



- Molecule 1: A20P50



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	43.85Å 69.15Å 84.70Å 99.14° 100.28° 97.09°	Depositor
Resolution (Å)	33.72 – 1.84 33.72 – 1.84	Depositor EDS
% Data completeness (in resolution range)	94.3 (33.72-1.84) 94.3 (33.72-1.84)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 1.84Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.182 , 0.220 0.183 , 0.219	Depositor DCC
R_{free} test set	3937 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	27.3	Xtrriage
Anisotropy	0.205	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 62.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5697	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSO, CL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.84	2/2768 (0.1%)	0.82	3/3751 (0.1%)
1	B	0.75	2/2723 (0.1%)	0.81	3/3688 (0.1%)
All	All	0.80	4/5491 (0.1%)	0.81	6/7439 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	86[A]	CYS	CB-SG	-6.86	1.70	1.82
1	A	86[B]	CYS	CB-SG	-6.86	1.70	1.82
1	B	342	VAL	CB-CG2	-5.61	1.41	1.52
1	B	229	VAL	CB-CG1	-5.02	1.42	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	341	LEU	CB-CG-CD2	-9.87	94.22	111.00
1	B	48	LEU	CB-CG-CD1	-7.66	97.98	111.00
1	A	318	ASP	CB-CG-OD1	5.64	123.38	118.30
1	A	48	LEU	CA-CB-CG	5.27	127.42	115.30
1	B	278	ARG	NE-CZ-NH2	-5.18	117.71	120.30
1	A	147	LEU	CA-CB-CG	5.07	126.96	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2694	0	2650	38	0
1	B	2664	0	2619	39	0
2	A	8	0	12	7	0
3	A	2	0	0	2	0
3	B	4	0	0	0	0
4	A	169	0	0	5	0
4	B	156	0	0	2	0
All	All	5697	0	5281	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:45:ARG:HE	2:A:1358:EDO:H11	1.10	1.08
1:B:50:MET:HE2	1:B:51:PHE:H	1.31	0.95
1:B:48:LEU:HD11	1:B:342:VAL:HG21	1.49	0.92
1:B:57:CYS:HG	1:B:353:TYR:HH	1.10	0.91
1:B:66:LYS:O	1:B:96:LYS:NZ	2.04	0.91
1:A:45:ARG:NE	2:A:1358:EDO:H11	1.90	0.86
1:B:52:ARG:HH21	1:B:54:CYS:HB3	1.42	0.81
1:B:31:LYS:H	1:B:31:LYS:HE2	1.45	0.81
1:B:52:ARG:HH11	1:B:263:LEU:HD21	1.46	0.81
1:B:11:TYR:HA	1:B:341:LEU:HD21	1.66	0.77
1:B:50:MET:HE2	1:B:51:PHE:N	2.03	0.73
1:A:48:LEU:HD22	1:A:261[B]:VAL:HG11	1.71	0.73
3:A:1359:CL:CL	4:A:2104:HOH:O	2.43	0.72
1:A:277:ASN:OD1	4:A:2053:HOH:O	2.12	0.68
1:B:336:PRO:O	1:B:340:ASN:HB2	1.96	0.66
1:A:69:ILE:HD12	1:A:93[B]:VAL:CG2	2.27	0.65
1:B:32:PRO:HG2	1:B:36:ILE:HG12	1.79	0.65
1:B:49:GLU:HG2	1:B:262:THR:HG22	1.81	0.63
1:B:48:LEU:CD1	1:B:342:VAL:HG21	2.26	0.63
1:A:52:ARG:HA	1:A:52:ARG:HE	1.65	0.60
1:B:21:ILE:HG12	1:B:339:ILE:HB	1.84	0.60
1:A:49:GLU:HB2	1:A:333:ALA:HB2	1.85	0.59
1:A:82:LYS:HE2	1:A:213:LYS:HB3	1.84	0.59
1:B:227:LEU:HB3	1:B:229:VAL:HG23	1.85	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:50:MET:HE3	1:B:50:MET:HA	1.85	0.57
1:B:53:THR:OG1	1:B:65:HIS:NE2	2.18	0.57
1:B:34:ASN:OD1	1:B:36:ILE:HG23	2.05	0.56
1:A:45:ARG:HE	2:A:1358:EDO:C1	2.02	0.56
1:B:319:HIS:HE1	4:B:2066:HOH:O	1.89	0.55
1:B:50:MET:HA	1:B:50:MET:CE	2.36	0.55
1:A:45:ARG:HH21	2:A:1358:EDO:H22	1.72	0.54
1:B:10:LEU:CD2	1:B:342:VAL:HG22	2.37	0.54
1:B:335:LEU:HD22	1:B:339:ILE:HG13	1.89	0.54
1:A:22:ARG:NH2	4:A:2005:HOH:O	2.33	0.53
1:A:227:LEU:HD13	3:A:1360:CL:CL	2.46	0.52
1:A:80:GLN:NE2	1:A:212:ASP:OD1	2.36	0.52
1:B:52:ARG:NH1	1:B:263:LEU:HD21	2.22	0.52
1:A:53:THR:HG21	1:A:65:HIS:CE1	2.45	0.51
1:A:69:ILE:HD12	1:A:93[B]:VAL:HG21	1.93	0.50
1:A:53:THR:HG21	1:A:65:HIS:NE2	2.26	0.49
1:A:86[B]:CYS:HB3	1:A:89:VAL:HG22	1.93	0.49
1:A:17:LYS:NZ	1:A:344:ASP:OD2	2.42	0.48
1:B:133:THR:HG22	1:B:134:ASP:H	1.78	0.48
1:A:87:ARG:NH2	1:A:315:GLN:OE1	2.45	0.48
1:A:53:THR:HG23	1:A:61:ARG:HG2	1.96	0.47
1:A:69:ILE:HD12	1:A:93[B]:VAL:HG22	1.97	0.47
1:B:263:LEU:HB3	4:B:2063:HOH:O	2.15	0.47
1:A:52:ARG:NH1	1:A:54:CYS:HB3	2.30	0.46
1:B:10:LEU:HG	1:B:342:VAL:HG22	1.97	0.46
1:A:213:LYS:HB2	1:A:227:LEU:N	2.30	0.46
1:A:121:VAL:HB	2:A:1357:EDO:C1	2.45	0.46
1:A:187:GLN:HG2	1:A:188:TYR:CE2	2.50	0.46
1:B:52:ARG:NH2	1:B:54:CYS:HB3	2.23	0.45
1:B:133:THR:HG22	1:B:134:ASP:N	2.32	0.45
1:B:186:LEU:HD22	1:B:186:LEU:H	1.82	0.45
1:A:299:LYS:HB3	1:A:299:LYS:HE2	1.63	0.45
1:A:121:VAL:HB	2:A:1357:EDO:H11	1.98	0.45
1:A:325:ILE:O	1:A:325:ILE:HG13	2.17	0.45
1:A:334:ASN:O	1:A:336:PRO:HD3	2.17	0.44
1:A:60:PHE:HE2	1:A:349:VAL:HG13	1.81	0.44
1:A:24:ARG:HD2	4:A:2008:HOH:O	2.17	0.44
1:A:48:LEU:HD23	1:A:49:GLU:N	2.32	0.44
1:A:95:LEU:HB2	1:A:259:PRO:HG2	2.01	0.43
1:B:95:LEU:HB2	1:B:259:PRO:HG2	2.00	0.43
1:A:45:ARG:HH21	2:A:1358:EDO:C2	2.31	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:287:LYS:HE2	4:A:2023:HOH:O	2.18	0.42
1:B:52:ARG:HH21	1:B:54:CYS:CB	2.23	0.42
1:B:31:LYS:H	1:B:31:LYS:CE	2.22	0.42
1:B:335:LEU:HA	1:B:336:PRO:HD3	1.88	0.42
1:B:287:LYS:HA	1:B:287:LYS:HD2	1.74	0.42
1:B:170:ASN:O	1:B:174[B]:MET:HG2	2.19	0.42
1:B:52:ARG:HD2	1:B:52:ARG:HA	1.69	0.41
1:A:292:THR:OG1	1:A:294:PRO:HD2	2.20	0.41
1:A:284:GLU:OE1	1:A:284:GLU:HA	2.21	0.41
1:B:119:ASP:O	1:B:120:LEU:HB2	2.20	0.41
1:B:19:VAL:O	1:B:23:GLU:HG3	2.21	0.41
1:A:134:ASP:OD1	1:A:136:ARG:HD3	2.22	0.40
1:B:200:CYS:HB2	1:B:205:ARG:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/366 (89%)	316 (97%)	10 (3%)	0	100	100
1	B	316/366 (86%)	305 (96%)	9 (3%)	2 (1%)	25	12
All	All	642/732 (88%)	621 (97%)	19 (3%)	2 (0%)	41	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	133	THR
1	B	34	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	290/335 (87%)	274 (94%)	16 (6%)	21	7
1	B	286/335 (85%)	266 (93%)	20 (7%)	15	3
All	All	576/670 (86%)	540 (94%)	36 (6%)	17	4

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	LEU
1	A	38	HIS
1	A	52	ARG
1	A	53	THR
1	A	87	ARG
1	A	90	ARG
1	A	134	ASP
1	A	136	ARG
1	A	147	LEU
1	A	213	LYS
1	A	227	LEU
1	A	241	GLN
1	A	284	GLU
1	A	335	LEU
1	A	343	ASP
1	A	348	LEU
1	B	6	LEU
1	B	10	LEU
1	B	31	LYS
1	B	36	ILE
1	B	38	HIS
1	B	50	MET
1	B	52	ARG
1	B	120	LEU
1	B	132	GLU
1	B	134	ASP
1	B	164	TRP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	173	LYS
1	B	186	LEU
1	B	262	THR
1	B	282	ARG
1	B	287	LYS
1	B	319	HIS
1	B	339	ILE
1	B	341	LEU
1	B	347	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSO	B	103	1	3,6,7	0.96	0	1,6,8	0.31	0
1	CSO	A	103	1	3,6,7	0.93	0	1,6,8	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	B	103	1	-	0/1/5/7	-
1	CSO	A	103	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	1357	-	3,3,3	0.31	0	2,2,2	0.64	0
2	EDO	A	1358	-	3,3,3	0.35	0	2,2,2	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	1357	-	-	0/1/1/1	-
2	EDO	A	1358	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1357	EDO	2	0
2	A	1358	EDO	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	328/366 (89%)	0.61	52 (15%) 1 1	11, 32, 66, 78	9 (2%)
1	B	323/366 (88%)	0.54	42 (13%) 3 2	13, 34, 71, 91	15 (4%)
All	All	651/732 (88%)	0.57	94 (14%) 2 2	11, 33, 68, 91	24 (3%)

All (94) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	57	CYS	9.2
1	B	58	PRO	7.6
1	B	356	TRP	6.9
1	A	58	PRO	6.4
1	B	60	PHE	6.2
1	B	54	CYS	6.0
1	A	161	THR	5.8
1	A	33	THR	5.6
1	B	33	THR	5.5
1	A	179	THR	5.0
1	A	321	THR	5.0
1	A	54	CYS	4.9
1	B	36	ILE	4.8
1	B	63	ILE	4.8
1	A	185	GLY	4.6
1	B	134	ASP	4.5
1	B	56	PHE	4.4
1	A	59	GLN	4.3
1	A	180	PRO	4.3
1	A	160	ASP	4.3
1	A	63	ILE	4.3
1	B	152	PHE	4.2
1	A	337	LYS	4.0
1	A	248	ILE	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	153	VAL	4.0
1	A	57	CYS	4.0
1	A	162	ARG	4.0
1	A	32	PRO	4.0
1	A	207	ILE	3.9
1	A	227	LEU	3.8
1	A	56	PHE	3.7
1	B	150	GLN	3.7
1	B	248	ILE	3.6
1	B	34	ASN	3.5
1	B	179	THR	3.5
1	A	60	PHE	3.4
1	B	164	TRP	3.4
1	A	268	PRO	3.3
1	A	8	GLN	3.3
1	A	6	LEU	3.3
1	A	319	HIS	3.3
1	B	166	ASP	3.2
1	B	32	PRO	3.2
1	A	267	GLY	3.2
1	A	249	VAL	3.2
1	A	199	LEU	3.1
1	B	249	VAL	3.1
1	B	207	ILE	3.0
1	B	55	GLN	3.0
1	A	214	MET	3.0
1	A	196	ILE	2.9
1	A	203	LEU	2.9
1	B	111	TYR	2.9
1	A	163	ASN	2.9
1	B	186	LEU	2.8
1	B	180	PRO	2.8
1	B	250	LEU	2.8
1	B	59	GLN	2.8
1	B	185	GLY	2.7
1	B	261	VAL	2.7
1	A	186	LEU	2.7
1	A	322	THR	2.7
1	B	208	ILE	2.7
1	A	250	LEU	2.7
1	A	34	ASN	2.7
1	A	208	ILE	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	30	PHE	2.6
1	B	337	LYS	2.6
1	B	353	TYR	2.6
1	B	151	GLU	2.6
1	A	232	ILE	2.6
1	A	218	LEU	2.5
1	B	209	VAL	2.5
1	B	6	LEU	2.5
1	A	233	TYR	2.5
1	A	107	ALA	2.4
1	A	323	HIS	2.4
1	A	104	LEU	2.4
1	B	35	GLY	2.4
1	B	267	GLY	2.4
1	A	62	GLU	2.4
1	A	108	THR	2.4
1	B	108	THR	2.4
1	B	133	THR	2.4
1	A	353	TYR	2.3
1	B	357	GLN	2.3
1	B	31	LYS	2.3
1	A	122	LEU	2.2
1	A	178	ASP	2.2
1	A	200	CYS	2.1
1	A	11	TYR	2.1
1	B	149	SER	2.1
1	A	55	GLN	2.0
1	A	320	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSO	B	103	7/8	0.97	0.11	23,23,34,43	0
1	CSO	A	103	7/8	0.98	0.11	15,18,29,42	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	EDO	A	1358	4/4	0.93	0.18	18,26,31,40	0
3	CL	A	1359	1/1	0.97	0.06	29,29,29,29	0
3	CL	A	1360	1/1	0.97	0.09	49,49,49,49	0
2	EDO	A	1357	4/4	0.99	0.10	22,25,26,28	0
3	CL	B	1358	1/1	0.99	0.09	18,18,18,18	0
3	CL	B	1361	1/1	0.99	0.06	36,36,36,36	0
3	CL	B	1360	1/1	1.00	0.04	22,22,22,22	0
3	CL	B	1359	1/1	1.00	0.07	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.