



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 28, 2024 – 05:50 am BST

PDB ID : 4BA9
Title : The structural basis for the coordination of Y-family Translesion DNA Polymerases by Rev1
Authors : Grummitt, C.G.; Kilkenny, M.L.; Frey, A.; Roe, S.M.; Oliver, A.W.; Sale, J.E.; Pearl, L.H.
Deposited on : 2012-09-12
Resolution : 2.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1271 (2.76-2.72)
Clashscore	141614	1322 (2.76-2.72)
Ramachandran outliers	138981	1297 (2.76-2.72)
Sidechain outliers	138945	1298 (2.76-2.72)
RSRZ outliers	127900	1243 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	114	
1	B	114	
1	C	114	
1	D	114	
1	E	114	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	114	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a red segment on the left labeled '9%', a large green segment in the middle labeled '88%', and a yellow segment on the right labeled '9%'. The bar is set against a light gray background.</p>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5313 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA POLYMERASE KAPPA, DNA REPAIR PROTEIN REV1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	104	845	540	138	161	6	0	0	0
1	B	104	849	542	138	163	6	0	0	0
1	C	107	881	560	144	169	8	0	2	0
1	D	104	855	545	141	163	6	0	0	0
1	E	104	846	540	136	163	7	0	1	0
1	F	104	860	549	141	163	7	0	1	0

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	201	GLY	-	expression tag	UNP Q9UBT6
A	202	PRO	-	expression tag	UNP Q9UBT6
A	203	HIS	-	expression tag	UNP Q9UBT6
A	204	MET	-	expression tag	UNP Q9UBT6
A	218	GLY	-	linker	UNP Q9UBZ9
A	219	SER	-	linker	UNP Q9UBZ9
A	220	ASP	-	linker	UNP Q9UBZ9
A	1149	ALA	-	linker	UNP Q9UBZ9
A	1150	GLY	-	linker	UNP Q9UBZ9
A	1151	SER	-	linker	UNP Q9UBZ9
A	1152	ALA	-	linker	UNP Q9UBZ9
A	1153	GLY	-	linker	UNP Q9UBZ9
A	1154	ASP	-	linker	UNP Q9UBZ9
A	1155	GLY	-	linker	UNP Q9UBZ9
A	1156	ALA	-	linker	UNP Q9UBZ9
A	1157	SER	-	linker	UNP Q9UBZ9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	201	GLY	-	expression tag	UNP Q9UBT6
B	202	PRO	-	expression tag	UNP Q9UBT6
B	203	HIS	-	expression tag	UNP Q9UBT6
B	204	MET	-	expression tag	UNP Q9UBT6
B	218	GLY	-	linker	UNP Q9UBZ9
B	219	SER	-	linker	UNP Q9UBZ9
B	220	ASP	-	linker	UNP Q9UBZ9
B	1149	ALA	-	linker	UNP Q9UBZ9
B	1150	GLY	-	linker	UNP Q9UBZ9
B	1151	SER	-	linker	UNP Q9UBZ9
B	1152	ALA	-	linker	UNP Q9UBZ9
B	1153	GLY	-	linker	UNP Q9UBZ9
B	1154	ASP	-	linker	UNP Q9UBZ9
B	1155	GLY	-	linker	UNP Q9UBZ9
B	1156	ALA	-	linker	UNP Q9UBZ9
B	1157	SER	-	linker	UNP Q9UBZ9
C	201	GLY	-	expression tag	UNP Q9UBT6
C	202	PRO	-	expression tag	UNP Q9UBT6
C	203	HIS	-	expression tag	UNP Q9UBT6
C	204	MET	-	expression tag	UNP Q9UBT6
C	218	GLY	-	linker	UNP Q9UBZ9
C	219	SER	-	linker	UNP Q9UBZ9
C	220	ASP	-	linker	UNP Q9UBZ9
C	1149	ALA	-	linker	UNP Q9UBZ9
C	1150	GLY	-	linker	UNP Q9UBZ9
C	1151	SER	-	linker	UNP Q9UBZ9
C	1152	ALA	-	linker	UNP Q9UBZ9
C	1153	GLY	-	linker	UNP Q9UBZ9
C	1154	ASP	-	linker	UNP Q9UBZ9
C	1155	GLY	-	linker	UNP Q9UBZ9
C	1156	ALA	-	linker	UNP Q9UBZ9
C	1157	SER	-	linker	UNP Q9UBZ9
D	201	GLY	-	expression tag	UNP Q9UBT6
D	202	PRO	-	expression tag	UNP Q9UBT6
D	203	HIS	-	expression tag	UNP Q9UBT6
D	204	MET	-	expression tag	UNP Q9UBT6
D	218	GLY	-	linker	UNP Q9UBZ9
D	219	SER	-	linker	UNP Q9UBZ9
D	220	ASP	-	linker	UNP Q9UBZ9
D	1149	ALA	-	linker	UNP Q9UBZ9
D	1150	GLY	-	linker	UNP Q9UBZ9
D	1151	SER	-	linker	UNP Q9UBZ9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	1152	ALA	-	linker	UNP Q9UBZ9
D	1153	GLY	-	linker	UNP Q9UBZ9
D	1154	ASP	-	linker	UNP Q9UBZ9
D	1155	GLY	-	linker	UNP Q9UBZ9
D	1156	ALA	-	linker	UNP Q9UBZ9
D	1157	SER	-	linker	UNP Q9UBZ9
E	201	GLY	-	expression tag	UNP Q9UBT6
E	202	PRO	-	expression tag	UNP Q9UBT6
E	203	HIS	-	expression tag	UNP Q9UBT6
E	204	MET	-	expression tag	UNP Q9UBT6
E	218	GLY	-	linker	UNP Q9UBZ9
E	219	SER	-	linker	UNP Q9UBZ9
E	220	ASP	-	linker	UNP Q9UBZ9
E	1149	ALA	-	linker	UNP Q9UBZ9
E	1150	GLY	-	linker	UNP Q9UBZ9
E	1151	SER	-	linker	UNP Q9UBZ9
E	1152	ALA	-	linker	UNP Q9UBZ9
E	1153	GLY	-	linker	UNP Q9UBZ9
E	1154	ASP	-	linker	UNP Q9UBZ9
E	1155	GLY	-	linker	UNP Q9UBZ9
E	1156	ALA	-	linker	UNP Q9UBZ9
E	1157	SER	-	linker	UNP Q9UBZ9
F	201	GLY	-	expression tag	UNP Q9UBT6
F	202	PRO	-	expression tag	UNP Q9UBT6
F	203	HIS	-	expression tag	UNP Q9UBT6
F	204	MET	-	expression tag	UNP Q9UBT6
F	218	GLY	-	linker	UNP Q9UBZ9
F	219	SER	-	linker	UNP Q9UBZ9
F	220	ASP	-	linker	UNP Q9UBZ9
F	1149	ALA	-	linker	UNP Q9UBZ9
F	1150	GLY	-	linker	UNP Q9UBZ9
F	1151	SER	-	linker	UNP Q9UBZ9
F	1152	ALA	-	linker	UNP Q9UBZ9
F	1153	GLY	-	linker	UNP Q9UBZ9
F	1154	ASP	-	linker	UNP Q9UBZ9
F	1155	GLY	-	linker	UNP Q9UBZ9
F	1156	ALA	-	linker	UNP Q9UBZ9
F	1157	SER	-	linker	UNP Q9UBZ9

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	B	2	Total Mg 2 2	0	0
2	C	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	E	2	Total Mg 2 2	0	0
2	F	1	Total Mg 1 1	0	0

- Molecule 3 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Ni 2 2	0	0
3	B	2	Total Ni 2 2	0	0
3	D	2	Total Ni 2 2	0	0
3	E	2	Total Ni 2 2	0	0
3	F	2	Total Ni 2 2	0	0


- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	29	Total O 29 29	0	0
4	B	47	Total O 47 47	0	0
4	C	21	Total O 21 21	0	0
4	D	34	Total O 34 34	0	0
4	E	12	Total O 12 12	0	0
4	F	16	Total O 16 16	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA POLYMERASE KAPPA, DNA REPAIR PROTEIN REV1

Chain A: 




- Molecule 1: DNA POLYMERASE KAPPA, DNA REPAIR PROTEIN REV1

Chain B: 





- Molecule 1: DNA POLYMERASE KAPPA, DNA REPAIR PROTEIN REV1

Chain C: 





- Molecule 1: DNA POLYMERASE KAPPA, DNA REPAIR PROTEIN REV1

Chain D: 





- Molecule 1: DNA POLYMERASE KAPPA, DNA REPAIR PROTEIN REV1

Chain E: 



- Molecule 1: DNA POLYMERASE KAPPA, DNA REPAIR PROTEIN REV1

Chain F:  9% 88% 9%



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	105.75Å 105.75Å 424.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.82 – 2.73 84.08 – 2.73	Depositor EDS
% Data completeness (in resolution range)	98.7 (42.82-2.73) 98.8 (84.08-2.73)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.05 (at 2.73Å)	Xtrriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.214 , 0.250 0.226 , 0.262	Depositor DCC
R_{free} test set	1256 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	59.8	Xtrriage
Anisotropy	0.297	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 62.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5313	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/858	0.58	0/1154
1	B	0.49	0/862	0.58	0/1159
1	C	0.58	0/900	0.68	0/1208
1	D	0.51	0/868	0.61	0/1166
1	E	0.48	0/862	0.58	0/1161
1	F	0.46	0/876	0.58	0/1176
All	All	0.50	0/5226	0.60	0/7024

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	845	0	838	2	0
1	B	849	0	842	9	0
1	C	881	0	879	11	0
1	D	855	0	853	4	0
1	E	846	0	829	5	0
1	F	860	0	862	2	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	2	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	2	0	0	0	0
2	F	1	0	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	D	2	0	0	0	0
3	E	2	0	0	0	0
3	F	2	0	0	0	0
4	A	29	0	0	0	0
4	B	47	0	0	4	0
4	C	21	0	0	1	0
4	D	34	0	0	0	0
4	E	12	0	0	0	0
4	F	16	0	0	0	0
All	All	5313	0	5103	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1194[B]:CYS:SG	1:C:1236:VAL:HG22	1.71	1.29
1:C:217:ARG:HB2	1:C:218:GLY:HA2	1.44	0.99
1:C:1194[B]:CYS:SG	1:C:1236:VAL:CG2	2.52	0.97
1:B:1242:GLN:OXT	1:E:1233:LEU:HD21	1.80	0.81
1:C:217:ARG:HB2	1:C:218:GLY:CA	2.13	0.79
1:B:1174:GLU:O	1:B:1178:THR:HB	1.93	0.69
1:A:1174:GLU:O	1:A:1178:THR:HB	1.95	0.67
1:E:1174:GLU:O	1:E:1178:THR:HB	1.95	0.66
1:D:1174:GLU:O	1:D:1178:THR:HB	1.96	0.65
1:F:1174:GLU:O	1:F:1178:THR:HB	1.98	0.62
1:B:1180:SER:HA	4:B:2016:HOH:O	2.03	0.58
1:B:1210:ILE:HD11	1:B:1236:VAL:HG11	1.88	0.54
1:C:1173:ARG:HG3	1:C:1212:TYR:OH	2.08	0.53
1:C:1160:ALA:HB3	1:C:1171:LEU:HD13	1.90	0.53
1:C:1178:THR:HB	1:C:1179:ILE:HD12	1.94	0.50
1:F:1210:ILE:HD11	1:F:1236:VAL:HG11	1.94	0.49
1:B:1221:VAL:HG22	4:B:2035:HOH:O	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1210:ILE:HD11	1:A:1236:VAL:HG11	1.96	0.48
1:B:1242:GLN:OXT	1:E:1233:LEU:CD2	2.58	0.47
1:C:1175:TRP:HD1	1:C:1179:ILE:HD13	1.79	0.47
1:B:1242:GLN:HA	1:E:1233:LEU:HD11	1.97	0.47
1:C:1194[B]:CYS:HG	1:C:1236:VAL:HG22	1.68	0.46
1:B:202:PRO:HB3	1:D:1235:ASN:OD1	2.15	0.46
1:D:1210:ILE:HD11	1:D:1236:VAL:HG11	1.97	0.46
4:B:2037:HOH:O	1:D:1221:VAL:HG12	2.16	0.46
1:E:1210:ILE:HD11	1:E:1236:VAL:HG11	1.98	0.45
1:C:1172:LEU:O	1:C:1176:ILE:HD12	2.20	0.42
1:B:1183:MET:HB3	4:B:2020:HOH:O	2.21	0.40
1:C:1164:GLU:HB3	4:C:2008:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	100/114 (88%)	99 (99%)	1 (1%)	0	100	100
1	B	100/114 (88%)	99 (99%)	1 (1%)	0	100	100
1	C	105/114 (92%)	100 (95%)	5 (5%)	0	100	100
1	D	100/114 (88%)	99 (99%)	1 (1%)	0	100	100
1	E	101/114 (89%)	100 (99%)	1 (1%)	0	100	100
1	F	101/114 (89%)	99 (98%)	2 (2%)	0	100	100
All	All	607/684 (89%)	596 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	95/102 (93%)	95 (100%)	0	100	100
1	B	96/102 (94%)	96 (100%)	0	100	100
1	C	101/102 (99%)	97 (96%)	4 (4%)	31	52
1	D	97/102 (95%)	96 (99%)	1 (1%)	76	85
1	E	95/102 (93%)	95 (100%)	0	100	100
1	F	98/102 (96%)	98 (100%)	0	100	100
All	All	582/612 (95%)	577 (99%)	5 (1%)	78	87

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	217	ARG
1	C	220	ASP
1	C	1169	LYS
1	C	1173	ARG
1	D	1173	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	1237	GLN
1	E	1241	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 18 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	104/114 (91%)	0.45	3 (2%) 51 58	38, 76, 100, 109	0
1	B	104/114 (91%)	0.33	3 (2%) 51 58	34, 56, 90, 107	0
1	C	107/114 (93%)	0.76	14 (13%) 3 3	41, 69, 91, 104	0
1	D	104/114 (91%)	0.29	1 (0%) 82 86	39, 53, 82, 105	0
1	E	104/114 (91%)	0.54	8 (7%) 13 15	42, 68, 94, 118	0
1	F	104/114 (91%)	0.72	10 (9%) 8 8	41, 80, 109, 114	0
All	All	627/684 (91%)	0.51	39 (6%) 20 23	34, 67, 99, 118	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	1241	GLN	5.6
1	F	1216	LEU	5.0
1	F	1217	MET	4.8
1	E	1221	VAL	4.5
1	F	1173	ARG	4.4
1	C	1209	VAL	4.1
1	C	1229	PHE	3.9
1	C	1161	GLY	3.8
1	E	1217	MET	3.5
1	C	1233	LEU	3.4
1	C	1163	VAL	3.3
1	F	1229	PHE	3.2
1	F	1176	ILE	3.1
1	E	1218	GLN	3.1
1	C	1162	ALA	3.0
1	C	1228	ALA	2.9
1	B	1241	GLN	2.7
1	F	1171	LEU	2.6
1	F	1159	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	1232	ILE	2.5
1	B	1190	VAL	2.4
1	C	1179	ILE	2.4
1	E	1209	VAL	2.3
1	E	1203	LEU	2.3
1	A	1209	VAL	2.3
1	E	1163	VAL	2.3
1	F	1175	TRP	2.3
1	C	218	GLY	2.2
1	C	219	SER	2.2
1	F	1206	LEU	2.2
1	E	1204	GLU	2.2
1	A	1168	VAL	2.2
1	A	1206	LEU	2.2
1	F	1172	LEU	2.1
1	C	1206	LEU	2.1
1	E	1165	PHE	2.1
1	B	1210	ILE	2.1
1	C	1194[A]	CYS	2.0
1	C	1208	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MG	E	2243	1/1	0.88	0.20	74,74,74,74	0
2	MG	D	2243	1/1	0.94	0.27	79,79,79,79	0
2	MG	E	2244	1/1	0.94	0.28	64,64,64,64	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MG	F	2243	1/1	0.95	0.20	61,61,61,61	0
2	MG	B	2244	1/1	0.96	0.27	57,57,57,57	0
2	MG	A	2243	1/1	0.97	0.17	67,67,67,67	0
2	MG	B	2243	1/1	0.98	0.17	59,59,59,59	0
2	MG	C	2243	1/1	0.98	0.21	63,63,63,63	0
3	NI	B	2245	1/1	0.99	0.23	47,47,47,47	0
3	NI	B	2246	1/1	0.99	0.17	44,44,44,44	0
3	NI	D	2245	1/1	0.99	0.22	52,52,52,52	0
3	NI	E	2245	1/1	0.99	0.22	53,53,53,53	0
3	NI	E	2246	1/1	0.99	0.14	37,37,37,37	1
3	NI	F	2244	1/1	0.99	0.19	59,59,59,59	0
3	NI	F	2245	1/1	0.99	0.16	40,40,40,40	1
3	NI	D	2244	1/1	1.00	0.20	57,57,57,57	0
3	NI	A	2244	1/1	1.00	0.24	48,48,48,48	0
3	NI	A	2245	1/1	1.00	0.16	57,57,57,57	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.