



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 29, 2024 – 08:11 pm BST

PDB ID : 4COI
Title : Crystal structure of the anaerobic ribonucleotide reductase from *Thermotoga maritima* with glycerol in the active site
Authors : Aurelius, O.; Johansson, R.; Bagenholm, V.; Beck, T.; Balhuizen, A.; Lundin, D.; Sjoberg, B.M.; Mulliez, E.; Logan, D.T.
Deposited on : 2014-01-28
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	651	 3% 88% 6% 7%
1	B	651	 3% 90% 6%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 10496 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ANAEROBIC RIBONUCLEOSIDE-TRIPHOSPHATE REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	606	Total 4969	C 3210	N 818	O 914	S 27	0	1	0
1	B	611	Total 5026	C 3245	N 829	O 925	S 27	0	4	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Zn 1	0	0
2	B	1	Total 1	Zn 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	272	Total O 272 272	0	0
4	B	197	Total O 197 197	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	77.89Å 98.82Å 86.98Å 90.00° 112.42° 90.00°	Depositor
Resolution (Å)	24.30 – 1.94 24.30 – 1.94	Depositor EDS
% Data completeness (in resolution range)	97.7 (24.30-1.94) 97.7 (24.30-1.94)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 1.95Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.181 , 0.217 0.199 , 0.235	Depositor DCC
R_{free} test set	2210 reflections (2.52%)	wwPDB-VP
Wilson B-factor (Å ²)	27.8	Xtrriage
Anisotropy	0.182	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 53.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10496	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/5092	0.52	0/6871
1	B	0.34	0/5157	0.51	0/6958
All	All	0.34	0/10249	0.51	0/13829

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4969	0	4896	25	0
1	B	5026	0	4962	14	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	18	0	24	3	0
3	B	12	0	16	1	0
4	A	272	0	0	2	1
4	B	197	0	0	1	0
All	All	10496	0	9898	35	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:LYS:NZ	1:B:595:GLU:O	2.21	0.73
1:B:612:LYS:NZ	1:B:615:GLU:OE2	2.26	0.68
1:A:603:GLU:OE1	4:A:2267:HOH:O	2.17	0.55
1:A:467:GLU:OE2	1:A:541:ARG:NH2	2.40	0.54
1:A:199:ASP:N	1:A:199:ASP:OD1	2.40	0.52
1:A:146:SER:OG	1:B:218:GLN:NE2	2.40	0.52
1:A:221:ARG:HD2	1:B:221:ARG:NH1	2.27	0.50
1:A:566:THR:HG23	1:A:569:GLU:H	1.77	0.49
1:B:34:ASP:HB2	1:B:640:ARG:HH12	1.78	0.48
1:A:187:VAL:CG1	1:A:205:ILE:HD11	2.45	0.47
1:A:220:ILE:HG21	1:B:164:MET:HB3	1.98	0.46
1:B:596:ASP:OD2	4:B:2194:HOH:O	2.20	0.46
1:A:25:GLU:HG3	1:A:512:PHE:CE1	2.50	0.46
1:A:138:LEU:N	4:A:2070:HOH:O	2.48	0.46
1:A:374:VAL:HG22	1:A:401:ILE:CD1	2.46	0.46
1:A:330:CYS:O	1:A:331:ARG:HB3	2.16	0.45
1:B:530:LEU:HD13	1:B:557:HIS:CE1	2.51	0.45
1:A:475:ASP:HB2	3:A:1641:GOL:H32	1.99	0.43
3:A:1642:GOL:O3	3:A:1642:GOL:O1	2.36	0.43
1:A:5:TYR:CE2	1:B:3:VAL:HG21	2.54	0.43
1:B:528:VAL:HG21	1:B:539:ARG:HG2	1.99	0.43
1:A:303:ARG:NH2	1:A:568:GLU:OE1	2.52	0.42
1:A:122:MET:HB2	1:A:123:PRO:HD2	2.01	0.42
1:A:374:VAL:HG22	1:A:401:ILE:HD12	2.02	0.42
1:B:122:MET:HB2	1:B:123:PRO:HD2	2.01	0.42
1:A:330:CYS:O	1:A:331:ARG:CB	2.67	0.41
1:B:607:VAL:O	3:B:1642:GOL:O2	2.38	0.41
1:A:157:GLN:HB2	3:A:1642:GOL:H11	2.03	0.41
1:A:187:VAL:HG11	1:A:205:ILE:HD11	2.03	0.41
1:A:87:TRP:CD1	1:A:99:ALA:HB1	2.56	0.40
1:A:71:ILE:HD11	1:A:221:ARG:NH1	2.36	0.40
1:A:376:LEU:N	1:A:377:PRO:CD	2.85	0.40
1:B:571:PHE:CE2	1:B:575:LYS:HE2	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:471:ASP:OD2	4:A:2211:HOH:O[1_454]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	601/651 (92%)	592 (98%)	9 (2%)	0	100	100
1	B	607/651 (93%)	597 (98%)	10 (2%)	0	100	100
All	All	1208/1302 (93%)	1189 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	536/577 (93%)	532 (99%)	4 (1%)	84	81
1	B	544/577 (94%)	540 (99%)	4 (1%)	84	81
All	All	1080/1154 (94%)	1072 (99%)	8 (1%)	84	81

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	115	ASP
1	A	297	LEU
1	A	318	TYR
1	A	518	GLN
1	B	115	ASP
1	B	221	ARG
1	B	318	TYR
1	B	518	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	1642	-	5,5,5	0.30	0	5,5,5	0.87	0
3	GOL	A	1640	-	5,5,5	0.41	0	5,5,5	0.19	0
3	GOL	B	1642	-	5,5,5	0.34	0	5,5,5	0.26	0
3	GOL	B	1641	-	5,5,5	0.40	0	5,5,5	0.13	0
3	GOL	A	1641	-	5,5,5	0.38	0	5,5,5	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1642	-	-	0/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1640	-	-	4/4/4/4	-
3	GOL	B	1642	-	-	2/4/4/4	-
3	GOL	B	1641	-	-	4/4/4/4	-
3	GOL	A	1641	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1640	GOL	C1-C2-C3-O3
3	A	1640	GOL	O2-C2-C3-O3
3	B	1641	GOL	C1-C2-C3-O3
3	B	1642	GOL	C1-C2-C3-O3
3	A	1640	GOL	O1-C1-C2-C3
3	A	1641	GOL	C1-C2-C3-O3
3	B	1641	GOL	O2-C2-C3-O3
3	B	1642	GOL	O2-C2-C3-O3
3	A	1641	GOL	O2-C2-C3-O3
3	B	1641	GOL	O1-C1-C2-C3
3	B	1641	GOL	O1-C1-C2-O2
3	A	1640	GOL	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1642	GOL	2	0
3	B	1642	GOL	1	0
3	A	1641	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	606/651 (93%)	0.09	21 (3%) 44 51	15, 26, 52, 105	0
1	B	611/651 (93%)	0.13	22 (3%) 42 50	14, 29, 53, 82	0
All	All	1217/1302 (93%)	0.11	43 (3%) 44 51	14, 27, 53, 105	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	638	TYR	8.2
1	A	636	ILE	6.2
1	A	639	PRO	5.6
1	A	561	GLY	5.2
1	A	564	PHE	4.7
1	B	516	GLU	4.1
1	A	631	LYS	4.0
1	A	635	GLU	3.9
1	A	563	SER	3.7
1	A	562	GLU	3.2
1	A	634	ARG	3.0
1	B	631	LYS	3.0
1	B	408	VAL	3.0
1	B	50	ILE	2.9
1	B	325	ALA	2.9
1	B	603	GLU	2.9
1	A	24	TYR	2.8
1	A	607	VAL	2.8
1	B	9	ARG	2.7
1	B	24	TYR	2.7
1	A	63	VAL	2.7
1	B	634	ARG	2.6
1	A	457	ILE	2.6
1	B	443	VAL	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	608	CYS	2.6
1	B	633	ARG	2.5
1	B	328	SER	2.5
1	A	198	ARG	2.5
1	A	331	ARG	2.5
1	A	637	GLU	2.5
1	B	565	LYS	2.4
1	B	83	LEU	2.4
1	A	613	VAL	2.3
1	B	470	VAL	2.3
1	B	55	ILE	2.3
1	B	376	LEU	2.2
1	A	458	ASP	2.2
1	A	440	VAL	2.1
1	B	324	ASP	2.1
1	B	569	GLU	2.1
1	A	629	PHE	2.1
1	B	629	PHE	2.1
1	B	287	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	B	1642	6/6	0.49	0.29	54,56,58,59	0
3	GOL	A	1640	6/6	0.76	0.16	25,32,35,38	0
3	GOL	A	1642	6/6	0.84	0.15	52,59,60,61	0
3	GOL	B	1641	6/6	0.85	0.12	32,37,43,47	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	A	1641	6/6	0.87	0.21	42,45,48,50	0
2	ZN	B	652	1/1	0.99	0.02	35,35,35,35	0
2	ZN	A	652	1/1	0.99	0.02	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.