

Full wwPDB X-ray Structure Validation Report (i)

Oct 29, 2024 - 01:13 PM EDT

PDB ID	:	$4\mathrm{ECL}$
Title	:	Crystal structure of the cytoplasmic domain of vancomycin resistance serine
		racemase VanTg
Authors	:	Stogios, P.J.; Wawrzak, Z.; Minasov, G.; Evdokimova, E.; Egorova, O.;
		Cosme, J.; Di Leo, R.; Krishnamoorthy, M.; Meziane-Cherif, D.; Courvalin, P.;
		Savchenko, A.; Anderson, W.F.; Center for Structural Genomics of Infectious
		Diseases (CSGID)
Deposited on	:	2012-03-26
Resolution	:	2.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$		
R _{free}	164625	12358 (2.04-2.00)		
Clashscore	180529	13897 (2.04-2.00)		
Ramachandran outliers	177936	13770 (2.04-2.00)		
Sidechain outliers	177891	13769 (2.04-2.00)		
RSRZ outliers	164620	12358 (2.04-2.00)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	374	89%	9%	••
1	В	374	88%	11%	•
1	С	374	88%	11%	
1	D	374	92%	7%	•

Validation Pipeline (wwPDB-VP) : 2.39



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 12319 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		Atoms					ZeroOcc	AltConf	Trace
1	Δ	379	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	Se	0	0	0
L	Π	512	2889	1836	492	548	6	7	0		0
1	В	360	Total	С	Ν	Ο	\mathbf{S}	Se	0	3	0
	D	509	2892	1842	493	545	5	7	0	5	0
1	С	370	Total	С	Ν	Ο	\mathbf{S}	Se	0	7	0
		570	2920	1858	499	550	5	8			0
1	1 D	260	Total	C	Ν	0	S	Se	0	1	0
	309	2890	1839	488	550	6	$\overline{7}$		4	0	

• Molecule 1 is a protein called Serine racemase.

• Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	2	Total Cl 2 2	0	0
2	В	2	Total Cl 2 2	0	0
2	С	1	Total Cl 1 1	0	0
2	D	3	Total Cl 3 3	0	0

• Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O_4S).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
3	С	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
3	D	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	169	Total O 172 172	0	3
4	В	162	Total O 164 164	0	2
4	С	185	Total O 187 187	0	3
4	D	177	Total O 177 177	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Serine racemase



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	78.68Å 82.34Å 117.47Å	Deperitor
a, b, c, α , β , γ	90.00° 90.14° 90.00°	Depositor
$\mathbf{P}_{\text{oscolution}}(\hat{\mathbf{A}})$	19.91 - 2.02	Depositor
Resolution (A)	19.91 - 2.02	EDS
% Data completeness	98.2 (19.91-2.02)	Depositor
(in resolution range)	$90.7\ (19.91-2.02)$	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$< I/\sigma(I) > 1$	$1.40 (at 2.02 \text{\AA})$	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
B B c	0.181 , 0.233	Depositor
It, Itfree	0.184 , 0.232	DCC
R_{free} test set	1837 reflections (2.05%)	wwPDB-VP
Wilson B-factor $(Å^2)$	17.6	Xtriage
Anisotropy	1.595	Xtriage
Bulk solvent $k_{sol}(e/A^3), B_{sol}(A^2)$	0.37, 41.0	EDS
L-test for $twinning^2$	$< L >=0.47, < L^2>=0.30$	Xtriage
	0.010 for -k,-h,-l	
Estimated twinning fraction	0.009 for k,h,-l	Xtriage
	0.135 for h,-k,-l	
F_o, F_c correlation	0.98	EDS
Total number of atoms	12319	wwPDB-VP
Average B, all atoms $(Å^2)$	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 10.44% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, $\mathrm{SO4}$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.35	0/2927	0.52	0/3943	
1	В	0.35	0/2935	0.54	0/3951	
1	С	0.35	0/2975	0.52	0/4004	
1	D	0.36	0/2933	0.52	0/3951	
All	All	0.35	0/11770	0.53	0/15849	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2889	0	2965	33	0
1	В	2892	0	2984	29	0
1	С	2920	0	3019	24	0
1	D	2890	0	2966	18	0
2	А	2	0	0	0	0
2	В	2	0	0	0	0
2	С	1	0	0	0	0
2	D	3	0	0	0	0
3	A	5	0	0	1	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	5	0	0	1	0
3	С	5	0	0	0	0
3	D	5	0	0	0	0
4	А	172	0	0	2	0
4	В	164	0	0	1	0
4	С	187	0	0	2	0
4	D	177	0	0	1	0
All	All	12319	0	11934	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:A:314:MSE:HE2	1:B:38:LYS:HG3	1.49	0.92	
1:A:314:MSE:HE3	1:B:39:ALA:HB3	1.54	0.88	
1:D:358:ASN:HD22	1:D:358:ASN:H	1.39	0.69	
1:A:358:ASN:HD22	1:A:358:ASN:H	1.39	0.69	
3:B:403:SO4:O3	4:B:656:HOH:O	2.11	0.68	
3:A:403:SO4:O1	4:A:638:HOH:O	2.12	0.67	
1:A:40:GLU:OE2	1:A:71:ARG:NH2	2.27	0.64	
1:B:27:MSE:HG2	1:B:218:ILE:HD12	1.80	0.62	
1:C:132:ARG:HD3	1:D:316:GLN:HE21	1.64	0.61	
1:B:71:ARG:NH2	1:B:75:TYR:OH	2.30	0.60	
1:D:321:VAL:HG21	1:D:327:VAL:HG11	1.86	0.58	
1:C:55:GLN:NE2	4:C:659:HOH:O	2.26	0.57	
1:D:358:ASN:H	1:D:358:ASN:ND2	2.02	0.57	
1:C:18:HIS:HE1	1:C:240:ASP:O	1.87	0.56	
1:A:39:ALA:CB	1:B:314:MSE:HE3	2.35	0.56	
1:C:358:ASN:H	1:C:358:ASN:HD22	1.52	0.56	
1:C:231:THR:HG22	1:C:347:PRO:HB2	1.87	0.56	
1:C:355:SER:OG	1:C:359:GLU:OE1	2.21	0.56	
1:C:311:LYS:NZ	4:C:507:HOH:O	2.38	0.56	
1:D:34:MSE:CE	1:D:219:ARG:HG2	2.36	0.56	
1:D:23:LEU:HD13	1:D:220[A]:VAL:HG21	1.89	0.55	
1:D:27:MSE:HG2	1:D:31:CYS:HB3	1.88	0.55	
1:D:34:MSE:HE2	1:D:219:ARG:HG2	1.88	0.55	
1:B:222[A]:VAL:HG22	1:B:227:VAL:HG23	1.88	0.55	
1:C:71[A]:ARG:NE	1:C:75:TYR:OH	2.37	0.55	
1:C:314:MSE:SE	1:D:38:LYS:HE3	2.58	0.54	



4ECL

Atom-1	Atom-2	Interatomic	Clash
	Atom-2	distance (Å)	overlap (Å)
1:A:314:MSE:CE	1:B:38:LYS:HG3	2.30	0.54
1:B:34:MSE:CE	1:B:219:ARG:HG2	2.39	0.53
1:A:268:ALA:HB1	1:A:294:CYS:SG	2.48	0.53
1:B:160:HIS:HD2	1:B:219:ARG:HH21	1.57	0.52
1:B:18:HIS:HE1	1:B:240:ASP:O	1.94	0.51
1:C:46:MSE:HE1	1:C:68:GLU:HG2	1.92	0.51
1:A:266:SER:C	1:A:268:ALA:H	2.13	0.51
1:A:267:ARG:H	1:A:267:ARG:HE	1.57	0.51
1:A:290:ARG:NH1	4:A:658:HOH:O	2.44	0.51
1:A:358:ASN:HD22	1:A:358:ASN:N	2.09	0.50
1:A:174:PHE:HA	1:A:177:LYS:HE3	1.94	0.50
1:A:38:LYS:HG3	1:B:314:MSE:HE1	1.91	0.50
1:B:37:VAL:HG11	1:B:50:THR:HG22	1.94	0.49
1:B:273:ARG:HG3	1:B:274:ASP:O	2.13	0.49
1:A:39:ALA:HB3	1:B:314:MSE:HE3	1.93	0.49
1:A:200:HIS:HB3	1:A:217:TYR:HB2	1.96	0.48
1:D:316:GLN:NE2	4:D:501:HOH:O	2.45	0.47
1:A:47:TYR:HD1	1:A:75:TYR:CG	2.33	0.47
1:A:314:MSE:CE	1:B:39:ALA:HB3	2.36	0.46
1:A:171:ASP:OD2	1:B:267:ARG:NH2	2.49	0.46
1:C:131:HIS:HD2	1:D:262:SER:O	1.98	0.46
1:C:345:THR:HB	1:C:347:PRO:HD2	1.97	0.46
1:B:209:ASN:O	1:B:236:LYS:HD2	2.16	0.46
1:C:34:MSE:CE	1:C:219:ARG:HG2	2.46	0.45
1:A:38:LYS:HG3	1:B:314:MSE:CE	2.46	0.45
1:D:128:THR:O	1:D:178:GLN:HG2	2.17	0.45
1:D:220[A]:VAL:HG22	1:D:224:LEU:H	1.82	0.45
1:A:97:LYS:HB3	1:A:97:LYS:HE2	1.89	0.45
1:B:158:PHE:HB2	1:B:200:HIS:O	2.17	0.45
1:A:266:SER:C	1:A:268:ALA:N	2.70	0.44
1:B:284:TYR:CE1	1:B:314:MSE:HG3	2.53	0.44
1:C:6:ARG:HG3	1:C:281:PRO:HB2	1.99	0.44
1:C:105:ILE:HD12	1:D:253:LEU:HD12	1.99	0.44
1:C:130[B]:MSE:SE	1:D:264:GLY:HA2	2.68	0.44
1:B:34:MSE:HE2	1:B:219:ARG:HG2	2.00	0.44
1:A:13:LEU:HD13	1:A:52:TYR:CD1	2.53	0.44
1:A:231:THR:HA	1:A:232:ASN:HA	1.67	0.43
1:A:253:LEU:HG	1:A:254:ILE:N	2.33	0.43
1:C:37:VAL:O	1:C:46:MSE:HE2	2.19	0.43
1:C:130[B]:MSE:HE3	1:C:132:ARG:HD2	2.01	0.43
1:A:140:LYS:HD2	1:A:188:TRP:CG	2.54	0.42



Atom 1	Atom 2	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:C:40:GLU:OE2	1:C:71[A]:ARG:NH2	2.32	0.42	
1:C:207:LEU:HB2	1:C:218:ILE:HD13	2.02	0.42	
1:A:104:LEU:HB2	1:A:124:ILE:HA	2.02	0.42	
1:C:186:LEU:HD13	1:C:196:ILE:HG21	2.00	0.42	
1:A:314:MSE:HE3	1:B:39:ALA:CB	2.36	0.42	
1:B:31:CYS:SG	1:B:218:ILE:HG13	2.60	0.42	
1:A:355:SER:OG	1:A:356:ILE:N	2.51	0.42	
1:B:250:LYS:HE2	1:B:332:ILE:HG12	2.01	0.42	
1:B:44:HIS:CD2	1:B:245:LEU:HB2	2.54	0.41	
1:B:22:THR:HG22	1:B:239:LEU:HD13	2.02	0.41	
1:B:277:ILE:HD12	1:B:309:VAL:HG21	2.02	0.41	
1:C:132:ARG:HD3	1:D:316:GLN:NE2	2.32	0.41	
1:D:325:PRO:HA	1:D:326:ASN:HA	1.58	0.41	
1:C:173:ALA:O	1:C:177:LYS:HG2	2.20	0.41	
1:D:284:TYR:HA	1:D:288:PHE:O	2.21	0.41	
1:A:39:ALA:HB2	1:B:314:MSE:HE3	2.01	0.41	
1:C:262:SER:OG	1:C:263:VAL:N	2.53	0.41	
1:A:229:SER:HA	1:A:357:THR:HB	2.02	0.41	
1:A:236:LYS:HA	1:A:236:LYS:HD3	1.87	0.41	
1:A:265:TYR:O	1:A:265:TYR:CG	2.74	0.41	
1:B:27:MSE:HE1	1:B:33:LEU:HA	2.03	0.41	
1:A:264:GLY:O	1:A:266:SER:N	2.54	0.40	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	370/374~(99%)	357~(96%)	10 (3%)	3~(1%)	16	11
1	В	368/374~(98%)	357~(97%)	11 (3%)	0	100	100
1	С	373/374~(100%)	364~(98%)	9 (2%)	0	100	100



001000	contributed from proceed by page									
Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentile	es			
1	D	369/374~(99%)	353~(96%)	16 (4%)	0	100 100)			
All	All	1480/1496~(99%)	1431 (97%)	46 (3%)	3(0%)	44 41				

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	265	TYR
1	А	229	SER
1	А	267	ARG

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric Out		Percentiles
1	А	317/311~(102%)	313~(99%)	4 (1%)	65 70
1	В	318/311~(102%)	311~(98%)	7 (2%)	47 49
1	С	323/311~(104%)	312~(97%)	11 (3%)	32 31
1	D	319/311~(103%)	313~(98%)	6~(2%)	52 55
All	All	1277/1244 (103%)	1249~(98%)	28 (2%)	48 49

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	\mathbf{Res}	Type
1	А	138	GLU
1	А	267	ARG
1	А	315	ASP
1	А	358	ASN
1	В	28	SER
1	В	230	SER
1	В	234	LYS
1	В	256	LYS
1	В	328	LYS
1	В	372	ARG
1	В	373	ARG



Mol	Chain	Res	Type
1	С	99	GLU
1	С	115	LYS
1	С	160	HIS
1	С	170[A]	ASN
1	С	170[B]	ASN
1	С	231	THR
1	С	234	LYS
1	С	273	ARG
1	С	276	LEU
1	С	297	SER
1	С	315	ASP
1	D	3	ASP
1	D	28	SER
1	D	160	HIS
1	D	265	TYR
1	D	328	LYS
1	D	358	ASN

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such side chains are listed below:

Mol	Chain	Res	Type
1	А	358	ASN
1	В	18	HIS
1	В	24	GLN
1	С	18	HIS
1	С	102	GLN
1	С	358	ASN
1	D	316	GLN
1	D	358	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Chain		Dec Link	Bond lengths		Bond angles									
	туре	Unam	nes	nes	rtes	nes	nes	Res Link	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
3	SO4	D	404	-	4,4,4	0.27	0	6,6,6	0.22	0				
3	SO4	С	402	-	4,4,4	0.19	0	6,6,6	0.39	0				
3	SO4	А	403	-	4,4,4	0.29	0	6,6,6	0.17	0				
3	SO4	В	403	-	4,4,4	0.32	0	6,6,6	0.12	0				

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	А	403	SO4	1	0
3	В	403	SO4	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2		$Z{>}2$	$OWAB(Å^2)$	Q < 0.9
1	А	365/374~(97%)	-1.87	0 10	00	100	19, 32, 59, 103	0
1	В	362/374~(96%)	-1.85	0 10	00	100	17, 33, 59, 100	3~(0%)
1	С	363/374~(97%)	-1.86	0 10	00	100	16, 31, 57, 103	6(1%)
1	D	362/374~(96%)	-1.91	0 10	00	100	13, 29, 62, 101	4 (1%)
All	All	1452/1496~(97%)	-1.88	0 10	00	100	13, 31, 60, 103	13 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	CL	А	401	1/1	1.00	0.01	24,24,24,24	1
2	CL	А	402	1/1	1.00	0.02	20,20,20,20	1
2	CL	В	401	1/1	1.00	0.03	33,33,33,33	1
2	CL	В	402	1/1	1.00	0.04	13,13,13,13	1



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	CL	С	401	1/1	1.00	0.02	33,33,33,33	1
2	CL	D	401	1/1	1.00	0.02	$53,\!53,\!53,\!53$	0
2	CL	D	402	1/1	1.00	0.01	36, 36, 36, 36	0
2	CL	D	403	1/1	1.00	0.01	39,39,39,39	0
3	SO4	А	403	5/5	1.00	0.02	$23,\!26,\!31,\!35$	0
3	SO4	В	403	5/5	1.00	0.01	$22,\!22,\!26,\!27$	5
3	SO4	С	402	5/5	1.00	0.02	23,26,30,30	0
3	SO4	D	404	5/5	1.00	0.02	24,26,31,34	0

Continued from previous page...

6.5 Other polymers (i)

There are no such residues in this entry.

