



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 13, 2024 – 10:42 AM EDT

PDB ID : 4G9K
Title : Structure of the Ndi1 protein from *Saccharomyces cerevisiae*
Authors : Iwata, M.; Lee, Y.; Yamashita, T.; Yagi, T.; Iwata, S.; Cameron, A.D.; Maher, M.J.
Deposited on : 2012-07-24
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.36.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	471	 7% 86% 11% ..
1	B	471	 9% 87% 11% .

2 Entry composition i


There are 3 unique types of molecules in this entry. The entry contains 7482 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rotenone-insensitive NADH-ubiquinone oxidoreductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	463	Total 3679	C 2381	N 620	O 673	S 5	0	1	0
1	B	463	Total 3679	C 2381	N 620	O 673	S 5	0	1	0

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 53	C 27	N 9	O 15	P 2	0	0
2	B	1	Total 53	C 27	N 9	O 15	P 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	9	Total O 9 9	0	0
3	B	9	Total O 9 9	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Rotenone-insensitive NADH-ubiquinone oxidoreductase



- Molecule 1: Rotenone-insensitive NADH-ubiquinone oxidoreductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	116.12Å 165.56Å 70.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.32 – 2.70 39.32 – 2.70	Depositor EDS
% Data completeness (in resolution range)	73.7 (39.32-2.70) 73.8 (39.32-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.24 (at 2.69Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.234 , 0.268 0.217 , 0.248	Depositor DCC
R_{free} test set	1401 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	71.4	Xtrriage
Anisotropy	0.171	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7482	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/3770	0.48	0/5103
1	B	0.40	0/3770	0.49	0/5103
All	All	0.40	0/7540	0.49	0/10206

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3679	0	3737	37	0
1	B	3679	0	3737	34	0
2	A	53	0	31	1	0
2	B	53	0	31	1	0
3	A	9	0	0	0	0
3	B	9	0	0	0	0
All	All	7482	0	7536	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:502:LEU:HD12	1:A:506:LYS:HA	1.65	0.79
1:B:502:LEU:HD12	1:B:506:LYS:HA	1.66	0.78
1:B:176:VAL:HG22	1:B:384:ASN:HD22	1.63	0.63
1:A:176:VAL:HG22	1:A:384:ASN:HD22	1.65	0.61
1:A:88:PHE:CZ	1:A:487:LEU:HD22	2.38	0.59
1:B:380:ALA:HB1	1:B:385:ALA:HB2	1.85	0.59
1:B:290:GLN:NE2	1:B:302:LEU:HD11	2.20	0.57
1:A:380:ALA:HB1	1:A:385:ALA:HB2	1.88	0.56
1:B:101:THR:HG23	1:B:502:LEU:HD22	1.88	0.56
1:A:239:THR:HG23	2:A:601:FAD:HM73	1.89	0.55
1:B:88:PHE:CZ	1:B:487:LEU:HD22	2.41	0.55
1:A:290:GLN:NE2	1:A:302:LEU:HD11	2.22	0.53
1:A:101:THR:HG23	1:A:502:LEU:HD22	1.92	0.51
1:A:104:GLU:HB3	1:B:513:LEU:HD21	1.92	0.51
1:A:102:VAL:HG11	1:A:107:ILE:HG23	1.93	0.50
1:B:427:ILE:HG22	1:B:431:PHE:CE1	2.47	0.50
1:A:427:ILE:HG22	1:A:431:PHE:CE1	2.47	0.49
1:B:132:ILE:HD12	1:B:350:LEU:HD22	1.95	0.49
1:A:132:ILE:HD12	1:A:350:LEU:HD22	1.95	0.49
1:A:116:LEU:HD13	1:A:150:LEU:HD13	1.94	0.49
1:A:173:ILE:HD12	1:A:173:ILE:N	2.27	0.49
1:B:173:ILE:N	1:B:173:ILE:HD12	2.29	0.48
1:B:116:LEU:HD13	1:B:150:LEU:HD13	1.95	0.48
1:A:279:ASN:C	1:A:279:ASN:HD22	2.18	0.47
1:B:280:MET:HE3	1:B:280:MET:H	1.79	0.47
1:A:376:ASN:O	1:A:427:ILE:HD12	2.14	0.47
1:B:61:SER:HB3	1:B:111:ILE:HD11	1.96	0.47
1:B:102:VAL:HG11	1:B:107:ILE:HG23	1.96	0.47
1:A:483:LEU:HD21	1:A:495:VAL:HG23	1.97	0.47
1:A:97:ALA:HB3	1:A:98:PRO:HD3	1.96	0.47
1:B:239:THR:HG23	2:B:601:FAD:HM73	1.96	0.47
1:A:279:ASN:HD22	1:A:280:MET:N	2.13	0.46
1:B:197:GLU:H	1:B:200:ASN:HD22	1.62	0.46
1:A:216:ASN:HB2	1:A:262:LEU:HD11	1.98	0.46
1:A:289:ALA:HA	1:A:450:LEU:HD13	1.96	0.46
1:B:376:ASN:O	1:B:427:ILE:HD12	2.16	0.45
1:B:97:ALA:HB3	1:B:98:PRO:HD3	1.99	0.45
1:B:216:ASN:HB2	1:B:262:LEU:HD11	1.98	0.45
1:B:174:SER:C	1:B:384:ASN:HD21	2.20	0.44
1:A:196:LYS:H	1:A:200:ASN:HD22	1.64	0.44
1:A:513:LEU:HD21	1:B:104:GLU:HB3	1.98	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:289:ALA:HA	1:B:450:LEU:HD13	2.00	0.43
1:A:61:SER:HB3	1:A:111:ILE:HD11	2.00	0.43
1:A:447:LEU:HD23	1:A:457:ALA:HB2	2.01	0.43
1:B:279:ASN:HD22	1:B:279:ASN:C	2.23	0.43
1:B:101:THR:HG23	1:B:502:LEU:CD2	2.49	0.42
1:A:99:VAL:HG21	1:A:247:LEU:HA	2.02	0.42
1:B:483:LEU:HD21	1:B:495:VAL:HG23	2.00	0.42
1:A:447:LEU:HD23	1:A:457:ALA:CB	2.50	0.42
1:A:274:LEU:HB3	1:A:275:PRO:HD2	2.01	0.42
1:B:486:ILE:O	1:B:492:ARG:NH2	2.53	0.42
1:B:99:VAL:HG21	1:B:247:LEU:HA	2.02	0.42
1:A:254:ASP:OD1	1:A:506:LYS:HB2	2.19	0.42
1:B:254:ASP:OD1	1:B:506:LYS:HB2	2.20	0.42
1:A:377:ASN:HD22	1:A:377:ASN:H	1.68	0.41
1:A:486:ILE:O	1:A:492:ARG:NH2	2.52	0.41
1:A:101:THR:HG23	1:A:502:LEU:CD2	2.51	0.41
1:A:280:MET:HE3	1:A:280:MET:H	1.86	0.41
1:A:72:ILE:O	1:A:118:LYS:NZ	2.52	0.41
1:B:99:VAL:CG1	1:B:250:TYR:HB2	2.51	0.41
1:A:104:GLU:CB	1:B:513:LEU:HD21	2.50	0.41
1:A:174:SER:C	1:A:384:ASN:HD21	2.23	0.41
1:B:274:LEU:HB3	1:B:275:PRO:HD2	2.01	0.41
1:B:380:ALA:CB	1:B:385:ALA:HB2	2.49	0.41
1:A:380:ALA:CB	1:A:385:ALA:HB2	2.51	0.40
1:B:279:ASN:HD22	1:B:280:MET:N	2.19	0.40
1:A:102:VAL:HG12	1:A:103:ASP:O	2.21	0.40
1:B:46:ILE:N	1:B:46:ILE:HD12	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	461/471 (98%)	443 (96%)	16 (4%)	2 (0%)	34	60
1	B	461/471 (98%)	443 (96%)	16 (4%)	2 (0%)	34	60
All	All	922/942 (98%)	886 (96%)	32 (4%)	4 (0%)	34	60

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	44	LYS
1	B	44	LYS
1	A	507	ARG
1	B	507	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	396/402 (98%)	390 (98%)	6 (2%)	65	86
1	B	396/402 (98%)	390 (98%)	6 (2%)	65	86
All	All	792/804 (98%)	780 (98%)	12 (2%)	65	86

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	279	ASN
1	A	280	MET
1	A	314	LEU
1	A	366	VAL
1	A	377	ASN
1	A	460	ARG
1	B	279	ASN
1	B	280	MET
1	B	314	LEU
1	B	366	VAL
1	B	377	ASN
1	B	460	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (25) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	137	ASN
1	A	200	ASN
1	A	248	GLN
1	A	252	HIS
1	A	253	GLN
1	A	279	ASN
1	A	290	GLN
1	A	377	ASN
1	A	384	ASN
1	A	394	GLN
1	A	415	ASN
1	A	442	ASN
1	B	137	ASN
1	B	156	HIS
1	B	200	ASN
1	B	248	GLN
1	B	252	HIS
1	B	253	GLN
1	B	279	ASN
1	B	290	GLN
1	B	377	ASN
1	B	384	ASN
1	B	394	GLN
1	B	415	ASN
1	B	442	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	B	601	-	54,58,58	1.35	5 (9%)	71,89,89	1.46	12 (16%)
2	FAD	A	601	-	54,58,58	1.32	4 (7%)	71,89,89	1.36	12 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	B	601	-	-	3/30/50/50	0/6/6/6
2	FAD	A	601	-	-	0/30/50/50	0/6/6/6

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	FAD	C9A-C5X	5.96	1.50	1.41
2	B	601	FAD	C9A-C5X	5.50	1.50	1.41
2	B	601	FAD	C8-C7	3.56	1.49	1.40
2	A	601	FAD	C8-C7	3.51	1.49	1.40
2	B	601	FAD	C4X-N5	2.52	1.36	1.30
2	B	601	FAD	C4-N3	-2.30	1.34	1.38
2	B	601	FAD	C2A-N3A	2.27	1.35	1.32
2	A	601	FAD	C4X-N5	2.21	1.35	1.30
2	A	601	FAD	C4-N3	-2.09	1.34	1.38

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	FAD	N3A-C2A-N1A	-4.81	122.14	128.67
2	B	601	FAD	N3A-C2A-N1A	-4.41	122.68	128.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	FAD	C4'-C3'-C2'	-4.30	106.41	113.57
2	B	601	FAD	C4-C4X-N5	4.03	123.78	118.21
2	A	601	FAD	C1B-N9A-C4A	-3.07	121.25	126.64
2	A	601	FAD	C4-C4X-N5	3.02	122.38	118.21
2	A	601	FAD	C4X-C10-N10	2.65	120.28	116.48
2	A	601	FAD	O2-C2-N1	-2.65	117.40	121.80
2	B	601	FAD	C4X-C10-N10	2.61	120.22	116.48
2	B	601	FAD	C4A-C5A-N7A	-2.60	106.59	109.34
2	B	601	FAD	C1B-N9A-C4A	-2.46	122.31	126.64
2	A	601	FAD	O2P-P-O1P	2.46	123.90	112.44
2	B	601	FAD	C10-N1-C2	2.42	122.09	116.85
2	B	601	FAD	C4X-C10-N1	-2.42	118.66	124.59
2	B	601	FAD	O2-C2-N1	-2.39	117.83	121.80
2	A	601	FAD	O4-C4-C4X	-2.38	120.25	126.53
2	A	601	FAD	C4X-C10-N1	-2.36	118.81	124.59
2	B	601	FAD	C5X-N5-C4X	2.36	121.90	118.09
2	B	601	FAD	C10-C4X-N5	-2.35	120.00	124.81
2	A	601	FAD	C4A-C5A-N7A	-2.23	106.98	109.34
2	B	601	FAD	O2P-P-O1P	2.20	122.68	112.44
2	A	601	FAD	C10-N1-C2	2.16	121.53	116.85
2	A	601	FAD	C4X-C4-N3	2.15	118.72	113.25
2	A	601	FAD	C4-N3-C2	-2.13	121.85	125.64

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	601	FAD	PA-O3P-P-O5'
2	B	601	FAD	P-O3P-PA-O1A
2	B	601	FAD	P-O3P-PA-O2A


There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	FAD	1	0
2	A	601	FAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	463/471 (98%)	0.39	34 (7%) 15 13	39, 63, 102, 142	0
1	B	463/471 (98%)	0.47	42 (9%) 9 7	38, 64, 125, 157	0
All	All	926/942 (98%)	0.43	76 (8%) 11 9	38, 64, 112, 157	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	427	ILE	10.1
1	A	50	HIS	9.1
1	A	49	GLN	8.4
1	B	50	HIS	8.4
1	B	417	GLN	7.5
1	B	52	ASP	7.2
1	B	49	GLN	7.1
1	A	322	ASP	6.8
1	B	411	ALA	6.7
1	B	431	PHE	6.4
1	A	153	PRO	6.4
1	B	430	LEU	6.3
1	A	154	GLU	6.1
1	B	154	GLU	5.5
1	A	417	GLN	5.3
1	B	153	PRO	5.1
1	B	416	PHE	5.1
1	A	320	HIS	4.8
1	B	322	ASP	4.4
1	A	416	PHE	4.3
1	B	464	ARG	4.2
1	A	51	SER	4.1
1	B	51	SER	4.0
1	B	429	LEU	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	319	LYS	3.8
1	B	426	LYS	3.8
1	A	468	THR	3.7
1	A	221	GLY	3.7
1	A	352	LYS	3.6
1	B	150	LEU	3.4
1	A	151	TYR	3.4
1	B	410	MET	3.4
1	B	412	GLN	3.3
1	A	470	GLY	3.3
1	B	433	GLU	3.3
1	B	324	LYS	3.3
1	A	119	LYS	3.3
1	B	371	GLN	3.3
1	A	52	ASP	3.3
1	A	321	GLU	3.2
1	B	463	LYS	3.2
1	B	321	GLU	3.1
1	B	119	LYS	3.1
1	B	151	TYR	3.1
1	A	155	ASN	3.1
1	A	462	GLY	3.0
1	A	152	GLN	3.0
1	A	469	GLY	3.0
1	B	320	HIS	2.9
1	B	162	ALA	2.9
1	A	264	GLU	2.9
1	B	468	THR	2.9
1	B	414	PRO	2.9
1	A	362	ARG	2.8
1	B	434	ASN	2.8
1	B	469	GLY	2.8
1	A	135	ASP	2.7
1	B	408	ASP	2.7
1	B	376	ASN	2.7
1	B	171	TYR	2.6
1	B	369	PHE	2.6
1	A	323	GLY	2.6
1	A	121	ASN	2.5
1	A	324	LYS	2.5
1	B	367	ASN	2.3
1	B	163	GLU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	428	ASP	2.2
1	A	464	ARG	2.1
1	B	252	HIS	2.1
1	A	43	MET	2.1
1	B	264	GLU	2.1
1	A	150	LEU	2.0
1	A	303	ARG	2.0
1	A	361	LYS	2.0
1	A	61	SER	2.0
1	B	147	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FAD	A	601	53/53	0.97	0.22	36,44,47,49	0
2	FAD	B	601	53/53	0.97	0.20	38,42,57,58	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD B 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.