



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 19, 2024 – 11:36 AM EDT

PDB ID : 4HA0  
Title : Structure of Geobacillus kaustophilus lactonase, mutant R230D with Zn<sup>2+</sup>  
Authors : Xue, B.; Chow, J.Y.; Yew, W.S.; Robinson, R.C.  
Deposited on : 2012-09-25  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	
1	B	330	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5369 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphotriesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	323	2540	1615	432	477	16	0	0	0
1	B	323	2540	1615	432	477	16	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	EXPRESSION TAG	UNP Q5KZU5
A	-2	SER	-	EXPRESSION TAG	UNP Q5KZU5
A	-1	HIS	-	EXPRESSION TAG	UNP Q5KZU5
A	0	ASN	-	EXPRESSION TAG	UNP Q5KZU5
A	230	ASP	ARG	ENGINEERED MUTATION	UNP Q5KZU5
B	-3	GLY	-	EXPRESSION TAG	UNP Q5KZU5
B	-2	SER	-	EXPRESSION TAG	UNP Q5KZU5
B	-1	HIS	-	EXPRESSION TAG	UNP Q5KZU5
B	0	ASN	-	EXPRESSION TAG	UNP Q5KZU5
B	230	ASP	ARG	ENGINEERED MUTATION	UNP Q5KZU5

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Fe	0	0
			1	1		
2	B	1	Total	Fe	0	0
			1	1		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is HYDROXIDE ION (three-letter code: OH) (formula: HO).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	O	0	0
			1	1		
4	B	1	Total	O	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	137	Total	O	0	0
			137	137		
5	B	146	Total	O	0	0
			146	146		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphotriesterase



- Molecule 1: Phosphotriesterase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.97Å 76.22Å 134.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.98 – 1.90 29.97 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.5 (19.98-1.90) 97.4 (29.97-1.90)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.14 (at 1.91Å)	Xtrriage
Refinement program	PHENIX 1.7.3_928	Depositor
R, $R_{free}$	0.259 , 0.303 0.268 , 0.303	Depositor DCC
$R_{free}$ test set	2851 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.1	Xtrriage
Anisotropy	0.292	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 51.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5369	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 89.56 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.2567e-08. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, KCX, OH, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.62	0/2586	0.62	1/3500 (0.0%)
1	B	0.62	1/2586 (0.0%)	0.62	0/3500
All	All	0.62	1/5172 (0.0%)	0.62	1/7000 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	100	TYR	CD1-CE1	-5.16	1.31	1.39

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	115	LEU	CA-CB-CG	5.37	127.65	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2540	0	2477	33	0
1	B	2540	0	2478	29	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	137	0	0	1	1
5	B	146	0	0	0	0
All	All	5369	0	4955	61	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:234:GLN:HG2	1:A:241:THR:HA	1.60	0.82
1:B:259:LYS:HG2	1:B:314:GLN:HE21	1.56	0.71
1:B:100:TYR:CE2	1:B:102:GLY:HA3	2.34	0.62
1:A:78:PRO:HG2	1:A:134:ILE:HG21	1.84	0.59
1:B:78:PRO:HG2	1:B:134:ILE:HG21	1.86	0.57
1:B:112:ARG:HA	1:B:115:LEU:HG	1.89	0.55
1:B:208:CYS:HB3	1:B:228:PHE:CD2	2.42	0.54
1:A:208:CYS:HB3	1:A:228:PHE:CD2	2.43	0.53
1:A:265:HIS:HB2	1:A:267:THR:HG23	1.90	0.53
1:A:43:ARG:HA	5:A:593:HOH:O	2.10	0.51
1:B:265:HIS:HB2	1:B:267:THR:HG23	1.92	0.51
1:B:250:LEU:HD11	1:B:305:GLU:HG3	1.91	0.51
1:A:250:LEU:HD11	1:A:305:GLU:HG3	1.95	0.49
1:B:282:PHE:CD1	1:B:282:PHE:N	2.81	0.48
1:B:210:ASN:O	1:B:216:HIS:HE1	1.97	0.47
1:A:237:VAL:HG12	1:A:237:VAL:O	2.15	0.47
1:A:210:ASN:O	1:A:216:HIS:HE1	1.97	0.47
1:B:106:PRO:O	1:B:110:GLN:HG3	2.14	0.47
1:B:19:LYS:HG2	1:B:63:ILE:HD13	1.96	0.47
1:A:111:PHE:O	1:A:114:LEU:HB2	2.15	0.47
1:A:282:PHE:CD1	1:A:282:PHE:N	2.81	0.47
1:A:19:LYS:HG2	1:A:63:ILE:HD13	1.97	0.47
1:B:152:ARG:HE	1:B:152:ARG:HB2	1.51	0.46
1:A:207:MET:O	1:A:216:HIS:NE2	2.48	0.46
1:B:55:ALA:HB3	1:B:88:THR:HG21	1.98	0.46
1:A:239:ALA:HA	1:A:240:PRO:HD2	1.83	0.45
1:A:55:ALA:HB3	1:A:88:THR:HG21	1.98	0.45

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:203:VAL:HG22	1:A:225:TYR:HB2	1.97	0.45
1:B:145:KCX:OQ2	1:B:178:HIS:HB2	2.17	0.45
1:B:203:VAL:HG22	1:B:225:TYR:HB2	1.99	0.45
1:A:57:LYS:HD3	1:A:292:GLU:OE1	2.17	0.45
1:B:57:LYS:HD3	1:B:292:GLU:OE1	2.17	0.44
1:A:39:ARG:NE	1:B:124:ASP:OD2	2.36	0.44
1:A:100:TYR:CE2	1:A:102:GLY:HA3	2.53	0.44
1:B:303:LYS:HE2	1:B:303:LYS:HB3	1.68	0.44
1:A:152:ARG:HE	1:A:152:ARG:HB2	1.51	0.44
1:A:277:THR:HG22	1:A:278:LEU:N	2.32	0.44
1:A:282:PHE:N	1:A:282:PHE:HD1	2.16	0.44
1:A:106:PRO:O	1:A:110:GLN:HG3	2.18	0.44
1:B:130:LEU:HD11	1:B:144:ILE:HD11	1.99	0.44
1:A:41:THR:CG2	1:A:42:PHE:N	2.81	0.43
1:B:296:VAL:HG23	1:B:297:ASN:OD1	2.18	0.43
1:A:207:MET:HG3	1:A:226:ILE:HB	2.01	0.43
1:A:299:ILE:HB	1:A:300:PRO:HD3	2.01	0.43
1:A:277:THR:CG2	1:A:278:LEU:N	2.81	0.43
1:B:8:VAL:HA	1:B:139:ILE:HG23	2.01	0.43
1:B:299:ILE:HB	1:B:300:PRO:HD3	2.01	0.43
1:A:61:HIS:CD2	1:A:295:PHE:HB2	2.53	0.43
1:B:55:ALA:HB1	1:B:90:LEU:HD22	2.00	0.43
1:B:207:MET:O	1:B:216:HIS:NE2	2.52	0.42
1:A:280:GLU:N	1:A:281:PRO:HD2	2.35	0.42
1:B:207:MET:HG3	1:B:226:ILE:HB	2.02	0.41
1:B:282:PHE:N	1:B:282:PHE:HD1	2.15	0.41
1:A:3:GLU:HG3	1:A:4:MET:HG2	2.03	0.41
1:A:60:ARG:C	1:A:61:HIS:ND1	2.74	0.41
1:A:303:LYS:HB3	1:A:303:LYS:HE2	1.70	0.41
1:B:39:ARG:HH11	1:B:39:ARG:HD3	1.76	0.41
1:A:8:VAL:HA	1:A:139:ILE:HG23	2.02	0.41
1:B:41:THR:CG2	1:B:42:PHE:N	2.83	0.40
1:B:3:GLU:HG3	1:B:4:MET:HG2	2.04	0.40
1:A:259:LYS:HG2	1:A:314:GLN:HE21	1.86	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:558:HOH:O	5:A:559:HOH:O 3_545	2.08	0.12

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	320/330 (97%)	313 (98%)	6 (2%)	1 (0%)	41	31
1	B	320/330 (97%)	312 (98%)	7 (2%)	1 (0%)	41	31
All	All	640/660 (97%)	625 (98%)	13 (2%)	2 (0%)	41	31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	69	PRO
1	B	69	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/267 (98%)	259 (98%)	4 (2%)	65	62
1	B	263/267 (98%)	259 (98%)	4 (2%)	65	62
All	All	526/534 (98%)	518 (98%)	8 (2%)	65	62

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	HIS
1	A	99	TYR
1	A	178	HIS
1	A	264	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	41	THR
1	B	99	TYR
1	B	115	LEU
1	B	264	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	314	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	KCX	B	145	2,1,3	10,11,12	1.20	0	6,12,14	1.18	0
1	KCX	A	145	2,1,3	10,11,12	0.93	0	6,12,14	1.27	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	B	145	2,1,3	-	0/9/10/12	-
1	KCX	A	145	2,1,3	-	0/9/10/12	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	145	KCX	CE-NZ-CX	-2.35	117.99	121.98

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	145	KCX	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic and 2 are modelled with single atom - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	322/330 (97%)	1.03	61 (18%) <b>1</b> <b>1</b>	20, 50, 85, 128	0
1	B	322/330 (97%)	0.50	18 (5%) <b>24</b> <b>27</b>	21, 38, 62, 83	0
All	All	644/660 (97%)	0.77	79 (12%) <b>4</b> <b>4</b>	20, 43, 82, 128	0

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	237	VAL	8.3
1	B	238	GLY	6.3
1	A	227	ALA	6.0
1	A	263	LEU	5.4
1	A	262	MET	4.9
1	A	176	ILE	4.9
1	B	280	GLU	4.9
1	B	236	MET	4.8
1	A	237	VAL	4.7
1	A	22	ILE	4.6
1	A	280	GLU	4.6
1	A	203	VAL	4.4
1	A	226	ILE	4.4
1	B	282	PHE	4.4
1	A	177	THR	4.1
1	A	205	GLY	3.8
1	A	204	ILE	3.7
1	B	285	MET	3.7
1	A	229	ASP	3.6
1	A	236	MET	3.4
1	A	261	ILE	3.4
1	A	225	TYR	3.4
1	A	146	LEU	3.3
1	B	235	GLY	3.3

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	283	ALA	3.3
1	A	289	TRP	3.2
1	B	22	ILE	3.2
1	A	264	SER	3.2
1	A	42	PHE	3.2
1	A	231	PHE	3.1
1	A	291	VAL	3.1
1	B	175	ILE	3.0
1	A	52	VAL	2.9
1	A	238	GLY	2.9
1	A	179	THR	2.8
1	A	295	PHE	2.8
1	A	70	THR	2.7
1	A	294	LEU	2.7
1	A	67	VAL	2.7
1	A	178	HIS	2.7
1	A	298	ILE	2.6
1	A	285	MET	2.6
1	B	42	PHE	2.6
1	A	144	ILE	2.6
1	A	282	PHE	2.6
1	A	243	GLU	2.6
1	B	16	GLN	2.6
1	B	278	LEU	2.5
1	A	296	VAL	2.5
1	A	16	GLN	2.4
1	A	232	GLY	2.4
1	A	224	VAL	2.4
1	A	21	LEU	2.4
1	A	60	ARG	2.4
1	B	279	PRO	2.4
1	A	93	ILE	2.4
1	B	146	LEU	2.3
1	A	74	CYS	2.3
1	A	265	HIS	2.3
1	A	88	THR	2.3
1	A	45	ASP	2.3
1	A	311	VAL	2.2
1	A	305	GLU	2.2
1	A	228	PHE	2.2
1	A	211	THR	2.2
1	B	43	ARG	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	95	ALA	2.2
1	A	268	VAL	2.2
1	B	176	ILE	2.1
1	A	49	ARG	2.1
1	A	147	ALA	2.1
1	A	69	PRO	2.1
1	B	284	GLU	2.1
1	A	323	LEU	2.1
1	A	207	MET	2.1
1	A	287	LYS	2.1
1	A	266	ASP	2.1
1	A	23	HIS	2.0
1	A	247	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	KCX	A	145	12/13	0.93	0.23	29,32,44,45	0
1	KCX	B	145	12/13	0.94	0.21	20,23,30,39	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	OH	A	403	1/1	0.91	0.38	41,41,41,41	0
4	OH	B	403	1/1	0.96	0.25	34,34,34,34	0
2	FE	A	401	1/1	0.97	0.11	29,29,29,29	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ZN	A	402	1/1	0.99	0.12	33,33,33,33	1
3	ZN	B	402	1/1	0.99	0.10	30,30,30,30	1
2	FE	B	401	1/1	1.00	0.11	23,23,23,23	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.