



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 18, 2024 – 10:34 PM EST

PDB ID : 4I1B
Title : FUNCTIONAL IMPLICATIONS OF INTERLEUKIN-1BETA BASED ON THE THREE-DIMENSIONAL STRUCTURE
Authors : Veerapandian, B.; Poulos, T.L.; Gilliland, G.L.; Raag, R.; Svensson, L.A.; Masui, Y.; Hirai, Y.
Deposited on : 1990-03-27
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	153	54% 33% 12% .

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 1298 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INTERLEUKIN-1 BETA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	151	1207	765	199	235	8	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	91	91	91	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: INTERLEUKIN-1 BETA

Chain A:  54% 33% 12%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	55.14Å 55.14Å 76.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.00	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.00)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	RESTRAIN	Depositor
R, R_{free}	0.190 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1298	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.02	1/1229 (0.1%)	1.46	21/1651 (1.3%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	141	GLN	CA-CB	-5.22	1.42	1.53

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	54	ASP	CB-CG-OD1	9.08	126.48	118.30
1	A	95	MET	CG-SD-CE	8.66	114.06	100.20
1	A	152	SER	C-N-CA	8.60	143.20	121.70
1	A	85	VAL	CA-CB-CG2	8.42	123.53	110.90
1	A	20	MET	CG-SD-CE	7.10	111.57	100.20
1	A	44	MET	CG-SD-CE	6.89	111.22	100.20
1	A	141	GLN	N-CA-CB	6.76	122.76	110.60
1	A	60	LEU	CB-CG-CD1	6.73	122.44	111.00
1	A	148	MET	CG-SD-CE	6.38	110.41	100.20
1	A	100	VAL	CA-CB-CG2	6.27	120.31	110.90
1	A	19	VAL	CA-CB-CG2	5.88	119.71	110.90
1	A	58	VAL	CA-CB-CG2	5.87	119.71	110.90
1	A	40	VAL	CA-CB-CG2	5.83	119.65	110.90
1	A	130	MET	CG-SD-CE	5.66	109.26	100.20
1	A	106	ILE	C-N-CA	5.65	135.83	121.70
1	A	141	GLN	CA-CB-CG	5.53	125.57	113.40
1	A	43	SER	N-CA-CB	5.50	118.75	110.50
1	A	141	GLN	CB-CA-C	5.47	121.34	110.40
1	A	3	VAL	CA-CB-CG2	5.41	119.02	110.90
1	A	37	GLU	OE1-CD-OE2	-5.25	117.00	123.30
1	A	47	VAL	CA-CB-CG2	5.08	118.52	110.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1207	0	1202	35	0
2	A	91	0	0	1	0
All	All	1298	0	1202	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:106:ILE:HG22	1:A:106:ILE:O	1.60	0.96
1:A:106:ILE:O	1:A:106:ILE:CG2	2.26	0.83
1:A:21:SER:HB2	1:A:27:LYS:HG3	1.66	0.78
1:A:23:PRO:HB2	1:A:24:TYR:CD1	2.21	0.76
1:A:109:LYS:HD3	1:A:145:ASP:OD1	1.92	0.70
1:A:21:SER:HB2	1:A:27:LYS:HD2	1.78	0.64
1:A:29:LEU:HD12	1:A:129:ASN:ND2	2.12	0.64
1:A:27:LYS:HE3	1:A:129:ASN:O	1.98	0.63
1:A:21:SER:HB2	1:A:27:LYS:CG	2.33	0.59
1:A:23:PRO:HB2	1:A:24:TYR:CE1	2.39	0.57
1:A:21:SER:CB	1:A:27:LYS:HG3	2.37	0.55
1:A:121:TYR:CE1	1:A:138:LYS:HD2	2.43	0.54
1:A:21:SER:HB2	1:A:27:LYS:CD	2.38	0.53
1:A:47:VAL:HG11	1:A:95:MET:HG3	1.91	0.53
1:A:121:TYR:CZ	1:A:138:LYS:HD2	2.44	0.53
1:A:19:VAL:HA	1:A:40:VAL:HG23	1.92	0.52
1:A:21:SER:OG	1:A:22:GLY:N	2.42	0.51
1:A:97:LYS:HE3	1:A:116:GLN:HA	1.93	0.51
1:A:78:PRO:HB3	1:A:117:PHE:CD2	2.48	0.49
1:A:8:CYS:HA	1:A:149:GLN:O	2.14	0.48
1:A:139:GLY:O	1:A:141:GLN:HG2	2.13	0.48
1:A:4:ARG:O	1:A:46:PHE:HD2	1.96	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:45:SER:O	1:A:58:VAL:HA	2.14	0.47
1:A:119:ASN:ND2	2:A:243:HOH:O	2.47	0.47
1:A:6:LEU:HA	1:A:6:LEU:HD12	1.78	0.47
1:A:7:ASN:ND2	1:A:153:SER:HB2	2.31	0.46
1:A:77:LYS:HE2	1:A:77:LYS:HB3	1.66	0.45
1:A:34:GLN:O	1:A:34:GLN:HG2	2.02	0.45
1:A:134:LEU:HD12	1:A:134:LEU:HA	1.90	0.44
1:A:7:ASN:ND2	1:A:153:SER:CB	2.80	0.44
1:A:55:LYS:HG2	1:A:102:ASN:HB3	2.00	0.43
1:A:34:GLN:O	1:A:34:GLN:CG	2.66	0.42
1:A:126:GLN:HE21	1:A:143:ILE:HD11	1.85	0.42
1:A:52:SER:OG	1:A:53:ASN:N	2.54	0.40
1:A:71:CYS:O	1:A:98:ARG:HB2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	149/153 (97%)	131 (88%)	15 (10%)	3 (2%)	7 3

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	75	ASP
1	A	107	ASN
1	A	139	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	139/140 (99%)	116 (84%)	23 (16%)	2 1

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	VAL
1	A	10	LEU
1	A	20	MET
1	A	21	SER
1	A	23	PRO
1	A	29	LEU
1	A	34	GLN
1	A	35	ASP
1	A	36	MET
1	A	39	GLN
1	A	50	GLU
1	A	51	GLU
1	A	54	ASP
1	A	63	LYS
1	A	64	GLU
1	A	72	VAL
1	A	74	LYS
1	A	77	LYS
1	A	88	LYS
1	A	107	ASN
1	A	126	GLN
1	A	129	ASN
1	A	137	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	ASN
1	A	48	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	119	ASN
1	A	126	GLN
1	A	129	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.