



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 29, 2024 – 07:47 PM EDT

PDB ID : 4IYP  
Title : structure of the nPP2Ac-alpha4 complex  
Authors : Jiang, L.; Stanevich, V.; Satyshur, K.A.; Xing, Y.  
Deposited on : 2013-01-29  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3657 (2.80-2.80)
Clashscore	180529	4123 (2.80-2.80)
Ramachandran outliers	177936	4071 (2.80-2.80)
Sidechain outliers	177891	4073 (2.80-2.80)
RSRZ outliers	164620	3659 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	222	 78% 13% 8%
2	C	151	 55% 23% 21%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2708 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Immunoglobulin-binding protein 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	205	1679	1057	293	322	2	5	0	0	0

There are 17 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP P78318
A	0	SER	-	expression tag	UNP P78318
A	1	MSE	-	expression tag	UNP P78318
A	?	-	LYS	deletion	UNP P78318
A	?	-	THR	deletion	UNP P78318
A	?	-	MET	deletion	UNP P78318
A	?	-	ASN	deletion	UNP P78318
A	?	-	ASN	deletion	UNP P78318
A	?	-	SER	deletion	UNP P78318
A	?	-	ALA	deletion	UNP P78318
A	?	-	GLU	deletion	UNP P78318
A	?	-	ASN	deletion	UNP P78318
A	?	-	HIS	deletion	UNP P78318
A	?	-	THR	deletion	UNP P78318
A	?	-	ALA	deletion	UNP P78318
A	?	-	ASN	deletion	UNP P78318
A	?	-	SER	deletion	UNP P78318

- Molecule 2 is a protein called Serine/threonine-protein phosphatase 2A catalytic subunit alpha isoform.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
2	C	119	995	643	158	189	3	2	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	3	GLY	-	expression tag	UNP P67775
C	4	SER	-	expression tag	UNP P67775
C	5	MSE	-	expression tag	UNP P67775


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	27	Total O 27 27	0	0
3	C	7	Total O 7 7	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Immunoglobulin-binding protein 1



- Molecule 2: Serine/threonine-protein phosphatase 2A catalytic subunit alpha isoform



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	116.13Å 116.13Å 81.23Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	34.43 – 2.80 34.43 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (34.43-2.80) 99.5 (34.43-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.66 (at 2.81Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, $R_{free}$	0.179 , 0.222 0.187 , 0.235	Depositor DCC
$R_{free}$ test set	796 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	73.1	Xtrriage
Anisotropy	0.100	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 69.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.033 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2708	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	84.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/1700	0.61	1/2279 (0.0%)
2	C	0.43	0/1008	0.63	1/1348 (0.1%)
All	All	0.45	0/2708	0.62	2/3627 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	7	LEU	CA-CB-CG	7.28	132.04	115.30
2	C	108	ARG	NE-CZ-NH1	5.14	122.87	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1679	0	1682	16	0
2	C	995	0	961	15	0
3	A	27	0	0	0	0
3	C	7	0	0	0	0
All	All	2708	0	2643	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:83:MSE:HE1	2:C:100:LEU:HB2	1.53	0.90
2:C:83:MSE:HG2	2:C:92:TYR:HE1	1.57	0.68
2:C:66:MSE:HG2	2:C:67:GLU:N	2.21	0.56
2:C:108:ARG:CG	2:C:108:ARG:HH11	2.22	0.53
1:A:159:ILE:HG12	2:C:40:THR:HG22	1.91	0.53
2:C:83:MSE:HG2	2:C:92:TYR:CE1	2.42	0.53
2:C:66:MSE:HG2	2:C:67:GLU:H	1.77	0.50
1:A:68:LEU:HD12	1:A:71:ILE:HD12	1.94	0.48
1:A:175:MSE:HE3	1:A:188:ARG:HD2	1.95	0.48
1:A:12:LEU:HB3	1:A:13:PRO:HD3	1.95	0.47
1:A:161:ARG:HH21	1:A:206:GLU:HG3	1.80	0.46
1:A:71:ILE:HD11	1:A:119:TYR:CZ	2.51	0.46
1:A:218:LEU:HD12	1:A:218:LEU:HA	1.76	0.46
1:A:184:ALA:HB1	1:A:188:ARG:HB3	1.98	0.46
2:C:39:LEU:HD23	2:C:39:LEU:HA	1.85	0.44
2:C:61:GLN:HE21	2:C:99:LEU:HD23	1.83	0.44
1:A:88:LEU:HA	1:A:88:LEU:HD23	1.77	0.44
1:A:54:GLU:O	1:A:58:GLN:HG2	2.19	0.43
1:A:32:PRO:HG2	1:A:35:SER:HB3	2.00	0.43
2:C:107:TYR:HB3	2:C:110:ARG:HB2	2.00	0.43
2:C:17:LEU:HD11	2:C:98:THR:HG22	1.99	0.43
1:A:205:LEU:HD23	1:A:205:LEU:HA	1.90	0.42
1:A:110:ILE:HD13	1:A:110:ILE:HA	1.86	0.42
1:A:82:PRO:HB2	1:A:112:TYR:HB2	2.02	0.41
2:C:24:SER:O	2:C:28:VAL:HG23	2.20	0.41
2:C:36:LYS:HD3	2:C:152:TYR:CE2	2.55	0.41
1:A:23:LEU:HD13	1:A:88:LEU:HG	2.01	0.41
1:A:37:ILE:H	1:A:37:ILE:HD12	1.85	0.41
2:C:83:MSE:CE	2:C:100:LEU:HB2	2.37	0.40
2:C:108:ARG:HH11	2:C:108:ARG:HG3	1.85	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	201/222 (90%)	196 (98%)	5 (2%)	0	100	100
2	C	107/151 (71%)	93 (87%)	11 (10%)	3 (3%)	4	14
All	All	308/373 (83%)	289 (94%)	16 (5%)	3 (1%)	13	39

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	87	VAL
2	C	85	ASP
2	C	116	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	183/191 (96%)	171 (93%)	12 (7%)	14	39
2	C	108/135 (80%)	92 (85%)	16 (15%)	2	8
All	All	291/326 (89%)	263 (90%)	28 (10%)	7	22

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	LEU
1	A	9	LEU
1	A	11	ARG
1	A	12	LEU
1	A	104	ARG
1	A	148	LEU
1	A	155	ARG
1	A	177	SER
1	A	188	ARG
1	A	194	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	195	LEU
1	A	218	LEU
2	C	7	THR
2	C	8	LYS
2	C	26	SER
2	C	42	GLU
2	C	63	HIS
2	C	68	LEU
2	C	86	TYR
2	C	97	VAL
2	C	106	ARG
2	C	108	ARG
2	C	123	ILE
2	C	124	THR
2	C	126	VAL
2	C	127	TYR
2	C	131	ASP
2	C	153	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	200/222 (90%)	-0.57	2 (1%) 79 73	38, 68, 118, 155	0
2	C	117/151 (77%)	0.15	11 (9%) 15 12	61, 98, 141, 159	0
All	All	317/373 (84%)	-0.30	13 (4%) 42 34	38, 82, 134, 159	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	150	ALA	5.4
2	C	57	ASP	4.7
2	C	55	CYS	4.6
2	C	69	PHE	3.2
2	C	6	PHE	3.1
2	C	60	GLY	2.9
1	A	148	LEU	2.6
2	C	123	ILE	2.6
2	C	127	TYR	2.2
2	C	85	ASP	2.1
2	C	7	THR	2.1
2	C	86	TYR	2.1
2	C	67	GLU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.