



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 13, 2024 – 09:37 AM EST

PDB ID : 4J3C  
Title : Crystal structure of 16S ribosomal RNA methyltransferase RsmE  
Authors : Eswaramoorthy, S.; Chamala, S.; Evans, B.; Foti, F.; Gizzi, A.; Hillerich, B.; Kar, A.; Laffleur, J.; Seidel, R.; Villigas, G.; Zencheck, W.; Al Obaidi, N.; Almo, S.C.; Swaminathan, S.; New York Structural Genomics Research Consortium (NYSGRC)  
Deposited on : 2013-02-05  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	
1	B	267	

## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 3612 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribosomal RNA small subunit methyltransferase E.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	232	1766	1118	317	323	3	5	0	0	0
1	B	234	1786	1129	324	325	3	5	0	0	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MSE	-	expression tag	UNP Q92RS8
A	-20	HIS	-	expression tag	UNP Q92RS8
A	-19	HIS	-	expression tag	UNP Q92RS8
A	-18	HIS	-	expression tag	UNP Q92RS8
A	-17	HIS	-	expression tag	UNP Q92RS8
A	-16	HIS	-	expression tag	UNP Q92RS8
A	-15	HIS	-	expression tag	UNP Q92RS8
A	-14	SER	-	expression tag	UNP Q92RS8
A	-13	SER	-	expression tag	UNP Q92RS8
A	-12	GLY	-	expression tag	UNP Q92RS8
A	-11	VAL	-	expression tag	UNP Q92RS8
A	-10	ASP	-	expression tag	UNP Q92RS8
A	-9	LEU	-	expression tag	UNP Q92RS8
A	-8	GLY	-	expression tag	UNP Q92RS8
A	-7	THR	-	expression tag	UNP Q92RS8
A	-6	GLU	-	expression tag	UNP Q92RS8
A	-5	ASN	-	expression tag	UNP Q92RS8
A	-4	LEU	-	expression tag	UNP Q92RS8
A	-3	TYR	-	expression tag	UNP Q92RS8
A	-2	PHE	-	expression tag	UNP Q92RS8
A	-1	GLN	-	expression tag	UNP Q92RS8
A	0	SER	-	expression tag	UNP Q92RS8
B	-21	MSE	-	expression tag	UNP Q92RS8
B	-20	HIS	-	expression tag	UNP Q92RS8
B	-19	HIS	-	expression tag	UNP Q92RS8

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-18	HIS	-	expression tag	UNP Q92RS8
B	-17	HIS	-	expression tag	UNP Q92RS8
B	-16	HIS	-	expression tag	UNP Q92RS8
B	-15	HIS	-	expression tag	UNP Q92RS8
B	-14	SER	-	expression tag	UNP Q92RS8
B	-13	SER	-	expression tag	UNP Q92RS8
B	-12	GLY	-	expression tag	UNP Q92RS8
B	-11	VAL	-	expression tag	UNP Q92RS8
B	-10	ASP	-	expression tag	UNP Q92RS8
B	-9	LEU	-	expression tag	UNP Q92RS8
B	-8	GLY	-	expression tag	UNP Q92RS8
B	-7	THR	-	expression tag	UNP Q92RS8
B	-6	GLU	-	expression tag	UNP Q92RS8
B	-5	ASN	-	expression tag	UNP Q92RS8
B	-4	LEU	-	expression tag	UNP Q92RS8
B	-3	TYR	-	expression tag	UNP Q92RS8
B	-2	PHE	-	expression tag	UNP Q92RS8
B	-1	GLN	-	expression tag	UNP Q92RS8
B	0	SER	-	expression tag	UNP Q92RS8


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	42	Total O 42 42	0	0
2	B	18	Total O 18 18	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ribosomal RNA small subunit methyltransferase E



- Molecule 1: Ribosomal RNA small subunit methyltransferase E



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.08Å 152.08Å 38.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.00 50.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.2 (50.00-2.00) 99.3 (50.00-2.00)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.01 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.5.0110	Depositor
R, $R_{free}$	0.246 , 0.287 0.256 , 0.294	Depositor DCC
$R_{free}$ test set	1979 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.8	Xtrriage
Anisotropy	0.547	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 39.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3612	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.74% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.72	1/1790 (0.1%)	0.76	1/2419 (0.0%)
1	B	0.61	0/1810	0.72	0/2448
All	All	0.67	1/3600 (0.0%)	0.74	1/4867 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	139	CYS	CB-SG	5.34	1.91	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	243	ASP	CB-CG-OD1	5.13	122.92	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1766	0	1778	34	0
1	B	1786	0	1798	27	0
2	A	42	0	0	2	0
2	B	18	0	0	0	0
All	All	3612	0	3576	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:114:MSE:HE3	1:B:150:PRO:HB3	1.54	0.89
1:A:35:VAL:HG13	1:A:36:LEU:HA	1.61	0.83
1:B:98:LEU:CD2	1:B:227:ASP:HB3	2.09	0.81
1:A:225:ARG:HH22	1:B:100:GLN:HE21	1.30	0.79
1:A:35:VAL:HA	1:A:37:ARG:H	1.49	0.78
1:B:90:LEU:HB3	1:B:197:GLU:OE2	1.85	0.75
1:A:179:LEU:O	1:A:183:GLU:HG2	1.88	0.73
1:A:25:THR:C	1:A:27:GLU:H	1.93	0.72
1:B:98:LEU:HD23	1:B:227:ASP:HB3	1.70	0.71
1:A:52:GLY:H	1:A:77:GLN:HE21	1.38	0.71
1:A:35:VAL:HA	1:A:37:ARG:N	2.08	0.69
1:B:94:ARG:O	1:B:98:LEU:HG	1.94	0.68
1:A:28:GLN:O	1:A:29:PHE:HB2	1.93	0.67
1:B:97:TYR:CE2	1:B:101:LYS:HG3	2.29	0.67
1:B:112:PRO:HB3	1:B:122:ILE:HD11	1.81	0.62
1:A:29:PHE:HB2	1:A:34:ASN:N	2.14	0.62
1:A:44:LEU:HD11	1:A:46:VAL:HG13	1.79	0.62
1:B:55:ARG:NH2	1:B:57:GLU:OE2	2.32	0.62
1:A:25:THR:O	1:A:27:GLU:N	2.34	0.61
1:A:44:LEU:CD1	1:A:46:VAL:HG13	2.30	0.61
1:A:27:GLU:C	1:A:29:PHE:H	2.05	0.60
1:A:152:LYS:HE2	2:A:323:HOH:O	2.01	0.60
1:B:12:ILE:HG22	1:B:47:PHE:O	2.03	0.58
1:A:225:ARG:HH22	1:B:100:GLN:NE2	1.99	0.57
1:B:42:SER:HB2	1:B:58:ILE:HD12	1.86	0.56
1:A:12:ILE:HD12	1:A:14:ASN:HB2	1.88	0.55
1:A:44:LEU:C	1:A:44:LEU:HD12	2.27	0.55
1:A:27:GLU:C	1:A:29:PHE:N	2.60	0.54
1:B:17:HIS:HA	1:B:54:TRP:CH2	2.43	0.53
1:B:8:GLN:NE2	1:B:31:TYR:OH	2.41	0.53
1:A:29:PHE:HB3	1:A:35:VAL:O	2.09	0.52
1:A:27:GLU:O	1:A:29:PHE:N	2.44	0.51
1:B:53:GLU:OE1	1:B:75:ARG:HD3	2.11	0.51
1:A:62:SER:O	1:A:65:GLN:N	2.44	0.50
1:A:25:THR:C	1:A:27:GLU:N	2.63	0.50
1:B:97:TYR:CE2	1:B:101:LYS:CG	2.95	0.49
1:A:224:LEU:HD21	2:A:319:HOH:O	2.13	0.47

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:26:ARG:C	1:A:28:GLN:H	2.17	0.47
1:B:12:ILE:HG23	1:B:48:ASN:HB3	1.94	0.47
1:A:17:HIS:HA	1:A:54:TRP:CH2	2.49	0.47
1:A:53:GLU:OE1	1:A:75:ARG:HD3	2.16	0.46
1:A:122:ILE:HA	1:A:122:ILE:HD13	1.73	0.46
1:A:113:VAL:HG11	1:A:153:LEU:HD13	1.98	0.46
1:A:42:SER:HB2	1:A:58:ILE:HD12	1.98	0.46
1:B:32:LEU:HA	1:B:36:LEU:HD12	1.99	0.44
1:B:169:CYS:SG	1:B:219:LEU:HD11	2.56	0.44
1:A:44:LEU:HD11	1:A:46:VAL:CG1	2.46	0.44
1:B:33:ILE:O	1:B:37:ARG:HA	2.18	0.44
1:A:45:LEU:HD22	1:A:74:THR:HG21	2.00	0.43
1:A:44:LEU:CD1	1:A:44:LEU:C	2.86	0.43
1:B:12:ILE:HD13	1:B:12:ILE:HG21	1.76	0.43
1:A:170:ASP:OD1	1:A:171:GLU:N	2.52	0.42
1:A:28:GLN:O	1:A:29:PHE:CB	2.65	0.42
1:B:12:ILE:HD11	1:B:22:HIS:HB2	2.00	0.42
1:B:238:GLN:HB3	1:B:244:TRP:CD1	2.55	0.42
1:B:125:LEU:O	1:B:129:ARG:HG3	2.20	0.41
1:B:125:LEU:HD13	1:B:129:ARG:HH21	1.85	0.41
1:B:86:LEU:HD23	1:B:111:GLN:HB3	2.02	0.41
1:B:227:ASP:OD1	1:B:227:ASP:N	2.54	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/267 (84%)	213 (95%)	7 (3%)	4 (2%)	7	3
1	B	226/267 (85%)	220 (97%)	6 (3%)	0	100	100
All	All	450/534 (84%)	433 (96%)	13 (3%)	4 (1%)	14	10

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	26	ARG
1	A	27	GLU
1	A	61	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/211 (85%)	169 (94%)	11 (6%)	15	12
1	B	182/211 (86%)	177 (97%)	5 (3%)	40	42
All	All	362/422 (86%)	346 (96%)	16 (4%)	24	22

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	PHE
1	A	36	LEU
1	A	44	LEU
1	A	60	MSE
1	A	62	SER
1	A	68	LEU
1	A	116	GLN
1	A	141	VAL
1	A	188	ARG
1	A	210	SER
1	A	223	ILE
1	B	46	VAL
1	B	90	LEU
1	B	124	SER
1	B	145	PRO
1	B	151	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	HIS
1	A	77	GLN
1	A	111	GLN
1	A	116	GLN
1	B	8	GLN
1	B	100	GLN
1	B	111	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	227/267 (85%)	0.49	26 (11%)	11   10	20, 39, 75, 94	0
1	B	229/267 (85%)	1.02	37 (16%)	5   5	26, 48, 72, 79	0
All	All	456/534 (85%)	0.76	63 (13%)	8   6	20, 44, 73, 94	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	31	TYR	5.8
1	B	169	CYS	4.7
1	A	29	PHE	4.6
1	A	35	VAL	4.4
1	B	90	LEU	4.0
1	A	34	ASN	4.0
1	B	122	ILE	3.6
1	B	62	SER	3.4
1	A	172	GLY	3.4
1	B	92	VAL	3.3
1	B	125	LEU	3.3
1	B	159	ASP	3.2
1	A	63	ARG	3.2
1	A	59	ALA	3.1
1	A	25	THR	2.9
1	A	159	ASP	2.9
1	B	174	ASP	2.9
1	A	61	PRO	2.8
1	B	63	ARG	2.8
1	A	28	GLN	2.8
1	A	38	LEU	2.8
1	B	66	ALA	2.8
1	A	97	TYR	2.8
1	B	35	VAL	2.7

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	163	ASP	2.7
1	B	61	PRO	2.7
1	A	18	ALA	2.7
1	B	34	ASN	2.6
1	B	156	LEU	2.6
1	B	49	GLY	2.5
1	A	176	GLN	2.4
1	B	24	ALA	2.4
1	B	94	ARG	2.4
1	B	123	GLY	2.4
1	A	138	GLN	2.3
1	B	30	ASN	2.3
1	A	68	LEU	2.3
1	B	186	ALA	2.3
1	B	33	ILE	2.3
1	A	36	LEU	2.3
1	B	29	PHE	2.2
1	A	26	ARG	2.2
1	A	223	ILE	2.2
1	B	83	LEU	2.2
1	B	15	ALA	2.2
1	B	121	LYS	2.2
1	B	93	GLY	2.2
1	B	127	ARG	2.2
1	A	74	THR	2.1
1	B	175	SER	2.1
1	B	128	VAL	2.1
1	A	15	ALA	2.1
1	B	54	TRP	2.1
1	B	36	LEU	2.1
1	A	175	SER	2.1
1	A	67	VAL	2.1
1	A	92	VAL	2.1
1	B	225	ARG	2.1
1	B	38	LEU	2.0
1	B	12	ILE	2.0
1	A	19	GLY	2.0
1	A	24	ALA	2.0
1	B	79	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.