



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 13, 2023 – 07:35 AM EDT

PDB ID : 4QDQ  
Title : Physical basis for Nrp2 ligand binding  
Authors : Parker, M.W.; Vander Kooi, C.W.  
Deposited on : 2014-05-14  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	
1	B	329	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5444 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neuropilin-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	316	2547	1602	463	471	11	0	2	0
1	B	316	2536	1596	459	470	11	0	1	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	272	GLY	-	expression tag	UNP O60462
A	273	SER	-	expression tag	UNP O60462
A	274	HIS	-	expression tag	UNP O60462
A	275	MET	-	expression tag	UNP O60462
A	596	SER	-	expression tag	UNP O60462
A	597	ILE	-	expression tag	UNP O60462
A	598	ILE	-	expression tag	UNP O60462
A	599	ARG	-	expression tag	UNP O60462
A	600	ARG	-	expression tag	UNP O60462
B	272	GLY	-	expression tag	UNP O60462
B	273	SER	-	expression tag	UNP O60462
B	274	HIS	-	expression tag	UNP O60462
B	275	MET	-	expression tag	UNP O60462
B	596	SER	-	expression tag	UNP O60462
B	597	ILE	-	expression tag	UNP O60462
B	598	ILE	-	expression tag	UNP O60462
B	599	ARG	-	expression tag	UNP O60462
B	600	ARG	-	expression tag	UNP O60462

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	169	Total	O	0	0
			169	169		
4	B	158	Total	O	0	0
			158	158		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Neuropilin-2

Chain A: 



- Molecule 1: Neuropilin-2

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	41.05Å 120.81Å 69.84Å 90.00° 103.29° 90.00°	Depositor
Resolution (Å)	19.75 – 1.95 19.75 – 1.95	Depositor EDS
% Data completeness (in resolution range)	90.6 (19.75-1.95) 90.7 (19.75-1.95)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.69 (at 1.94Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.210 , 0.241 0.214 , 0.241	Depositor DCC
$R_{free}$ test set	2138 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.9	Xtrriage
Anisotropy	0.232	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 45.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5444	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.54	2/2610 (0.1%)	0.59	0/3540
1	B	0.53	3/2599 (0.1%)	0.61	0/3526
All	All	0.53	5/5209 (0.1%)	0.60	0/7066

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	487	TRP	CD2-CE2	5.22	1.47	1.41
1	A	304	TRP	CD2-CE2	5.20	1.47	1.41
1	B	594	TRP	CD2-CE2	5.19	1.47	1.41
1	A	594	TRP	CD2-CE2	5.07	1.47	1.41
1	B	535	TRP	CD2-CE2	5.06	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2547	0	2490	24	0
1	B	2536	0	2478	9	0
2	A	18	0	24	6	0
2	B	6	0	8	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	169	0	0	3	0
4	B	158	0	0	2	0
All	All	5444	0	5000	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:528:TYR:CG	1:A:566:ILE:HD12	2.22	0.75
1:A:528:TYR:CD2	1:A:566:ILE:HD12	2.25	0.72
1:A:481:ALA:O	1:A:578:TRP:CZ2	2.45	0.69
1:A:444:LEU:HD11	1:B:334:ARG:HD3	1.80	0.64
1:A:435:SER:O	1:A:590:LEU:HD22	2.01	0.60
1:A:481:ALA:O	1:A:578:TRP:HZ2	1.86	0.59
1:A:523:LYS:NZ	1:A:550:GLU:OE1	2.37	0.57
1:B:467:LEU:HG	1:B:468:VAL:HG23	1.84	0.57
1:A:340:THR:HG22	1:A:401:PRO:HB3	1.90	0.53
1:A:341:ALA:HB3	1:A:425:PHE:HB2	1.91	0.53
1:A:476:PRO:HG2	1:A:479:PRO:HA	1.91	0.52
2:A:702:GOL:H31	4:A:942:HOH:O	2.13	0.48
1:A:561:ARG:HD2	2:A:701:GOL:H11	1.96	0.47
1:A:526:VAL:HB	1:A:538:ILE:HD12	1.97	0.47
1:A:352:THR:O	1:A:353:GLN:HB2	2.14	0.47
1:A:567:PRO:HB3	1:A:594:TRP:NE1	2.29	0.47
1:A:400:ALA:HA	1:A:401:PRO:HD2	1.83	0.46
1:B:336:LEU:HD11	1:B:468:VAL:HG21	1.97	0.45
1:A:547:LYS:HE3	2:A:701:GOL:H2	1.98	0.45
1:A:355:GLY:H	2:A:702:GOL:H32	1.82	0.45
1:B:325:ASN:O	4:B:836:HOH:O	2.21	0.44
1:A:478:ILE:O	1:A:481:ALA:HB2	2.18	0.44
1:A:542:ARG:HA	1:A:542:ARG:HD2	1.88	0.44
1:A:338:MET:O	1:A:426:GLY:HA3	2.18	0.44
1:A:481:ALA:O	1:A:578:TRP:CH2	2.70	0.43
2:A:701:GOL:H12	4:A:903:HOH:O	2.18	0.43
1:A:410:ARG:HB3	4:A:845:HOH:O	2.17	0.43
1:B:410:ARG:HB3	4:B:836:HOH:O	2.19	0.42
1:A:354:ASN:HA	2:A:702:GOL:H32	2.01	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:453:SER:HB3	1:B:487:TRP:CD1	2.55	0.42
1:B:460:TRP:HZ2	1:B:465:ALA:HB2	1.85	0.41
1:B:338:MET:CE	1:B:401:PRO:HB2	2.51	0.40
1:B:373:MET:HB2	1:B:554:HIS:CE1	2.56	0.40
1:A:528:TYR:CG	1:A:566:ILE:CD1	3.00	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/329 (95%)	304 (97%)	10 (3%)	0	100	100
1	B	313/329 (95%)	302 (96%)	11 (4%)	0	100	100
All	All	627/658 (95%)	606 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	275/282 (98%)	275 (100%)	0	100	100
1	B	274/282 (97%)	274 (100%)	0	100	100
All	All	549/564 (97%)	549 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	B	701	-	5,5,5	0.27	0	5,5,5	0.30	0
3	SO4	B	702	-	4,4,4	0.39	0	6,6,6	0.17	0
3	SO4	A	704	-	4,4,4	0.32	0	6,6,6	0.13	0
2	GOL	A	702	-	5,5,5	0.33	0	5,5,5	0.22	0
2	GOL	A	703	-	5,5,5	0.26	0	5,5,5	0.27	0
2	GOL	A	701	-	5,5,5	0.40	0	5,5,5	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	703	-	-	4/4/4/4	-
2	GOL	A	701	-	-	2/4/4/4	-
2	GOL	A	702	-	-	2/4/4/4	-
2	GOL	B	701	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	702	GOL	O2-C2-C3-O3
2	A	701	GOL	C1-C2-C3-O3
2	A	702	GOL	C1-C2-C3-O3
2	A	703	GOL	O1-C1-C2-O2
2	A	703	GOL	O2-C2-C3-O3
2	A	703	GOL	O1-C1-C2-C3
2	A	703	GOL	C1-C2-C3-O3
2	A	701	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	702	GOL	3	0
2	A	701	GOL	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	316/329 (96%)	0.30	16 (5%) 28 37	18, 29, 49, 73	0
1	B	316/329 (96%)	0.42	28 (8%) 9 15	20, 30, 62, 75	0
All	All	632/658 (96%)	0.36	44 (6%) 16 24	18, 29, 54, 75	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	459	LEU	8.3
1	B	517	ALA	7.9
1	A	458	TYR	7.2
1	B	479	PRO	6.8
1	B	597	ILE	6.6
1	B	480	GLN	6.0
1	B	544	GLN	5.8
1	A	597	ILE	5.8
1	A	517	ALA	5.4
1	B	459	LEU	5.3
1	B	595	THR	5.0
1	A	595	THR	4.8
1	B	276	GLN	4.7
1	B	518	ARG	4.5
1	B	596	SER	4.2
1	A	420	LEU	4.1
1	B	542	ARG	3.8
1	A	276	GLN	3.3
1	B	460	TRP	3.2
1	A	542	ARG	3.2
1	B	420	LEU	2.9
1	B	598	ILE	2.7
1	B	335	PHE	2.6
1	B	481	ALA	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	541	PRO	2.6
1	A	594	TRP	2.6
1	A	379	LYS	2.6
1	B	431	ASP	2.6
1	A	418	ILE	2.5
1	B	478	ILE	2.5
1	B	521	VAL	2.5
1	B	545	GLN	2.4
1	B	399	HIS	2.4
1	A	596	SER	2.3
1	B	543	THR	2.3
1	B	419	ALA	2.2
1	B	577	ARG	2.2
1	A	433	PRO	2.1
1	A	599	ARG	2.1
1	B	318	TRP	2.1
1	A	378	GLY	2.1
1	B	553	MET	2.0
1	A	435	SER	2.0
1	B	304	TRP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GOL	A	701	6/6	0.61	0.30	48,54,55,57	0
2	GOL	A	702	6/6	0.70	0.22	67,70,72,74	0
2	GOL	B	701	6/6	0.72	0.22	61,67,71,75	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GOL	A	703	6/6	0.81	0.19	52,54,56,60	0
3	SO4	B	702	5/5	0.91	0.19	61,62,65,68	0
3	SO4	A	704	5/5	0.95	0.13	68,69,74,74	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.