



Full wwPDB X-ray Structure Validation Report ⓘ

May 3, 2025 – 09:47 AM EDT

PDB ID : 4R6N / pdb_00004r6n
Title : Jacalin-carbohydrate interactions. Distortion of the ligand as a determinant of affinity
Authors : Abhinav, K.V.; Sharma, K.; Swaminathan, C.P.; Surolia, A.; Vijayan, M.
Deposited on : 2014-08-26
Resolution : 1.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	8422 (1.70-1.66)
Clashscore	180529	1005 (1.68-1.68)
Ramachandran outliers	177936	9065 (1.70-1.66)
Sidechain outliers	177891	9064 (1.70-1.66)
RSRZ outliers	164620	8421 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	133	 89% 11%
1	C	133	 87% 12%
1	E	133	 87% 13%
1	G	133	 87% 12%
2	B	19	 53% 32% 16%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	19	
2	F	19	
2	H	19	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	IPA	A	205	-	-	X	-
4	IPA	G	203	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5367 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Agglutinin alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	133	Total	C	N	O	S	0	0	0
			1036	677	159	198	2			
1	C	133	Total	C	N	O	S	0	0	0
			1040	679	160	199	2			
1	E	133	Total	C	N	O	S	0	0	0
			1040	679	160	199	2			
1	G	133	Total	C	N	O	S	0	0	0
			1037	678	159	198	2			

- Molecule 2 is a protein called Agglutinin beta-3 chain.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	16	Total	C	N	O	0	0	0
			114	73	20	21			
2	D	16	Total	C	N	O	0	1	0
			117	75	20	22			
2	F	16	Total	C	N	O	0	1	0
			119	77	21	21			
2	H	15	Total	C	N	O	0	0	0
			105	68	18	19			

- Molecule 3 is methyl beta-D-galactopyranoside (CCD ID: MBG) (formula: C₇H₁₄O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	7	6		
3	E	1	Total	C	O	0	0
			13	7	6		
3	G	1	Total	C	O	0	0
			13	7	6		

- Molecule 4 is ISOPROPYL ALCOHOL (CCD ID: IPA) (formula: C_3H_8O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	3	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	3	1		
4	A	1	Total	C	O	0	0
			4	3	1		
4	A	1	Total	C	O	0	0
			4	3	1		
4	C	1	Total	C	O	0	0
			4	3	1		
4	C	1	Total	C	O	0	0
			4	3	1		
4	C	1	Total	C	O	0	0
			4	3	1		
4	C	1	Total	C	O	0	0
			4	3	1		
4	D	1	Total	C	O	0	0
			4	3	1		
4	D	1	Total	C	O	0	0
			4	3	1		
4	E	1	Total	C	O	0	0
			4	3	1		
4	E	1	Total	C	O	0	0
			4	3	1		
4	E	1	Total	C	O	0	0
			4	3	1		
4	E	1	Total	C	O	0	0
			4	3	1		
4	G	1	Total	C	O	0	0
			4	3	1		
4	G	1	Total	C	O	0	0
			4	3	1		
4	H	1	Total	C	O	0	0
			4	3	1		
4	H	1	Total	C	O	0	0
			4	3	1		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0
5	D	1	Total C O 4 2 2	0	0
5	E	1	Total C O 4 2 2	0	0
5	E	1	Total C O 4 2 2	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	163	Total O 163 163	0	0
6	B	14	Total O 14 14	0	0
6	C	138	Total O 138 138	0	0
6	D	12	Total O 12 12	0	0
6	E	139	Total O 139 139	0	0
6	F	15	Total O 15 15	0	0
6	G	130	Total O 130 130	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	17	Total	O	0	0
			17	17		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Agglutinin alpha chain

Chain A: 



- Molecule 1: Agglutinin alpha chain

Chain C: 



- Molecule 1: Agglutinin alpha chain

Chain E: 




- Molecule 1: Agglutinin alpha chain

Chain G: 



- Molecule 2: Agglutinin beta-3 chain

Chain B: 



- Molecule 2: Agglutinin beta-3 chain

Chain D: 



● Molecule 2: Agglutinin beta-3 chain



● Molecule 2: Agglutinin beta-3 chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.53Å 80.90Å 63.21Å 90.00° 108.08° 90.00°	Depositor
Resolution (Å)	60.09 – 1.67 60.09 – 1.67	Depositor EDS
% Data completeness (in resolution range)	96.3 (60.09-1.67) 96.3 (60.09-1.67)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 1.67Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.170 , 0.221 0.170 , 0.220	Depositor DCC
R_{free} test set	3158 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	12.0	Xtriage
Anisotropy	0.248	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 50.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5367	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IPA, MBG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.29	0/1065	1.16	0/1444
1	C	1.25	1/1069 (0.1%)	1.08	0/1449
1	E	1.30	4/1069 (0.4%)	1.14	0/1449
1	G	1.26	1/1066 (0.1%)	1.17	2/1445 (0.1%)
2	B	1.24	0/116	1.11	0/157
2	D	1.28	0/122	1.31	1/165 (0.6%)
2	F	1.25	0/120	1.27	1/160 (0.6%)
2	H	1.28	0/107	1.20	0/145
All	All	1.27	6/4734 (0.1%)	1.15	4/6414 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	84	LEU	CA-C	-7.13	1.44	1.52
1	C	113	ILE	N-CA	5.34	1.52	1.46
1	E	91	LYS	N-CA	5.21	1.52	1.45
1	E	38	PRO	C-O	5.20	1.29	1.23
1	E	44	HIS	CA-C	-5.14	1.46	1.52
1	G	26	GLY	N-CA	5.13	1.52	1.45

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	10	THR	CB-CA-C	-8.80	95.39	110.09
1	G	111	GLY	CA-C-O	-6.60	117.55	122.37
2	D	8	GLN	N-CA-C	5.38	118.04	109.81
2	F	4	SER	N-CA-C	5.00	117.27	109.52

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1036	0	1003	16	0
1	C	1040	0	1009	15	0
1	E	1040	0	1009	13	0
1	G	1037	0	1005	15	0
2	B	114	0	116	5	0
2	D	117	0	121	12	0
2	F	119	0	123	3	0
2	H	105	0	108	1	0
3	A	13	0	14	0	0
3	E	13	0	14	0	0
3	G	13	0	14	0	0
4	A	16	0	32	9	0
4	C	16	0	32	4	0
4	D	8	0	16	5	0
4	E	16	0	32	7	0
4	G	8	0	16	6	0
4	H	8	0	16	1	0
5	A	4	0	6	0	0
5	C	4	0	6	0	0
5	D	4	0	6	3	0
5	E	8	0	12	0	0
6	A	163	0	0	5	0
6	B	14	0	0	0	0
6	C	138	0	0	9	0
6	D	12	0	0	1	0
6	E	139	0	0	3	0
6	F	15	0	0	0	0
6	G	130	0	0	1	0
6	H	17	0	0	0	0
All	All	5367	0	4710	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:102:IPA:H12	1:G:35:ASN:OD1	1.74	0.86
2:D:16:GLY:H	5:D:103:EDO:H11	1.48	0.79
1:C:126:TYR:HD1	6:C:432:HOH:O	1.65	0.79
4:G:202:IPA:H11	6:G:341:HOH:O	1.85	0.76
1:G:82:ARG:HH11	4:G:203:IPA:H13	1.56	0.70
4:A:205:IPA:H11	6:E:340:HOH:O	1.92	0.68
1:A:34:LEU:HD23	4:A:205:IPA:H33	1.77	0.67
1:E:94:GLY:H	4:E:203:IPA:C1	2.08	0.66
1:C:127:PHE:N	6:C:432:HOH:O	2.30	0.64
6:A:455:HOH:O	2:D:18:LYS:HD3	2.00	0.62
1:A:35:ASN:HB2	4:A:205:IPA:H13	1.82	0.61
1:G:73:GLY:HA3	4:G:203:IPA:H11	1.84	0.60
2:D:8:GLN:HB2	4:D:101:IPA:O2	2.01	0.60
2:B:15:TRP:HZ2	6:C:389:HOH:O	1.85	0.59
1:E:110:ASN:ND2	4:H:101:IPA:H31	2.18	0.58
1:A:91:LYS:HE3	6:A:424:HOH:O	2.04	0.58
1:E:94:GLY:H	4:E:203:IPA:H13	1.69	0.57
1:C:14:GLU:OE2	1:C:55:LYS:NZ	2.38	0.57
1:A:90:LYS:NZ	6:A:402:HOH:O	2.37	0.56
1:E:16:ASN:HB3	4:E:205:IPA:H12	1.87	0.55
1:C:72:THR:HG21	2:D:16:GLY:HA2	1.89	0.54
6:C:432:HOH:O	2:D:14:PRO:HB3	2.07	0.54
2:B:3:GLN:HB3	6:E:416:HOH:O	2.07	0.54
4:A:203:IPA:C3	1:E:10:THR:HG23	2.39	0.53
2:D:16:GLY:H	5:D:103:EDO:C1	2.20	0.53
1:E:17:LEU:HD12	1:E:17:LEU:C	2.34	0.53
4:E:205:IPA:H33	6:E:401:HOH:O	2.08	0.52
1:C:20:ASN:HB3	1:C:23:THR:HG1	1.73	0.52
1:A:91:LYS:CE	6:A:424:HOH:O	2.58	0.51
1:G:82:ARG:HH11	4:G:203:IPA:C1	2.22	0.51
1:E:94:GLY:H	4:E:203:IPA:H11	1.74	0.51
1:A:42:GLN:CB	6:A:453:HOH:O	2.59	0.50
1:C:20:ASN:HB3	1:C:23:THR:OG1	2.11	0.50
1:C:17:LEU:HG	1:C:54:VAL:CG2	2.41	0.50
2:D:11:ILE:HD11	4:D:101:IPA:H33	1.93	0.50
1:C:51:PHE:O	4:C:201:IPA:H31	2.12	0.49
4:C:203:IPA:H33	1:G:8:ALA:O	2.12	0.49
1:E:127:PHE:CE2	1:E:129:MET:HE2	2.48	0.49
1:C:127:PHE:CE2	1:C:129:MET:HE2	2.48	0.49
1:A:117:LYS:HE3	1:C:109:GLU:OE2	2.12	0.48
1:G:12:ILE:HD11	1:G:113:ILE:HG22	1.95	0.48
1:C:126:TYR:CD1	6:C:432:HOH:O	2.51	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:40:VAL:H	4:E:202:IPA:H11	1.77	0.48
1:G:10:THR:HG22	1:G:10:THR:O	2.13	0.48
1:C:10:THR:HG23	4:C:203:IPA:H32	1.96	0.48
1:G:82:ARG:HD2	4:G:203:IPA:H13	1.97	0.47
1:A:34:LEU:O	4:A:205:IPA:H32	2.14	0.47
1:G:82:ARG:CD	4:G:203:IPA:H13	2.45	0.46
6:C:432:HOH:O	2:D:14:PRO:CB	2.63	0.46
2:D:16:GLY:O	2:D:17:ALA:C	2.56	0.46
4:D:101:IPA:H2	6:D:207:HOH:O	2.16	0.46
1:A:133:LEU:HD13	2:D:8:GLN:HG3	1.98	0.45
1:A:47:PHE:HE2	4:A:204:IPA:O2	1.99	0.45
2:F:18[B]:LYS:O	2:F:18[B]:LYS:HG2	2.13	0.45
1:C:127:PHE:CA	6:C:432:HOH:O	2.64	0.45
1:E:40:VAL:H	4:E:202:IPA:C1	2.29	0.45
1:E:79:VAL:O	2:F:17:ALA:HB2	2.16	0.45
1:A:20:ASN:HB3	1:A:23:THR:OG1	2.17	0.44
1:A:79:VAL:O	2:B:17:ALA:HB2	2.16	0.44
2:B:8:GLN:NE2	2:H:8:GLN:HE22	2.15	0.44
2:F:17:ALA:O	2:F:18[A]:LYS:HB3	2.16	0.44
1:E:83:SER:HA	1:E:96:TYR:O	2.18	0.43
1:C:21:LYS:CE	6:C:333:HOH:O	2.65	0.43
1:G:39:TYR:C	1:G:39:TYR:CD1	2.96	0.43
2:D:16:GLY:N	5:D:103:EDO:H11	2.23	0.43
1:G:43:ASN:ND2	1:G:45:LYS:HD3	2.33	0.43
4:C:202:IPA:H32	6:C:404:HOH:O	2.18	0.43
1:A:35:ASN:HB2	4:A:205:IPA:C1	2.49	0.43
1:A:34:LEU:HD23	4:A:205:IPA:C3	2.45	0.42
2:D:6:ILE:HG23	1:G:10:THR:CG2	2.49	0.42
1:G:71:TYR:CE1	1:G:99:THR:HB	2.54	0.42
4:D:102:IPA:C1	1:G:35:ASN:OD1	2.58	0.42
1:A:85:THR:OG1	4:A:202:IPA:H32	2.19	0.42
2:B:18:LYS:HD3	2:B:18:LYS:HA	1.79	0.42
1:A:4:PHE:HB2	1:A:44:HIS:CE1	2.55	0.41
1:E:58:LEU:HD22	1:E:88:THR:HG21	2.03	0.41
1:C:18:SER:HB2	1:C:51:PHE:HB3	2.02	0.40
1:G:4:PHE:HB2	1:G:44:HIS:CE1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/133 (98%)	127 (97%)	4 (3%)	0	100	100
1	C	131/133 (98%)	125 (95%)	6 (5%)	0	100	100
1	E	131/133 (98%)	128 (98%)	3 (2%)	0	100	100
1	G	131/133 (98%)	126 (96%)	5 (4%)	0	100	100
2	B	14/19 (74%)	14 (100%)	0	0	100	100
2	D	15/19 (79%)	13 (87%)	2 (13%)	0	100	100
2	F	14/19 (74%)	14 (100%)	0	0	100	100
2	H	13/19 (68%)	13 (100%)	0	0	100	100
All	All	580/608 (95%)	560 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	112/113 (99%)	110 (98%)	2 (2%)	54	36
1	C	113/113 (100%)	111 (98%)	2 (2%)	54	36
1	E	113/113 (100%)	113 (100%)	0	100	100
1	G	112/113 (99%)	110 (98%)	2 (2%)	54	36
2	B	12/15 (80%)	11 (92%)	1 (8%)	9	2
2	D	13/15 (87%)	13 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	F	12/15 (80%)	9 (75%)	3 (25%)	0	0
2	H	11/15 (73%)	10 (91%)	1 (9%)	7	1
All	All	498/512 (97%)	487 (98%)	11 (2%)	50	28

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	THR
1	A	117	LYS
2	B	7	SER
1	C	21	LYS
1	C	100	SER
2	F	4	SER
2	F	18[A]	LYS
2	F	18[B]	LYS
1	G	10	THR
1	G	98	VAL
2	H	6	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	8	GLN
1	C	110	ASN
1	E	20	ASN
1	G	43	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

26 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	IPA	D	101	-	3,3,3	0.44	0	3,3,3	0.17	0
4	IPA	C	201	-	3,3,3	0.51	0	3,3,3	1.08	0
4	IPA	A	202	-	3,3,3	1.18	0	3,3,3	1.47	0
4	IPA	E	205	-	3,3,3	1.16	0	3,3,3	2.01	1 (33%)
4	IPA	E	204	-	3,3,3	0.64	0	3,3,3	0.35	0
4	IPA	H	101	-	3,3,3	0.51	0	3,3,3	0.34	0
3	MBG	E	201	-	13,13,13	1.21	1 (7%)	18,18,18	2.32	3 (16%)
4	IPA	H	102	-	3,3,3	0.63	0	3,3,3	0.14	0
5	EDO	E	206	-	3,3,3	0.56	0	2,2,2	0.20	0
4	IPA	A	203	-	3,3,3	0.36	0	3,3,3	0.43	0
4	IPA	G	203	-	3,3,3	0.54	0	3,3,3	0.28	0
4	IPA	C	204	-	3,3,3	0.48	0	3,3,3	0.65	0
4	IPA	C	203	-	3,3,3	0.40	0	3,3,3	0.35	0
3	MBG	A	201	-	13,13,13	1.05	1 (7%)	18,18,18	3.09	8 (44%)
5	EDO	A	206	-	3,3,3	0.70	0	2,2,2	0.55	0
3	MBG	G	201	-	13,13,13	1.00	1 (7%)	18,18,18	3.00	8 (44%)
4	IPA	D	102	-	3,3,3	0.50	0	3,3,3	0.66	0
4	IPA	E	202	-	3,3,3	0.66	0	3,3,3	0.95	0
4	IPA	E	203	-	3,3,3	0.49	0	3,3,3	0.98	0
5	EDO	C	205	-	3,3,3	0.86	0	2,2,2	0.60	0
4	IPA	A	204	-	3,3,3	0.55	0	3,3,3	0.33	0
4	IPA	A	205	-	3,3,3	0.64	0	3,3,3	0.64	0
4	IPA	G	202	-	3,3,3	0.59	0	3,3,3	0.99	0
5	EDO	E	207	-	3,3,3	0.58	0	2,2,2	0.28	0
5	EDO	D	103	-	3,3,3	0.31	0	2,2,2	1.27	0
4	IPA	C	202	-	3,3,3	0.69	0	3,3,3	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	206	-	-	0/1/1/1	-
3	MBG	G	201	-	-	0/4/24/24	0/1/1/1
5	EDO	D	103	-	-	1/1/1/1	-
5	EDO	C	205	-	-	0/1/1/1	-
3	MBG	E	201	-	-	0/4/24/24	0/1/1/1
5	EDO	E	207	-	-	1/1/1/1	-
5	EDO	E	206	-	-	0/1/1/1	-
3	MBG	A	201	-	-	0/4/24/24	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	201	MBG	C4-C5	2.72	1.58	1.53
3	A	201	MBG	C4-C5	2.62	1.58	1.53
3	G	201	MBG	O1-C1	-2.33	1.36	1.40

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	201	MBG	O1-C1-C2	7.38	116.65	108.14
3	G	201	MBG	C1-O5-C5	-6.80	100.43	113.72
3	A	201	MBG	C1-C2-C3	-6.33	96.69	110.01
3	G	201	MBG	O1-C1-C2	6.20	115.29	108.14
3	A	201	MBG	C1-O5-C5	-6.07	101.87	113.72
3	G	201	MBG	C1-C2-C3	-5.69	98.04	110.01
3	A	201	MBG	O1-C1-C2	5.22	114.17	108.14
3	A	201	MBG	O5-C1-O1	4.67	121.81	110.94
3	E	201	MBG	C1-O5-C5	-4.12	105.68	113.72
3	G	201	MBG	C6-C5-C4	3.87	122.51	113.02
3	A	201	MBG	C6-C5-C4	3.29	121.10	113.02
4	E	205	IPA	O2-C2-C1	-3.25	89.42	110.28
3	A	201	MBG	O5-C5-C6	3.14	114.22	106.44
3	E	201	MBG	C1-C2-C3	-2.95	103.81	110.01
3	G	201	MBG	C7-O1-C1	-2.73	109.11	113.26
3	A	201	MBG	C7-O1-C1	-2.58	109.34	113.26
3	G	201	MBG	O4-C4-C3	-2.48	104.53	110.38
3	G	201	MBG	O4-C4-C5	2.23	114.83	109.32
3	G	201	MBG	O2-C2-C3	-2.10	105.42	110.38
3	A	201	MBG	O3-C3-C2	-2.03	105.59	110.38

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	E	207	EDO	O1-C1-C2-O2
5	D	103	EDO	O1-C1-C2-O2

There are no ring outliers.

16 monomers are involved in 35 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	101	IPA	3	0
4	C	201	IPA	1	0
4	A	202	IPA	1	0
4	E	205	IPA	2	0
4	H	101	IPA	1	0
4	A	203	IPA	1	0
4	G	203	IPA	5	0
4	C	203	IPA	2	0
4	D	102	IPA	2	0
4	E	202	IPA	2	0
4	E	203	IPA	3	0
4	A	204	IPA	1	0
4	A	205	IPA	6	0
4	G	202	IPA	1	0
5	D	103	EDO	3	0
4	C	202	IPA	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	133/133 (100%)	-0.74	0	100 100	6, 10, 15, 22	0
1	C	133/133 (100%)	-0.62	0	100 100	6, 12, 22, 36	0
1	E	133/133 (100%)	-0.75	0	100 100	6, 10, 18, 29	0
1	G	133/133 (100%)	-0.70	0	100 100	6, 11, 19, 24	0
2	B	16/19 (84%)	-0.31	0	100 100	7, 13, 31, 35	0
2	D	16/19 (84%)	0.00	1 (6%)	27 29	8, 12, 43, 43	1 (6%)
2	F	16/19 (84%)	-0.44	0	100 100	6, 11, 23, 37	1 (6%)
2	H	15/19 (78%)	-0.12	1 (6%)	25 27	7, 12, 25, 48	0
All	All	595/608 (97%)	-0.65	2 (0%)	90 92	6, 11, 21, 48	2 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	17	ALA	5.1
2	H	18	LYS	2.7

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	IPA	E	203	4/4	0.71	0.20	13,17,19,29	0
4	IPA	D	102	4/4	0.74	0.16	21,24,27,33	0
4	IPA	C	202	4/4	0.76	0.17	31,33,41,43	0
4	IPA	A	205	4/4	0.78	0.17	19,28,28,32	0
4	IPA	H	102	4/4	0.78	0.15	39,40,43,47	0
4	IPA	G	202	4/4	0.80	0.17	22,29,31,32	0
4	IPA	C	201	4/4	0.82	0.17	22,29,32,37	0
4	IPA	E	205	4/4	0.85	0.12	10,10,11,11	0
4	IPA	D	101	4/4	0.86	0.17	29,32,32,33	0
4	IPA	A	202	4/4	0.86	0.15	14,17,18,20	0
5	EDO	D	103	4/4	0.86	0.13	32,34,36,38	0
4	IPA	A	204	4/4	0.87	0.16	34,35,36,37	0
4	IPA	E	204	4/4	0.87	0.11	24,27,28,32	0
4	IPA	A	203	4/4	0.88	0.12	30,31,34,41	0
4	IPA	E	202	4/4	0.89	0.10	13,16,17,18	0
4	IPA	G	203	4/4	0.89	0.10	22,32,32,34	0
5	EDO	E	206	4/4	0.89	0.11	29,32,37,44	0
4	IPA	H	101	4/4	0.91	0.10	33,33,33,34	0
4	IPA	C	203	4/4	0.91	0.09	23,23,23,28	0
5	EDO	E	207	4/4	0.91	0.09	21,24,26,35	0
4	IPA	C	204	4/4	0.93	0.09	25,27,28,29	0
5	EDO	A	206	4/4	0.94	0.08	22,24,24,25	0
3	MBG	E	201	13/13	0.95	0.07	10,16,22,28	0
3	MBG	G	201	13/13	0.96	0.06	11,13,18,19	0
3	MBG	A	201	13/13	0.96	0.07	9,16,24,28	0
5	EDO	C	205	4/4	0.96	0.06	13,13,14,16	0

6.5 Other polymers ⓘ

There are no such residues in this entry.